



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:16 AM GMT

PDB ID : 2GD2
Title : The 1,1-proton transfer reaction mechanism by alpha-methylacyl-CoA race-
mase is catalyzed by an aspartate/histidine pair and involves a smooth,
methionine-rich surface for binding the fatty acyl moiety
Authors : Bhaumik, P.; Wierenga, R.K.
Deposited on : 2006-03-15
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	<div> <div>10%</div> <div>87%</div> <div>11%</div> <div>..</div> </div>
1	B	360	<div> <div>9%</div> <div>81%</div> <div>15%</div> <div>..</div> </div>
1	C	360	<div> <div>9%</div> <div>86%</div> <div>11%</div> <div>..</div> </div>
1	D	360	<div> <div>9%</div> <div>86%</div> <div>11%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	2308	-	-	X	X
3	GOL	C	2307	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 12635 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called probable alpha-methylacyl-CoA racemase MCR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	6	0
			2701	1693	488	504	16			
1	B	354	Total	C	N	O	S	0	11	0
			2718	1702	492	508	16			
1	C	354	Total	C	N	O	S	0	11	0
			2724	1705	494	509	16			
1	D	354	Total	C	N	O	S	0	10	0
			2722	1704	495	507	16			

- Molecule 2 is ACETOACETYL-COENZYME A (three-letter code: CAA) (formula: $C_{25}H_{40}N_7O_{18}P_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			54	25	7	18	3 1		
2	B	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
2	D	1	Total	C	N	O	P	S	0	0
			54	25	7	18	3	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	394	Total	O	0	0
			394	394		
4	B	388	Total	O	0	0
			388	388		
4	C	372	Total	O	0	0
			372	372		
4	D	400	Total	O	0	0
			400	400		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: probable alpha-methylacyl-CoA racemase MCR




- Molecule 1: probable alpha-methylacyl-CoA racemase MCR




- Molecule 1: probable alpha-methylacyl-CoA racemase MCR



- Molecule 1: probable alpha-methylacyl-CoA racemase MCR





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	181.12Å 80.03Å 118.62Å 90.00° 91.58° 90.00°	Depositor
Resolution (Å)	19.80 – 1.70 19.80 – 1.70	Depositor EDS
% Data completeness (in resolution range)	91.3 (19.80-1.70) 91.3 (19.80-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.208 , 0.250 0.209 , 0.247	Depositor DCC
R_{free} test set	8538 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	15.5	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 55.0	EDS
Estimated twinning fraction	0.166 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 169869 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12635	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.78 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2435e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CAA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/2792	0.93	8/3795 (0.2%)
1	B	0.84	0/2818	0.97	12/3830 (0.3%)
1	C	0.85	0/2822	0.92	7/3835 (0.2%)
1	D	0.88	0/2812	0.93	9/3822 (0.2%)
All	All	0.85	0/11244	0.94	36/15282 (0.2%)

There are no bond length outliers.

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	85	ARG	NE-CZ-NH2	-10.66	114.97	120.30
1	A	185	ASP	CB-CG-OD2	10.37	127.63	118.30
1	A	295	ASP	CB-CG-OD2	9.06	126.45	118.30
1	B	295	ASP	CB-CG-OD2	8.76	126.19	118.30
1	C	185	ASP	CB-CG-OD2	8.41	125.86	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2701	0	2641	31	0
1	B	2718	0	2651	52	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2724	0	2662	43	0
1	D	2722	0	2660	35	0
2	A	54	0	36	3	0
2	B	48	0	31	8	0
2	C	48	0	31	5	0
2	D	54	0	36	3	0
3	B	6	0	8	11	0
3	C	6	0	8	4	0
4	A	394	0	0	14	0
4	B	388	0	0	6	0
4	C	372	0	0	13	1
4	D	400	0	0	10	1
All	All	12635	0	10764	151	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 151 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:2:ALA:HB1	4:D:2014:HOH:O	1.56	1.05
2:B:1752:CAA:S1P	3:B:2308:GOL:H31	2.12	0.89
1:C:322[B]:ALA:O	1:C:323[B]:ASN:O	1.94	0.86
1:C:176:GLN:HG3	4:D:1997:HOH:O	1.80	0.82
1:D:38:ARG:HG3	1:D:38:ARG:HH11	1.46	0.79

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:2554:HOH:O	4:D:1998:HOH:O[4_656]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	356/360 (99%)	345 (97%)	11 (3%)	0	100	100
1	B	361/360 (100%)	347 (96%)	11 (3%)	3 (1%)	24	7
1	C	361/360 (100%)	347 (96%)	12 (3%)	2 (1%)	30	12
1	D	360/360 (100%)	345 (96%)	15 (4%)	0	100	100
All	All	1438/1440 (100%)	1384 (96%)	49 (3%)	5 (0%)	52	26

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	323[A]	ASN
1	C	323[B]	ASN
1	B	323[A]	ASN
1	B	323[B]	ASN
1	B	151	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/275 (100%)	272 (99%)	4 (1%)	74	59
1	B	277/275 (101%)	270 (98%)	7 (2%)	55	34
1	C	278/275 (101%)	270 (97%)	8 (3%)	50	27
1	D	276/275 (100%)	268 (97%)	8 (3%)	50	27
All	All	1107/1100 (101%)	1080 (98%)	27 (2%)	60	36

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	58	THR
1	C	123[B]	GLN
1	D	203[B]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	103	ASN
1	B	85	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	263	ASN
1	C	21	HIS
1	D	103	ASN
1	B	134	ASN
1	D	134	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CAA	A	1751	1	45,56,56	1.20	3 (6%)	58,83,83	3.04	11 (18%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CAA	B	1752	-	40,50,56	1.09	3 (7%)	50,75,83	2.05	12 (24%)
3	GOL	B	2308	-	5,5,5	0.57	0	5,5,5	0.66	0
2	CAA	C	1753	1	40,50,56	1.09	4 (10%)	50,75,83	2.14	12 (24%)
3	GOL	C	2307	-	5,5,5	0.43	0	5,5,5	0.62	0
2	CAA	D	1754	-	45,56,56	1.22	6 (13%)	58,83,83	3.00	13 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CAA	A	1751	1	-	0/50/71/71	0/3/3/3
2	CAA	B	1752	-	-	0/44/64/71	0/3/3/3
3	GOL	B	2308	-	-	0/4/4/4	0/0/0/0
2	CAA	C	1753	1	-	0/44/64/71	0/3/3/3
3	GOL	C	2307	-	-	0/4/4/4	0/0/0/0
2	CAA	D	1754	-	-	2/50/71/71	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1751	CAA	C1-S1P	-3.46	1.69	1.76
2	D	1754	CAA	C1-S1P	-3.11	1.69	1.76
2	B	1752	CAA	C8A-N7A	2.00	1.38	1.34
2	C	1753	CAA	P2A-O4A	2.03	1.58	1.51
2	D	1754	CAA	O4B-C1B	2.06	1.43	1.41

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1751	CAA	O1-C1-S1P	-10.47	114.52	122.83
2	B	1752	CAA	N3A-C2A-N1A	-7.94	122.82	128.89
2	C	1753	CAA	N3A-C2A-N1A	-7.59	123.08	128.89
2	A	1751	CAA	N3A-C2A-N1A	-6.89	123.62	128.89
2	D	1754	CAA	C1B-N9A-C4A	-6.41	117.27	126.94

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1754	CAA	C4-C3-C2-C1
2	D	1754	CAA	O3-C3-C2-C1

There are no ring outliers.

6 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1751	CAA	3	0
2	B	1752	CAA	8	0
3	B	2308	GOL	11	0
2	C	1753	CAA	5	0
3	C	2307	GOL	4	0
2	D	1754	CAA	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/360 (98%)	0.67	35 (9%) 9 10	17, 25, 36, 46	0
1	B	354/360 (98%)	0.74	31 (8%) 12 13	19, 26, 38, 47	0
1	C	354/360 (98%)	0.69	33 (9%) 11 12	17, 23, 36, 49	0
1	D	354/360 (98%)	0.62	33 (9%) 11 12	15, 24, 36, 44	0
All	All	1416/1440 (98%)	0.68	132 (9%) 11 12	15, 24, 37, 49	0

The worst 5 of 132 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	GLY	7.8
1	C	347	ALA	7.4
1	C	45	ILE	6.8
1	C	349	ILE	6.4
1	C	359	ASP	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	B	2308	6/6	0.70	0.26	5.09	40,43,45,46	0
3	GOL	C	2307	6/6	0.85	0.15	1.30	42,44,45,46	0
2	CAA	A	1751	54/54	0.94	0.13	0.10	19,28,33,39	1
2	CAA	D	1754	54/54	0.93	0.14	0.07	19,28,35,42	1
2	CAA	B	1752	48/54	0.94	0.14	0.01	20,28,36,37	0
2	CAA	C	1753	48/54	0.94	0.12	-0.16	16,26,34,37	0

6.5 Other polymers

There are no such residues in this entry.