



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:51 PM GMT

PDB ID : 3GKA
Title : Crystal structure of N-ethylmaleimidine reductase from Burkholderia pseudo-mallei
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2009-03-10
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	361	 90% 6% . .
1	B	361	 90% 6% . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5735 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-ethylmaleimide reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	350	Total	C	N	O	S	0	1	0
			2626	1661	476	486	3			
1	B	351	Total	C	N	O	S	0	1	0
			2650	1674	483	489	4			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP Q63LY5
A	2	ALA	-	EXPRESSION TAG	UNP Q63LY5
A	3	HIS	-	EXPRESSION TAG	UNP Q63LY5
A	4	HIS	-	EXPRESSION TAG	UNP Q63LY5
A	5	HIS	-	EXPRESSION TAG	UNP Q63LY5
A	6	HIS	-	EXPRESSION TAG	UNP Q63LY5
A	7	HIS	-	EXPRESSION TAG	UNP Q63LY5
A	8	HIS	-	EXPRESSION TAG	UNP Q63LY5
B	1	MET	-	EXPRESSION TAG	UNP Q63LY5
B	2	ALA	-	EXPRESSION TAG	UNP Q63LY5
B	3	HIS	-	EXPRESSION TAG	UNP Q63LY5
B	4	HIS	-	EXPRESSION TAG	UNP Q63LY5
B	5	HIS	-	EXPRESSION TAG	UNP Q63LY5
B	6	HIS	-	EXPRESSION TAG	UNP Q63LY5
B	7	HIS	-	EXPRESSION TAG	UNP Q63LY5
B	8	HIS	-	EXPRESSION TAG	UNP Q63LY5

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 17	N 4	O 9	P 1	0	0
2	B	1	Total 31	C 17	N 4	O 9	P 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	184	Total O 184 184	0	0
3	B	213	Total O 213 213	0	0

- Molecule 1: N-ethylmaleimide reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.35Å 77.89Å 168.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 19.47 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-2.30) 99.9 (19.47-2.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.70 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.176 , 0.239 0.177 , 0.232	Depositor DCC
R_{free} test set	1506 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.090	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 29686 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5735	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2691	0.56	0/3675
1	B	0.43	0/2715	0.57	0/3704
All	All	0.43	0/5406	0.57	0/7379

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2626	0	2548	16	0
1	B	2650	0	2583	12	0
2	A	31	0	19	1	0
2	B	31	0	19	0	0
3	A	184	0	0	1	0
3	B	213	0	0	0	0
All	All	5735	0	5169	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:64:ALA:HB1	1:B:108:TRP:CD2	2.29	0.67
1:A:64:ALA:HB1	1:A:108:TRP:CD2	2.36	0.60
1:B:126:ALA:HB1	1:B:127:PRO:HD2	1.92	0.52
1:A:102:ARG:NH1	3:A:444:HOH:O	2.42	0.52
1:B:112:ARG:HB3	1:B:126:ALA:HB2	1.92	0.51
1:A:210:ARG:NH1	1:A:245:ASP:OD1	2.42	0.51
1:A:183:ALA:CB	1:A:233[B]:HIS:HD2	2.23	0.51
1:B:292:VAL:CG2	1:B:313:VAL:HG22	2.41	0.51
1:B:115:ASP:OD2	1:B:117:VAL:HG22	2.13	0.48
1:A:139:VAL:O	1:A:139:VAL:HG23	2.13	0.48
1:A:164:ARG:HG3	1:A:220:ALA:HB1	1.95	0.47
1:A:183:ALA:HB3	1:A:233[B]:HIS:HD2	1.79	0.46
1:B:268:PHE:C	1:B:268:PHE:CD2	2.89	0.45
1:A:80:ILE:O	1:A:80:ILE:HG13	2.17	0.45
1:B:257:ARG:HB3	1:B:261:ARG:HH22	1.81	0.45
1:A:280:GLN:OE1	1:A:309:GLN:HA	2.17	0.44
1:A:344:TYR:CZ	2:A:362:FMN:HM72	2.53	0.44
1:A:293:ASN:O	1:A:294:GLU:CB	2.66	0.44
1:B:140:ARG:HA	1:B:141:PRO:C	2.39	0.43
1:A:236:PRO:HB3	1:A:278:ILE:HD11	1.99	0.43
1:B:64:ALA:HB1	1:B:108:TRP:CG	2.54	0.42
1:A:139:VAL:HG21	1:A:143:ARG:NH1	2.34	0.42
1:A:13:PHE:CE1	1:A:330:LYS:HG2	2.55	0.41
1:B:152:GLU:HG3	1:B:155:GLU:OE2	2.19	0.41
1:A:301:ALA:HB1	1:A:313:VAL:HG21	2.02	0.41
1:B:199:ARG:HB2	1:B:204:GLY:HA3	2.03	0.40
1:A:285:ALA:HA	1:B:285:ALA:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/361 (97%)	337 (97%)	11 (3%)	1 (0%)	46	57
1	B	350/361 (97%)	337 (96%)	12 (3%)	1 (0%)	46	57
All	All	699/722 (97%)	674 (96%)	23 (3%)	2 (0%)	46	57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	294	GLU
1	B	294	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/267 (96%)	253 (99%)	2 (1%)	86	94
1	B	259/267 (97%)	252 (97%)	7 (3%)	52	70
All	All	514/534 (96%)	505 (98%)	9 (2%)	66	82

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	164	ARG
1	A	268	PHE
1	B	68	THR
1	B	90	ARG
1	B	140	ARG
1	B	143	ARG
1	B	165	ARG
1	B	268	PHE
1	B	327	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FMN	A	362	-	32,33,33	1.17	2 (6%)	34,50,50	1.65	5 (14%)
2	FMN	B	362	-	32,33,33	1.18	3 (9%)	34,50,50	1.65	5 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	362	-	-	0/18/18/18	0/3/3/3
2	FMN	B	362	-	-	0/18/18/18	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	362	FMN	C5A-N5	2.02	1.38	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	362	FMN	C4-N3	3.01	1.38	1.33
2	A	362	FMN	C4-N3	3.01	1.38	1.33
2	A	362	FMN	C4A-N5	3.33	1.38	1.33
2	B	362	FMN	C4A-N5	3.33	1.38	1.33

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	362	FMN	N3-C2-N1	-4.46	120.17	127.69
2	A	362	FMN	N3-C2-N1	-4.46	120.18	127.69
2	A	362	FMN	C4A-C4-N3	-2.83	119.82	123.52
2	B	362	FMN	C4A-C4-N3	-2.83	119.82	123.52
2	B	362	FMN	C4A-N5-C5A	2.75	119.97	116.72
2	A	362	FMN	C4A-N5-C5A	2.77	119.98	116.72
2	A	362	FMN	C5A-C9A-N10	3.24	120.01	117.58
2	B	362	FMN	C5A-C9A-N10	3.30	120.05	117.58
2	B	362	FMN	C4-N3-C2	5.98	120.15	115.16
2	A	362	FMN	C4-N3-C2	6.01	120.17	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	362	FMN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/361 (96%)	-0.36	1 (0%) 94 96	7, 13, 22, 29	0
1	B	351/361 (97%)	-0.36	3 (0%) 85 89	5, 12, 25, 37	0
All	All	701/722 (97%)	-0.36	4 (0%) 90 93	5, 13, 23, 37	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	9	MET	3.0
1	A	275	GLY	2.8
1	B	140	ARG	2.6
1	B	306	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FMN	B	362	31/31	0.97	0.10	-0.46	7,9,11,11	0
2	FMN	A	362	31/31	0.95	0.10	-0.62	7,9,11,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.