



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 18, 2016 – 01:29 PM EDT

PDB ID : 5GMC  
Title : Methylation at position 32 of tRNA catalyzed by TrmJ alters oxidative stress response in *Pseudomonas aeruginosa*  
Authors : Jaroensuk, J.; Atichartpongkul, S.; Chionh, Y.H.; Wong, Y.H.; Liew, C.W.; McBee, M.E.; Thongdee, N.; Prestwich, E.G.; DeMott, M.S.; Mongkolsuk, S.; Dedon, P.C.; Lescar, J.; Fuangthong, M.  
Deposited on : 2016-07-13  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	174	<div> <div>6%</div> <div> <div></div> <div>89%</div> <div>11%</div> <div>.</div> </div> </div>
1	B	174	<div> <div>6%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>..</div> </div> </div>
1	C	174	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div>.</div> </div> </div>
1	D	174	<div> <div>5%</div> <div> <div></div> <div>87%</div> <div>11%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA (cytidine/uridine-2'-O-)-methyltransferase TrmJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	174	Total	C	N	O	S	0	4	0
			1347	841	240	256	10			
1	B	171	Total	C	N	O	S	0	1	0
			1301	811	235	245	10			
1	C	169	Total	C	N	O	S	0	3	0
			1295	804	235	246	10			
1	D	172	Total	C	N	O	S	0	2	0
			1324	826	238	251	9			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	ASN	-	expression tag	UNP A0A072ZPM2
A	-5	LEU	-	expression tag	UNP A0A072ZPM2
A	-4	TYR	-	expression tag	UNP A0A072ZPM2
A	-3	PHE	-	expression tag	UNP A0A072ZPM2
A	-2	GLN	-	expression tag	UNP A0A072ZPM2
A	-1	SER	-	expression tag	UNP A0A072ZPM2
A	0	MET	-	expression tag	UNP A0A072ZPM2
B	-6	ASN	-	expression tag	UNP A0A072ZPM2
B	-5	LEU	-	expression tag	UNP A0A072ZPM2
B	-4	TYR	-	expression tag	UNP A0A072ZPM2
B	-3	PHE	-	expression tag	UNP A0A072ZPM2
B	-2	GLN	-	expression tag	UNP A0A072ZPM2
B	-1	SER	-	expression tag	UNP A0A072ZPM2
B	0	MET	-	expression tag	UNP A0A072ZPM2
C	-6	ASN	-	expression tag	UNP A0A072ZPM2
C	-5	LEU	-	expression tag	UNP A0A072ZPM2
C	-4	TYR	-	expression tag	UNP A0A072ZPM2
C	-3	PHE	-	expression tag	UNP A0A072ZPM2
C	-2	GLN	-	expression tag	UNP A0A072ZPM2
C	-1	SER	-	expression tag	UNP A0A072ZPM2
C	0	MET	-	expression tag	UNP A0A072ZPM2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-6	ASN	-	expression tag	UNP A0A072ZPM2
D	-5	LEU	-	expression tag	UNP A0A072ZPM2
D	-4	TYR	-	expression tag	UNP A0A072ZPM2
D	-3	PHE	-	expression tag	UNP A0A072ZPM2
D	-2	GLN	-	expression tag	UNP A0A072ZPM2
D	-1	SER	-	expression tag	UNP A0A072ZPM2
D	0	MET	-	expression tag	UNP A0A072ZPM2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	158	Total	O	0	0
			158	158		
2	B	163	Total	O	0	0
			163	163		
2	C	172	Total	O	0	0
			172	172		
2	D	136	Total	O	0	0
			136	136		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: tRNA (cytidine/uridine-2'-O-)-methyltransferase TrmJ




- Molecule 1: tRNA (cytidine/uridine-2'-O-)-methyltransferase TrmJ



- Molecule 1: tRNA (cytidine/uridine-2'-O-)-methyltransferase TrmJ



- Molecule 1: tRNA (cytidine/uridine-2'-O-)-methyltransferase TrmJ



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.44Å 65.04Å 93.19Å 90.00° 110.42° 90.00°	Depositor
Resolution (Å)	24.27 – 1.70 24.27 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.4 (24.27-1.70) 93.7 (24.27-1.70)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.03 (at 1.71Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, $R_{free}$	0.169 , 0.193 0.179 , 0.194	Depositor DCC
$R_{free}$ test set	3782 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.7	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 55.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.024 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5896	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.72	1/1374 (0.1%)	0.72	0/1868
1	B	0.76	0/1324	0.71	2/1799 (0.1%)
1	C	0.79	0/1317	0.72	0/1790
1	D	0.85	0/1348	0.77	0/1832
All	All	0.78	1/5363 (0.0%)	0.73	2/7289 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	51	SER	CB-OG	-6.07	1.34	1.42

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	87	PRO	N-CA-C	5.64	126.76	112.10
1	B	88	TRP	CA-CB-CG	5.36	123.88	113.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	27[B]	ASN	Mainchain

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	D	27[A]	ASN	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1347	0	1327	14	0
1	B	1301	0	1292	18	0
1	C	1295	0	1284	9	0
1	D	1324	0	1307	16	0
2	A	158	0	0	3	0
2	B	163	0	0	6	0
2	C	172	0	0	3	0
2	D	136	0	0	0	0
All	All	5896	0	5210	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:TRP:CE3	2:B:208:HOH:O	1.70	1.30
1:B:88:TRP:NE1	2:B:201:HOH:O	1.94	0.99
1:B:86:ILE:HB	1:B:87:PRO:HD3	1.49	0.93
1:C:117:GLU:O	2:C:203:HOH:O	1.91	0.88
1:C:43:HIS:HD2	1:C:45:ASP:H	1.26	0.84
1:D:43:HIS:HD2	1:D:45:ASP:H	1.30	0.80
1:A:43:HIS:HD2	1:A:45:ASP:H	1.27	0.79
1:B:88:TRP:CZ3	2:B:208:HOH:O	2.12	0.77
1:B:88:TRP:CE2	2:B:201:HOH:O	2.32	0.77
1:D:11:ASN:HD22	1:D:116:ARG:HE	1.35	0.74
1:B:86:ILE:HB	1:B:87:PRO:CD	2.18	0.73
1:A:109:GLU:OE1	2:A:201:HOH:O	2.06	0.73
1:B:88:TRP:CD1	1:B:132:HIS:NE2	2.57	0.72
1:D:101:LEU:O	1:D:105:GLU:HG3	1.93	0.69

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:11:ASN:HD22	1:C:116:ARG:HE	1.39	0.68
1:B:124:GLU:HG3	2:B:283:HOH:O	1.93	0.68
1:C:167:LYS:CE	2:C:201:HOH:O	2.26	0.67
1:A:123:ASN:O	1:A:127:GLN:HG3	1.95	0.65
1:B:88:TRP:CD1	1:B:132:HIS:CD2	2.84	0.65
1:A:43:HIS:CD2	1:A:45:ASP:H	2.13	0.65
1:B:11:ASN:HD22	1:B:116:ARG:HE	1.44	0.65
1:D:43:HIS:CD2	1:D:45:ASP:H	2.13	0.65
1:C:167:LYS:NZ	2:C:201:HOH:O	1.60	0.63
1:D:67[B]:GLU:CG	1:D:128:ARG:CZ	2.77	0.63
1:A:145:ASN:HD22	1:A:148:ALA:H	1.47	0.62
1:D:67[B]:GLU:HG3	1:D:128:ARG:CZ	2.31	0.61
1:A:159[B]:MET:HE2	2:B:333:HOH:O	2.01	0.59
1:C:43:HIS:CD2	1:C:45:ASP:H	2.13	0.58
1:A:11[B]:ASN:ND2	2:A:203:HOH:O	2.32	0.55
1:B:86:ILE:CB	1:B:87:PRO:HD3	2.30	0.55
1:D:26:LYS:HB2	1:D:56:ILE:HD12	1.90	0.54
1:B:88:TRP:HD1	1:B:132:HIS:CD2	2.24	0.54
1:B:167:LYS:HE3	1:B:167:LYS:HA	1.92	0.52
1:B:88:TRP:HB3	1:B:89:PRO:HD2	1.91	0.52
1:D:-2:GLN:NE2	1:D:165:GLN:HG2	2.25	0.51
1:B:166:GLY:O	1:B:167:LYS:HE3	2.09	0.51
1:A:165:GLN:HB3	1:A:167:LYS:HE3	1.93	0.51
1:B:11:ASN:HD22	1:B:116:ARG:NE	2.09	0.51
1:A:-6:ASN:O	1:A:-3:PHE:N	2.45	0.50
1:D:67[B]:GLU:HG2	1:D:128:ARG:CZ	2.42	0.50
1:C:11:ASN:HD22	1:C:116:ARG:NE	2.08	0.50
1:A:11[A]:ASN:ND2	2:A:207:HOH:O	2.45	0.49
1:A:-6:ASN:O	1:A:-5:LEU:C	2.48	0.48
1:D:167:LYS:HE3	1:D:167:LYS:HB3	1.64	0.45
1:D:26:LYS:HB2	1:D:56:ILE:CD1	2.46	0.45
1:B:88:TRP:HD1	1:B:132:HIS:NE2	2.09	0.45
1:D:11:ASN:HD22	1:D:116:ARG:NE	2.10	0.44
1:C:-1:SER:HA	1:C:161:TRP:HZ2	1.82	0.44
1:A:11[A]:ASN:HB3	1:A:116:ARG:HG2	2.01	0.43
1:D:155:TYR:O	1:D:159:MET:HG2	2.19	0.43
1:D:132:HIS:N	1:D:132:HIS:CD2	2.87	0.42
1:A:-6:ASN:C	1:A:-4:TYR:N	2.72	0.42
1:A:155:TYR:O	1:A:159[B]:MET:HG2	2.20	0.41
1:C:100[A]:CYS:SG	1:C:112:LEU:HD21	2.60	0.41
1:D:-3:PHE:CE2	1:D:1:MET:HE2	2.56	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100[B]:CYS:SG	1:B:112:LEU:HD21	2.61	0.41
1:D:-3:PHE:HE2	1:D:1:MET:HE2	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	176/174 (101%)	170 (97%)	6 (3%)	0	100	100
1	B	170/174 (98%)	164 (96%)	5 (3%)	1 (1%)	30	12
1	C	170/174 (98%)	166 (98%)	4 (2%)	0	100	100
1	D	172/174 (99%)	169 (98%)	3 (2%)	0	100	100
All	All	688/696 (99%)	669 (97%)	18 (3%)	1 (0%)	56	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	86	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	143/140 (102%)	140 (98%)	3 (2%)	61	42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	138/140 (99%)	137 (99%)	1 (1%)	88	82
1	C	138/140 (99%)	135 (98%)	3 (2%)	60	39
1	D	140/140 (100%)	137 (98%)	3 (2%)	61	42
All	All	559/560 (100%)	549 (98%)	10 (2%)	66	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-6	ASN
1	A	0	MET
1	A	61	ARG
1	B	167	LYS
1	C	84	ARG
1	C	85	ARG
1	C	90	LEU
1	D	-4	TYR
1	D	109	GLU
1	D	167	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-6	ASN
1	A	14	HIS
1	A	43	HIS
1	A	145	ASN
1	A	151	GLN
1	B	11	ASN
1	B	27	ASN
1	B	151	GLN
1	C	11	ASN
1	C	43	HIS
1	C	151	GLN
1	D	-2	GLN
1	D	11	ASN
1	D	43	HIS
1	D	127	GLN
1	D	151	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	174/174 (100%)	0.21	10 (5%) 27 29	13, 22, 42, 56	0
1	B	171/174 (98%)	0.22	11 (6%) 23 25	12, 19, 47, 95	0
1	C	169/174 (97%)	0.16	9 (5%) 30 32	14, 20, 51, 101	0
1	D	172/174 (98%)	0.03	8 (4%) 35 39	14, 25, 46, 61	0
All	All	686/696 (98%)	0.15	38 (5%) 29 31	12, 21, 46, 101	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	86	ILE	13.7
1	A	-5	LEU	12.3
1	B	88	TRP	12.0
1	D	-4	TYR	12.0
1	C	85	ARG	10.9
1	C	88	TRP	10.8
1	B	87	PRO	9.6
1	C	86	ILE	8.9
1	C	87	PRO	8.6
1	B	-3	PHE	8.2
1	B	85	ARG	7.6
1	A	-6	ASN	7.6
1	C	84	ARG	7.4
1	A	-4	TYR	6.8
1	A	167	LYS	5.5
1	C	83	ASP	5.1
1	D	-3	PHE	4.6
1	A	166	GLY	4.5
1	B	84	ARG	4.4
1	C	167	LYS	4.3
1	D	167	LYS	4.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	89	PRO	4.3
1	B	83	ASP	4.1
1	A	118	TYR	3.6
1	A	119	ALA	3.5
1	B	167	LYS	3.4
1	C	0	MET	3.2
1	D	83	ASP	3.1
1	D	166	GLY	2.7
1	D	82	ARG	2.6
1	B	0	MET	2.4
1	A	-3	PHE	2.3
1	D	-2	GLN	2.3
1	A	117[A]	GLU	2.2
1	A	83	ASP	2.2
1	D	0	MET	2.2
1	B	89	PRO	2.2
1	B	139	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.