



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:57 PM GMT

PDB ID : 4GN1
Title : Crystal Structure of the RA and PH domains of Lamellipodin
Authors : Chang, Y.C.E.; Wu, J.
Deposited on : 2012-08-16
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	<div> <div>4%</div> <div>82% 15% ..</div> </div>
1	B	255	<div> <div>5%</div> <div>79% 16% ..</div> </div>
1	C	255	<div> <div>3%</div> <div>78% 13% • 6%</div> </div>
1	D	255	<div> <div>5%</div> <div>77% 15% • 6%</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8481 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ras-associated and pleckstrin homology domains-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	1	0
			2091	1334	359	382	16			
1	B	250	Total	C	N	O	S	0	4	0
			2099	1341	358	382	18			
1	C	240	Total	C	N	O	S	0	1	0
			2011	1284	342	369	16			
1	D	240	Total	C	N	O	S	0	0	0
			2003	1279	341	368	15			

- Molecule 2 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	3	4		
2	B	1	Total	C	O	0	0
			7	3	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			7	3	4		
2	D	1	Total	C	O	0	0
			7	3	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	62	Total	O	0	0
			62	62		
3	B	59	Total	O	0	0
			59	59		
3	C	68	Total	O	0	0
			68	68		
3	D	60	Total	O	0	0
			60	60		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ras-associated and pleckstrin homology domains-containing protein 1




- Molecule 1: Ras-associated and pleckstrin homology domains-containing protein 1




- Molecule 1: Ras-associated and pleckstrin homology domains-containing protein 1



- Molecule 1: Ras-associated and pleckstrin homology domains-containing protein 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.34Å 93.47Å 80.72Å 90.00° 104.91° 90.00°	Depositor
Resolution (Å)	46.73 – 2.40 46.73 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.8 (46.73-2.40) 98.8 (46.73-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.61 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.253 , 0.303 0.253 , 0.303	Depositor DCC
R_{free} test set	2188 reflections (5.50%)	DCC
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	10 of 41923 reflections (0.024%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8481	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 82.82 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.5340e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	2/2136 (0.1%)	0.72	0/2874
1	B	0.62	2/2150 (0.1%)	0.72	0/2893
1	C	0.64	2/2052 (0.1%)	0.68	0/2763
1	D	0.65	4/2044 (0.2%)	0.72	1/2753 (0.0%)
All	All	0.63	10/8382 (0.1%)	0.71	1/11283 (0.0%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	404	TRP	CD2-CE2	5.75	1.48	1.41
1	B	413	TRP	CD2-CE2	5.67	1.48	1.41
1	C	496	TRP	CD2-CE2	5.58	1.48	1.41
1	D	312	TRP	CD2-CE2	5.57	1.48	1.41
1	D	496	TRP	CD2-CE2	5.30	1.47	1.41
1	D	413	TRP	CD2-CE2	5.25	1.47	1.41
1	B	404	TRP	CD2-CE2	5.24	1.47	1.41
1	C	340	TRP	CD2-CE2	5.23	1.47	1.41
1	D	404	TRP	CD2-CE2	5.19	1.47	1.41
1	A	312	TRP	CD2-CE2	5.14	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	468	LEU	CA-CB-CG	5.85	128.76	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2091	0	2105	24	0
1	B	2099	0	2118	31	0
1	C	2011	0	2005	21	0
1	D	2003	0	1997	23	0
2	A	7	0	2	0	0
2	B	7	0	2	0	0
2	C	7	0	2	1	0
2	D	7	0	2	0	0
3	A	62	0	0	2	0
3	B	59	0	0	0	0
3	C	68	0	0	0	0
3	D	60	0	0	1	0
All	All	8481	0	8233	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:400:GLU:HG3	1:B:417:TYR:OH	1.59	1.00
1:D:443:LEU:HD13	3:D:760:HOH:O	1.67	0.94
1:B:472:HIS:HD2	1:B:474:GLN:H	1.15	0.89
1:B:370:LYS:HE3	1:B:372:GLU:HB3	1.56	0.86
1:A:363:PRO:HD2	1:A:381:LYS:HG2	1.69	0.74
1:A:321:LEU:HB3	1:A:323:MET:HE2	1.69	0.73
1:D:468:LEU:HD13	1:D:485:LEU:HB2	1.71	0.72
1:B:472:HIS:CD2	1:B:474:GLN:H	2.04	0.71
1:A:356:LYS:HE3	1:A:357:TYR:CE2	2.27	0.69
1:A:367:LEU:O	1:A:397:PRO:HG2	1.91	0.69
1:C:279:ASP:OD1	1:C:281:SER:HB3	1.93	0.69
1:B:404:TRP:HB2	1:B:486:CYS:HB3	1.76	0.68
1:B:402:VAL:O	1:B:403:LEU:HD12	1.96	0.66
1:A:280:ASP:HB2	1:B:290:ARG:HH21	1.61	0.66
1:D:427:TYR:O	1:D:440:VAL:HB	1.96	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:321:LEU:HB3	1:A:323:MET:CE	2.29	0.63
1:B:285[A]:MET:SD	1:B:300:LEU:HD13	2.39	0.62
1:A:280:ASP:HB2	1:B:290:ARG:NH2	2.15	0.61
1:A:404:TRP:HB2	1:A:486:CYS:HB3	1.82	0.61
1:B:356:LYS:HE3	1:B:357:TYR:CE2	2.36	0.60
1:D:416:ARG:HG2	1:D:427:TYR:HD1	1.65	0.60
1:D:472:HIS:HE1	1:D:474:GLN:OE1	1.85	0.60
1:B:323:MET:CE	1:B:469:VAL:HG11	2.32	0.59
1:A:330:HIS:HD2	1:A:512:ASN:HD21	1.50	0.58
1:C:323:MET:HE1	1:C:469:VAL:HG21	1.86	0.58
1:B:323:MET:HE1	1:B:469:VAL:HG11	1.84	0.58
1:B:404:TRP:CD2	1:B:463:PRO:HG2	2.40	0.57
1:A:330:HIS:CD2	1:A:512:ASN:HD21	2.23	0.57
1:C:323:MET:CE	1:C:469:VAL:HG11	2.35	0.57
1:C:472:HIS:HE1	1:C:474:GLN:OE1	1.88	0.56
1:C:319:SER:CB	1:C:345:GLN:HB2	2.36	0.56
1:D:299:ASN:HB3	1:D:303:LYS:HZ2	1.70	0.56
1:B:404:TRP:CG	1:B:463:PRO:HG2	2.41	0.54
1:C:415:LYS:O	1:C:416:ARG:HD3	2.07	0.54
1:D:382:GLU:O	1:D:386:GLU:HG2	2.08	0.53
1:A:363:PRO:HB3	1:A:385:LEU:HD21	1.91	0.53
1:C:274:ARG:NH1	1:C:282:SER:OG	2.43	0.52
1:C:323:MET:HE1	1:C:469:VAL:HG11	1.92	0.51
1:B:318[A]:VAL:HG12	1:B:323:MET:HG3	1.92	0.51
1:C:318:VAL:HG22	1:C:321:LEU:HB2	1.92	0.51
1:D:416:ARG:HG2	1:D:427:TYR:CD1	2.44	0.51
1:B:363:PRO:HD2	1:B:381:LYS:HG2	1.93	0.51
1:B:320:GLU:HG2	1:B:477:LYS:HA	1.93	0.51
1:A:369:GLY:HA2	1:A:370:LYS:C	2.31	0.50
1:C:319:SER:OG	1:C:345:GLN:HB2	2.11	0.50
1:B:428:VAL:HG11	1:B:435:VAL:HB	1.94	0.49
1:B:472:HIS:HD2	1:B:474:GLN:N	1.97	0.49
1:D:394:VAL:HG21	1:D:506:GLY:HA2	1.95	0.48
1:D:394:VAL:HG21	1:D:506:GLY:CA	2.43	0.48
1:C:427:TYR:CD2	1:C:442:PHE:HB2	2.49	0.48
1:A:404:TRP:CD2	1:A:463:PRO:HG3	2.49	0.48
1:C:359:LEU:HA	1:C:366:TYR:CD1	2.48	0.48
1:C:285[B]:MET:SD	1:C:300:LEU:HG	2.55	0.47
1:A:323:MET:HE1	1:A:469:VAL:HG11	1.96	0.47
1:B:402:VAL:HG12	1:B:415:LYS:HD2	1.95	0.47
1:A:357:TYR:O	1:A:361:LYS:HG3	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:406:LYS:HG2	1:D:407:ASP:O	2.14	0.46
1:C:451:TYR:OH	2:C:601:MLI:O6	2.23	0.46
1:B:468:LEU:HD23	1:B:468:LEU:H	1.79	0.46
1:A:359:LEU:HB3	3:A:739:HOH:O	2.15	0.46
1:B:318[A]:VAL:O	1:B:318[A]:VAL:HG13	2.16	0.46
1:C:445:LEU:HD13	1:C:500:ILE:HG23	1.98	0.46
1:B:318[A]:VAL:HG13	1:B:321:LEU:HB2	1.98	0.45
1:D:319:SER:OG	1:D:345:GLN:HB2	2.17	0.45
1:B:359:LEU:HA	1:B:366:TYR:CD1	2.52	0.45
1:B:370:LYS:CE	1:B:372:GLU:HB3	2.38	0.45
1:D:442:PHE:CZ	1:D:483:LYS:HG2	2.50	0.45
1:D:404:TRP:CE3	1:D:415:LYS:HG3	2.52	0.45
1:A:323:MET:CE	1:A:469:VAL:HG11	2.46	0.45
1:D:310:LEU:H	1:D:310:LEU:HD23	1.80	0.45
1:B:411:LYS:HE2	1:B:459:LYS:O	2.17	0.44
1:D:356:LYS:HE3	1:D:357:TYR:CE2	2.53	0.44
1:D:406:LYS:HB3	1:D:484:TYR:HB2	1.99	0.44
1:D:359:LEU:HD12	1:D:359:LEU:C	2.39	0.43
1:C:414:LYS:HB3	1:C:414:LYS:HE3	1.59	0.43
1:B:472:HIS:CD2	1:B:474:GLN:HG3	2.53	0.43
1:A:363:PRO:CD	1:A:381:LYS:HG2	2.45	0.43
1:A:472:HIS:HE1	1:A:474:GLN:HG3	1.83	0.43
1:C:427:TYR:O	1:C:440:VAL:HG22	2.19	0.43
1:A:285:MET:HE1	1:A:299:ASN:HB3	2.00	0.43
1:A:310:LEU:HD13	1:A:517:LEU:HD13	2.02	0.42
1:C:356:LYS:HE3	1:C:357:TYR:CE2	2.54	0.42
1:C:314:LEU:HA	1:C:314:LEU:HD12	1.87	0.42
1:C:325:ARG:HB2	1:C:451:TYR:CD2	2.55	0.42
1:C:313:SER:OG	1:C:356:LYS:HE2	2.20	0.42
1:A:412:SER:HB2	1:D:515:GLU:HG2	2.01	0.42
1:B:303:LYS:HA	1:B:303:LYS:HD2	1.74	0.42
1:D:404:TRP:CZ3	1:D:415:LYS:HG3	2.55	0.42
1:D:367:LEU:HD11	1:D:385:LEU:HD23	2.02	0.41
1:B:310:LEU:HD13	1:B:311:ASP:OD1	2.20	0.41
1:B:434:LYS:HE3	1:B:434:LYS:HB2	1.96	0.41
1:D:404:TRP:HB2	1:D:486:CYS:HB3	2.03	0.40
1:A:274:ARG:HD2	3:A:757:HOH:O	2.22	0.40
1:A:294:ARG:HB2	1:A:329:ASP:HB3	2.03	0.40
1:D:285:MET:HE2	1:D:285:MET:HB2	1.93	0.40
1:B:472:HIS:NE2	1:B:474:GLN:HG3	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/255 (98%)	238 (95%)	12 (5%)	0	100	100
1	B	252/255 (99%)	239 (95%)	13 (5%)	0	100	100
1	C	237/255 (93%)	228 (96%)	9 (4%)	0	100	100
1	D	236/255 (92%)	227 (96%)	9 (4%)	0	100	100
All	All	975/1020 (96%)	932 (96%)	43 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/236 (99%)	220 (94%)	14 (6%)	24	37
1	B	236/236 (100%)	216 (92%)	20 (8%)	13	20
1	C	225/236 (95%)	211 (94%)	14 (6%)	23	35
1	D	224/236 (95%)	211 (94%)	13 (6%)	25	39
All	All	919/944 (97%)	858 (93%)	61 (7%)	21	32

All (61) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	268	VAL
1	A	278	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	285	MET
1	A	348	LEU
1	A	368	LEU
1	A	370	LYS
1	A	372	GLU
1	A	388[A]	CYS
1	A	388[B]	CYS
1	A	400	GLU
1	A	427	TYR
1	A	430	LYS
1	A	457	ARG
1	A	464	THR
1	B	270	LYS
1	B	278	SER
1	B	282	SER
1	B	290	ARG
1	B	300	LEU
1	B	303	LYS
1	B	310	LEU
1	B	348	LEU
1	B	359	LEU
1	B	368	LEU
1	B	370	LYS
1	B	373	THR
1	B	388[A]	CYS
1	B	388[B]	CYS
1	B	400	GLU
1	B	419	LEU
1	B	427	TYR
1	B	432	LYS
1	B	436	SER
1	B	457	ARG
1	C	281	SER
1	C	282	SER
1	C	283	LYS
1	C	318	VAL
1	C	370	LYS
1	C	388	CYS
1	C	414	LYS
1	C	421	ARG
1	C	427	TYR
1	C	428	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	440	VAL
1	C	446	ASP
1	C	454	GLN
1	C	478	LYS
1	D	280	ASP
1	D	285	MET
1	D	345	GLN
1	D	348	LEU
1	D	370	LYS
1	D	386	GLU
1	D	388	CYS
1	D	412	SER
1	D	419	LEU
1	D	427	TYR
1	D	446	ASP
1	D	468	LEU
1	D	478	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	330	HIS
1	A	508	GLN
1	B	472	HIS
1	B	474	GLN
1	D	305	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLI	A	601	-	0,6,6	0.00	-	0,7,7	0.00	-
2	MLI	B	601	-	0,6,6	0.00	-	0,7,7	0.00	-
2	MLI	C	601	-	0,6,6	0.00	-	0,7,7	0.00	-
2	MLI	D	601	-	0,6,6	0.00	-	0,7,7	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLI	A	601	-	-	0/0/4/4	0/0/0/0
2	MLI	B	601	-	-	0/0/4/4	0/0/0/0
2	MLI	C	601	-	-	0/0/4/4	0/0/0/0
2	MLI	D	601	-	-	0/0/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	601	MLI	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	251/255 (98%)	0.23	11 (4%) 38 39	13, 33, 65, 78	0
1	B	250/255 (98%)	0.27	14 (5%) 28 28	14, 34, 64, 88	0
1	C	240/255 (94%)	0.13	7 (2%) 55 54	14, 32, 58, 86	1 (0%)
1	D	240/255 (94%)	0.31	14 (5%) 26 27	13, 37, 66, 94	0
All	All	981/1020 (96%)	0.24	46 (4%) 35 36	13, 34, 64, 94	1 (0%)

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	428	VAL	7.3
1	B	433	ALA	6.9
1	B	438	ASP	5.0
1	A	438	ASP	4.7
1	B	440	VAL	4.3
1	D	485	LEU	3.9
1	D	428	VAL	3.9
1	A	518	LYS	3.5
1	A	433	ALA	3.5
1	B	431	GLY	3.4
1	B	432	LYS	3.3
1	B	434	LYS	3.3
1	B	397	PRO	3.2
1	A	396	VAL	3.2
1	D	398	GLU	3.1
1	A	409	GLY	3.1
1	B	441	CYS	3.0
1	C	427	TYR	2.9
1	A	397	PRO	2.9
1	C	426	TYR	2.8
1	B	404	TRP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	426	TYR	2.7
1	D	440	VAL	2.6
1	A	431	GLY	2.6
1	B	437	ARG	2.6
1	D	427	TYR	2.5
1	A	435	VAL	2.5
1	D	278	SER	2.5
1	B	442	PHE	2.4
1	D	416	ARG	2.4
1	B	435	VAL	2.3
1	D	377	ALA	2.3
1	D	429	PRO	2.2
1	B	423	SER	2.2
1	D	405	LEU	2.2
1	C	398	GLU	2.1
1	C	405	LEU	2.1
1	C	440	VAL	2.1
1	A	404	TRP	2.1
1	D	280	ASP	2.1
1	B	396	VAL	2.1
1	D	442	PHE	2.1
1	D	483	LYS	2.0
1	C	429	PRO	2.0
1	A	437	ARG	2.0
1	A	432	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MLI	D	601	7/7	0.96	0.19	1.72	28,30,30,31	0
2	MLI	B	601	7/7	0.94	0.19	1.50	21,23,26,29	0
2	MLI	C	601	7/7	0.93	0.16	0.62	29,31,34,35	0
2	MLI	A	601	7/7	0.97	0.15	0.52	19,21,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.