



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 04:57 PM GMT

PDB ID : 4GOJ
Title : The Crystal Structure of full length Arl3GppNHp in complex with UNC119a
Authors : Ismail, S.; Xiang-Chen, Y.; Miertzschke, M.; Vetter, I.; Koerner, C.; Wittinghofer, A.
Deposited on : 2012-08-20
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	<div> <div>3%</div> <div>80% 12% • 7%</div> </div>
1	B	189	<div> <div>9%</div> <div>79% 14% • 7%</div> </div>
2	C	240	<div> <div>2%</div> <div>56% 10% • 33%</div> </div>
2	D	240	<div> <div>•</div> <div>56% 9% • 33%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5712 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosylation factor-like protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	175	Total	C	N	O	S	0	0	0
			1382	876	238	264	4			
1	B	176	Total	C	N	O	S	0	0	0
			1392	881	241	266	4			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	183	GLU	-	EXPRESSION TAG	UNP Q9WUL7
A	184	HIS	-	EXPRESSION TAG	UNP Q9WUL7
A	185	HIS	-	EXPRESSION TAG	UNP Q9WUL7
A	186	HIS	-	EXPRESSION TAG	UNP Q9WUL7
A	187	HIS	-	EXPRESSION TAG	UNP Q9WUL7
A	188	HIS	-	EXPRESSION TAG	UNP Q9WUL7
A	189	HIS	-	EXPRESSION TAG	UNP Q9WUL7
B	183	GLU	-	EXPRESSION TAG	UNP Q9WUL7
B	184	HIS	-	EXPRESSION TAG	UNP Q9WUL7
B	185	HIS	-	EXPRESSION TAG	UNP Q9WUL7
B	186	HIS	-	EXPRESSION TAG	UNP Q9WUL7
B	187	HIS	-	EXPRESSION TAG	UNP Q9WUL7
B	188	HIS	-	EXPRESSION TAG	UNP Q9WUL7
B	189	HIS	-	EXPRESSION TAG	UNP Q9WUL7

- Molecule 2 is a protein called Protein unc-119 homolog A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	161	Total	C	N	O	S	0	1	0
			1350	872	228	243	7			
2	D	161	Total	C	N	O	S	0	2	0
			1348	871	225	245	7			

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter

Chemical structure of GNP (Guanosine Nucleotide Phosphate) showing the guanine base, ribose sugar, and three phosphate groups. Atoms are labeled with IDs and element symbols. Stereochemistry is indicated with wedges and dashes.

Atom IDs and Element Symbols:

- O6, O5, O4, O3, O2, O1, O: Oxygen
- N7, N6, N5, N4, N3, N2, N1: Nitrogen
- C8, C7, C6, C5, C4, C3, C2, C1: Carbon
- P1, P2, P3: Phosphorus
- H: Hydrogen

Key features include the guanine base (N1-C8), the ribose sugar (C1'-C5'), and the three phosphate groups (P1, P2, P3) attached to the 5' carbon of the sugar. Stereochemistry is indicated with wedges and dashes.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 32	C 10	N 6	O 13	P 3	0	0
3	B	1	Total 32	C 10	N 6	O 13	P 3	0	0


- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 4 | B | 1 | Total Mg
1 1 | 0 | 0 |
| 4 | A | 1 | Total Mg
1 1 | 0 | 0 |

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|------------------|---------|---------|
| 5 | A | 67 | Total O
67 67 | 0 | 0 |
| 5 | B | 29 | Total O
29 29 | 0 | 0 |
| 5 | C | 35 | Total O
35 35 | 0 | 0 |
| 5 | D | 43 | Total O
43 43 | 0 | 0 |


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ADP-ribosylation factor-like protein 3



• Molecule 1: ADP-ribosylation factor-like protein 3




• Molecule 2: Protein unc-119 homolog A



• Molecule 2: Protein unc-119 homolog A





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.75Å 103.30Å 119.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.97 – 2.10 29.97 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.97-2.10) 99.7 (29.97-2.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.58 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.6.0093	Depositor
R, R_{free}	0.214 , 0.248 0.213 , 0.249	Depositor DCC
R_{free} test set	2556 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	29.9	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 46.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 51107 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5712	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1403	0.58	1/1899 (0.1%)
1	B	0.38	0/1414	0.59	1/1913 (0.1%)
2	C	0.44	0/1391	0.57	0/1875
2	D	0.43	0/1391	0.57	1/1876 (0.1%)
All	All	0.41	0/5599	0.58	3/7563 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	59	PRO	N-CA-CB	5.65	110.08	103.30
1	A	7	LEU	CA-CB-CG	5.48	127.91	115.30
1	B	7	LEU	CA-CB-CG	5.40	127.72	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	59	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1382	0	1398	14	0
1	B	1392	0	1409	18	0
2	C	1350	0	1301	14	0
2	D	1348	0	1290	15	0
3	A	32	0	13	1	0
3	B	32	0	13	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	67	0	0	0	0
5	B	29	0	0	2	0
5	C	35	0	0	0	0
5	D	43	0	0	2	0
All	All	5712	0	5424	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:ILE:HD11	1:A:177:VAL:HG12	1.49	0.95
2:C:139:ARG:HH11	2:C:139:ARG:HG3	1.45	0.81
1:B:16:GLN:HB2	1:B:61:PHE:CE1	2.24	0.73
2:C:139:ARG:HH11	2:C:139:ARG:CG	2.02	0.72
1:A:87:ILE:CD1	1:A:177:VAL:HG12	2.20	0.70

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/189 (92%)	172 (99%)	1 (1%)	0	100	100
1	B	174/189 (92%)	168 (97%)	6 (3%)	0	100	100
2	C	158/240 (66%)	154 (98%)	4 (2%)	0	100	100
2	D	159/240 (66%)	158 (99%)	1 (1%)	0	100	100
All	All	664/858 (77%)	652 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	152/166 (92%)	146 (96%)	6 (4%)	39	39
1	B	154/166 (93%)	148 (96%)	6 (4%)	39	39
2	C	150/210 (71%)	142 (95%)	8 (5%)	28	25
2	D	149/210 (71%)	140 (94%)	9 (6%)	24	20
All	All	605/752 (80%)	576 (95%)	29 (5%)	31	29

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	66	LEU
2	C	127	ARG
2	D	143	VAL
2	C	69	GLN
2	C	139	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	82	ASN
2	D	82	ASN
2	C	170	GLN
1	B	156	GLN
2	C	158	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GNP	A	201	4	28,34,34	1.97	8 (28%)	33,54,54	2.01	9 (27%)
3	GNP	B	201	4	28,34,34	1.93	9 (32%)	33,54,54	2.14	8 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GNP	A	201	4	-	1/12/38/38	0/3/3/3
3	GNP	B	201	4	-	1/12/38/38	0/3/3/3

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	201	GNP	PG-O3G	-3.58	1.46	1.56
3	B	201	GNP	C6-C5	-3.55	1.34	1.41
3	B	201	GNP	PG-O2G	-3.51	1.46	1.56
3	A	201	GNP	C6-C5	-3.45	1.34	1.41
3	B	201	GNP	PG-O3G	-3.15	1.47	1.56

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	201	GNP	O1G-PG-N3B	-6.49	101.94	111.90
3	A	201	GNP	O1G-PG-N3B	-5.66	103.22	111.90
3	B	201	GNP	N3-C2-N1	-5.19	119.53	127.44
3	A	201	GNP	N3-C2-N1	-4.99	119.85	127.44
3	A	201	GNP	C5-C6-N1	-3.34	119.03	123.59

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	201	GNP	O1B-PB-N3B-PG
3	B	201	GNP	O1B-PB-N3B-PG

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	201	GNP	1	0
3	B	201	GNP	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	175/189 (92%)	0.13	5 (2%) 55 63	15, 25, 44, 59	0
1	B	176/189 (93%)	0.64	17 (9%) 10 14	21, 37, 59, 76	0
2	C	161/240 (67%)	0.23	5 (3%) 52 61	16, 32, 52, 63	0
2	D	161/240 (67%)	0.11	3 (1%) 70 75	20, 30, 46, 58	0
All	All	673/858 (78%)	0.28	30 (4%) 37 46	15, 31, 54, 76	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	149	ARG	4.8
1	B	14	PRO	4.6
2	D	208	ILE	4.1
1	B	146	HIS	3.9
1	B	43	SER	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MG	B	202	1/1	0.91	0.11	-0.82	26,26,26,26	0
3	GNP	B	201	32/32	0.97	0.10	-1.01	24,27,29,30	0
3	GNP	A	201	32/32	0.98	0.08	-1.08	16,21,23,25	0
4	MG	A	202	1/1	0.98	0.04	-3.02	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.