



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:40 PM GMT

PDB ID : 1GPW
Title : STRUCTURAL EVIDENCE FOR AMMONIA TUNNELING ACROSS THE (BETA/ALPHA)8 BARREL OF THE IMIDAZOLE GLYCEROL PHOSPHATE SYNTHASE BIENZYME COMPLEX.
Authors : Walker, M.; Beismann-Driemeyer, S.; Sterner, R.; Wilmanns, M.
Deposited on : 2001-11-12
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	253	
1	C	253	
1	E	253	
2	B	201	
2	D	201	
2	F	201	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISF PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			1937	1232	330	369	6			
1	C	253	Total	C	N	O	S	0	0	0
			1937	1230	331	370	6			
1	E	251	Total	C	N	O	S	0	0	0
			1927	1225	328	368	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	ASN	ASP	ENGINEERED MUTATION	UNP Q9X0C6
C	11	ASN	ASP	ENGINEERED MUTATION	UNP Q9X0C6
E	11	ASN	ASP	ENGINEERED MUTATION	UNP Q9X0C6

- Molecule 2 is a protein called AMIDOTRANSFERASE HISH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	200	Total	C	N	O	S	0	0	0
			1616	1029	284	295	8			
2	D	201	Total	C	N	O	S	0	0	0
			1612	1028	281	295	8			
2	F	201	Total	C	N	O	S	0	0	0
			1614	1029	282	295	8			

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	P	1	Total	O	P	0	0
			5	4	1		
3	P	1	Total	O	P	0	0
			5	4	1		
3	P	1	Total	O	P	0	0
			5	4	1		
3	P	1	Total	O	P	0	0
			5	4	1		
3	P	1	Total	O	P	0	0
			5	4	1		
3	P	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	118	Total	O	0	0
			118	118		
4	B	51	Total	O	0	0
			51	51		
4	C	93	Total	O	0	0
			93	93		
4	D	85	Total	O	0	0
			85	85		
4	E	38	Total	O	0	0
			38	38		
4	F	71	Total	O	0	0
			71	71		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: HISF PROTEIN




• Molecule 1: HISF PROTEIN




• Molecule 1: HISF PROTEIN






● Molecule 2: AMIDOTRANSFERASE HISH

Chain B: 50% 45%




● Molecule 2: AMIDOTRANSFERASE HISH

Chain D: 59% 36% 5%



● Molecule 2: AMIDOTRANSFERASE HISH

Chain F: 49% 46% 5%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.00 Å 91.90 Å 131.00 Å 90.00° 105.90° 90.00°	Depositor
Resolution (Å)	30.00 – 2.40	Depositor
% Data completeness (in resolution range)	97.8 (30.00-2.40)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.226 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11132	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1964	0.67	0/2648
1	C	0.43	0/1963	0.67	0/2647
1	E	0.46	0/1952	0.69	0/2630
2	B	0.45	0/1650	0.67	0/2218
2	D	0.44	0/1646	0.68	0/2214
2	F	0.43	0/1648	0.67	0/2217
All	All	0.44	0/10823	0.68	0/14574

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1937	0	1967	69	0
1	C	1937	0	1973	71	1
1	E	1927	0	1962	206	1
2	B	1616	0	1611	90	1
2	D	1612	0	1598	72	0
2	F	1614	0	1599	95	1
3	P	30	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	118	0	0	5	0
4	B	51	0	0	6	0
4	C	93	0	0	4	0
4	D	85	0	0	5	0
4	E	38	0	0	3	0
4	F	71	0	0	6	0
4	P	3	0	0	0	0
All	All	11132	0	10710	589	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 589 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:53:THR:HG21	1:E:57:GLU:HB2	1.34	1.07
1:E:9:CYS:HB2	1:E:222:LEU:HD11	1.34	1.06
1:E:53:THR:HG22	1:E:54:ALA:H	1.18	1.04
1:E:123:GLN:HE21	1:E:123:GLN:N	1.60	1.00
2:F:1:MET:HE2	2:F:194:ILE:HG23	1.41	0.99

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:55:SER:N	2:F:200:ARG:CB[1_456]	1.73	0.47
2:B:114:ARG:NH2	1:E:208:GLU:OE2[1_455]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/253 (99%)	231 (92%)	14 (6%)	6 (2%)	7	7
1	C	251/253 (99%)	228 (91%)	18 (7%)	5 (2%)	9	11
1	E	247/253 (98%)	184 (74%)	35 (14%)	28 (11%)	0	0
2	B	198/201 (98%)	167 (84%)	25 (13%)	6 (3%)	5	4
2	D	199/201 (99%)	171 (86%)	20 (10%)	8 (4%)	4	3
2	F	199/201 (99%)	177 (89%)	13 (6%)	9 (4%)	3	2
All	All	1345/1362 (99%)	1158 (86%)	125 (9%)	62 (5%)	3	2

5 of 62 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	24	GLU
1	A	55	SER
1	A	251	GLU
1	C	19	LYS
1	C	30	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	204/208 (98%)	193 (95%)	11 (5%)	27	43
1	C	205/208 (99%)	192 (94%)	13 (6%)	22	35
1	E	204/208 (98%)	184 (90%)	20 (10%)	10	14
2	B	176/177 (99%)	165 (94%)	11 (6%)	22	35
2	D	174/177 (98%)	166 (95%)	8 (5%)	33	51
2	F	174/177 (98%)	162 (93%)	12 (7%)	19	30
All	All	1137/1155 (98%)	1062 (93%)	75 (7%)	21	32

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	228	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	153	LEU
2	F	117	ARG
1	C	235	ARG
2	D	56	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	109	ASN
1	C	148	ASN
2	F	88	GLN
1	C	118	GLN
1	A	115	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	P	1	-	4,4,4	1.08	0	6,6,6	0.27	0
3	PO4	P	2	-	4,4,4	0.94	0	6,6,6	0.27	0
3	PO4	P	3	-	4,4,4	0.91	0	6,6,6	0.27	0
3	PO4	P	4	-	4,4,4	0.99	0	6,6,6	0.27	0
3	PO4	P	5	-	4,4,4	0.76	0	6,6,6	0.27	0
3	PO4	P	6	-	4,4,4	0.83	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	P	1	-	-	0/0/0/0	0/0/0/0
3	PO4	P	2	-	-	0/0/0/0	0/0/0/0
3	PO4	P	3	-	-	0/0/0/0	0/0/0/0
3	PO4	P	4	-	-	0/0/0/0	0/0/0/0
3	PO4	P	5	-	-	0/0/0/0	0/0/0/0
3	PO4	P	6	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.