



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:02 PM GMT

PDB ID : 4GXJ
Title : R283K DNA polymerase beta ternary complex with a templating 8OG and incoming dCTP analog
Authors : Freudenthal, B.D.; Beard, W.A.; Wilson, S.H.
Deposited on : 2012-09-04
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	
2	D	5	
3	P	10	
4	T	16	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3300 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	12	1	0
			2526	1602	438	477	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	283	LYS	ARG	ENGINEERED MUTATION	UNP P06746

- Molecule 2 is a DNA chain called DNA (5'-D(P*GP*TP*CP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	5	Total	C	N	O	P	0	0	0
			106	49	20	32	5			


- Molecule 3 is a DNA chain called DNA (5'-D(*GP*CP*TP*GP*AP*TP*GP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	10	Total	C	N	O	P	0	0	0
			205	98	40	58	9			

- Molecule 4 is a DNA chain called DNA (5'-D(*CP*CP*GP*AP*CP*(8OG)P*TP*CP*GP*CP*AP*TP*CP*AP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	T	16	Total	C	N	O	P	0	0	0
			322	153	60	94	15			

- Molecule 5 is 2'-DEOXY-5'-O-[(S)-{DIFLUORO[(S)-HYDROXY(PHOSPHONOOXY)PHOSPHORYL]METHYL}(HYDROXY)PHOSPHORYL]CYTIDINE (three-letter code: 6CF) (formula: C₁₀H₁₆F₂N₃O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	P	
			30	10	2	3	12	3	

- Molecule 6 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	P	1	Total	Mn	0	0
			1	1		
6	A	4	Total	Mn	0	0
			4	4		
6	T	2	Total	Mn	0	0
			2	2		

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Na	0	0
			1	1		
7	D	1	Total	Na	0	0
			1	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	73	Total	O	0	0
			73	73		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	D	4	Total 4	O 4	0	0
8	P	10	Total 10	O 10	0	0
8	T	15	Total 15	O 15	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase beta



- Molecule 2: DNA (5'-D(P*GP*TP*CP*GP*G)-3')



- Molecule 3: DNA (5'-D(*GP*CP*TP*GP*AP*TP*GP*CP*GP*A)-3')



- Molecule 4: DNA (5'-D(*CP*CP*GP*AP*CP*(8OG)P*TP*CP*GP*CP*AP*TP*CP*AP*GP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.08 Å 77.43 Å 55.12 Å 90.00° 114.45° 90.00°	Depositor
Resolution (Å)	21.06 – 2.20 21.06 – 2.20	Depositor EDS
% Data completeness (in resolution range)	86.5 (21.06-2.20) 86.1 (21.06-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 2.19 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7_650)	Depositor
R, R_{free}	0.231 , 0.292 0.241 , 0.286	Depositor DCC
R_{free} test set	2150 reflections (12.80%)	DCC
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.129	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.6	EDS
Estimated twinning fraction	0.047 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 21433 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3300	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MN, 6CF, 8OG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.38	0/2572	0.53	1/3455 (0.0%)
2	D	0.78	0/118	1.24	0/179
3	P	0.83	0/230	1.29	2/354 (0.6%)
4	T	0.65	0/333	1.40	5/508 (1.0%)
All	All	0.48	0/3253	0.79	8/4496 (0.2%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	7	DT	P-O5'-C5'	7.79	133.37	120.90
4	T	7	DT	OP1-P-OP2	6.67	129.61	119.60
3	P	6	DT	O4'-C1'-N1	-5.81	103.93	108.00
3	P	6	DT	N3-C4-O4	5.75	123.35	119.90
4	T	9	DG	O4'-C1'-N9	-5.74	103.98	108.00
4	T	12	DT	N3-C4-O4	5.59	123.25	119.90
1	A	314	ASP	CB-CG-OD2	5.21	122.99	118.30
4	T	12	DT	P-O5'-C5'	-5.20	112.57	120.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2526	0	2521	91	0
2	D	106	0	57	1	0
3	P	205	0	113	8	0
4	T	322	0	180	13	0
5	A	30	0	12	2	0
6	A	4	0	0	0	0
6	P	1	0	0	0	0
6	T	2	0	0	0	0
7	A	1	0	0	0	0
7	D	1	0	0	0	0
8	A	73	0	0	1	0
8	D	4	0	0	0	0
8	P	10	0	0	0	0
8	T	15	0	0	0	0
All	All	3300	0	2883	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (110) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:P:5:DA:H4'	3:P:6:DT:OP1	1.37	1.13
1:A:291:PHE:HD1	1:A:323:ILE:HG22	1.37	0.88
1:A:75:GLU:OE2	1:A:83:ARG:HG3	1.74	0.88
1:A:201:THR:H	1:A:204:SER:HB3	1.39	0.86
1:A:82:LEU:HD23	1:A:85:LEU:HG	1.54	0.86
1:A:290:GLY:HA2	1:A:301:LEU:HD22	1.60	0.82
3:P:4:DG:C6	3:P:5:DA:N6	2.48	0.81
1:A:301:LEU:HA	1:A:307:ALA:HB2	1.63	0.80
4:T:14:DA:H2''	4:T:15:DG:O5'	1.82	0.78
3:P:5:DA:C4'	3:P:6:DT:OP1	2.22	0.78
4:T:15:DG:H2''	4:T:16:DC:H5''	1.67	0.76
4:T:14:DA:H2'	4:T:15:DG:C8	2.21	0.75
4:T:5:DC:H4'	4:T:6:8OG:OP2	1.88	0.74
1:A:157:GLN:HG2	1:A:241:LEU:CD2	2.18	0.74
1:A:157:GLN:HG2	1:A:241:LEU:HD22	1.70	0.74
1:A:291:PHE:CD1	1:A:323:ILE:HG22	2.20	0.73
3:P:4:DG:C5	3:P:5:DA:N6	2.58	0.71
1:A:25:PHE:CE2	1:A:88:ILE:HD12	2.26	0.70
1:A:197:HIS:CE1	1:A:199:SER:HB3	2.29	0.68
1:A:195:LEU:HD21	1:A:214:VAL:HG21	1.78	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:GLY:HA2	1:A:301:LEU:HB2	1.78	0.65
1:A:145:ASP:OD2	8:A:546:HOH:O	2.15	0.64
3:P:8:DC:H2''	3:P:9:DG:H5'	1.78	0.64
1:A:290:GLY:CA	1:A:301:LEU:HB2	2.27	0.64
1:A:251:PRO:HG2	1:A:253:ARG:CZ	2.28	0.64
4:T:14:DA:H2''	4:T:15:DG:C5'	2.27	0.64
1:A:270:LEU:HD22	1:A:319:ILE:HD13	1.79	0.64
1:A:209:LYS:HB3	1:A:212:HIS:HB3	1.79	0.63
1:A:216:GLU:O	1:A:220:LYS:HG2	1.98	0.63
1:A:233:THR:HB	1:A:258:ARG:HH12	1.62	0.63
1:A:84:LYS:O	1:A:88:ILE:HG12	1.99	0.62
1:A:82:LEU:HD23	1:A:85:LEU:CG	2.29	0.62
1:A:25:PHE:CD2	1:A:88:ILE:HD12	2.35	0.62
1:A:292:THR:O	1:A:292:THR:HG23	1.99	0.62
1:A:162:VAL:O	1:A:166:VAL:HG23	2.00	0.62
1:A:197:HIS:ND1	1:A:198:PRO:HD2	2.16	0.61
1:A:290:GLY:HA2	1:A:301:LEU:CD2	2.31	0.60
1:A:299:ARG:NH1	1:A:310:PRO:HD3	2.19	0.57
1:A:197:HIS:CE1	1:A:198:PRO:HD2	2.39	0.57
1:A:291:PHE:HA	1:A:299:ARG:O	2.05	0.57
1:A:300:PRO:HD2	1:A:309:GLU:H	1.69	0.56
1:A:285:HIS:HA	1:A:288:GLU:HG3	1.87	0.55
1:A:172:GLU:CG	1:A:198:PRO:HG2	2.37	0.55
1:A:45:VAL:HG11	1:A:62:LEU:HB3	1.88	0.54
1:A:15:ILE:HG21	1:A:46:ILE:HD13	1.89	0.54
1:A:279:ASN:HB3	1:A:283:LYS:HE2	1.90	0.53
1:A:290:GLY:O	1:A:301:LEU:N	2.42	0.53
1:A:279:ASN:HB3	1:A:283:LYS:CE	2.38	0.53
1:A:279:ASN:O	1:A:283:LYS:HD2	2.09	0.53
3:P:8:DC:C2'	3:P:9:DG:H5'	2.38	0.53
1:A:270:LEU:CD2	1:A:319:ILE:HD13	2.38	0.53
1:A:231:GLY:HA3	4:T:10:DC:OP1	2.10	0.52
1:A:201:THR:OG1	1:A:203:GLU:HG2	2.09	0.52
1:A:45:VAL:HG21	1:A:63:PRO:O	2.11	0.51
1:A:201:THR:HB	1:A:263:ASP:OD2	2.11	0.51
4:T:3:DG:H4'	4:T:4:DA:OP1	2.10	0.51
4:T:1:DC:H2'	4:T:2:DC:C6	2.46	0.50
1:A:33:ILE:O	1:A:36:TYR:HB3	2.10	0.50
1:A:48:LYS:HD2	1:A:335:GLU:OE1	2.11	0.50
1:A:298:ILE:O	1:A:298:ILE:HG23	2.11	0.50
1:A:97:ILE:O	1:A:101:THR:HG23	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:PHE:HZ	1:A:122:LEU:HG	1.77	0.50
1:A:233:THR:HB	1:A:258:ARG:NH1	2.25	0.50
1:A:150:ILE:HG22	1:A:155:MET:HG2	1.95	0.49
1:A:196:THR:HG21	1:A:265:TYR:HD1	1.78	0.49
1:A:276:ASP:OD2	1:A:277:ILE:N	2.46	0.49
1:A:183:ARG:O	1:A:331:LYS:HA	2.13	0.48
1:A:150:ILE:HG12	1:A:253:ARG:NE	2.28	0.48
1:A:276:ASP:OD2	1:A:277:ILE:HG12	2.14	0.48
1:A:82:LEU:HD23	1:A:85:LEU:CD1	2.44	0.48
1:A:197:HIS:ND1	1:A:199:SER:HB3	2.29	0.48
1:A:133:ASN:OD1	1:A:136:GLN:HG3	2.14	0.48
1:A:22:LEU:CD2	1:A:85:LEU:HD21	2.45	0.47
1:A:254:ARG:NH2	1:A:256:ASP:OD1	2.47	0.47
1:A:157:GLN:HG2	1:A:241:LEU:HD21	1.95	0.47
1:A:127:LYS:HB2	1:A:127:LYS:HE2	1.66	0.46
1:A:172:GLU:HG2	1:A:198:PRO:HG2	1.96	0.46
1:A:167:LYS:HB3	1:A:167:LYS:HE3	1.67	0.46
4:T:1:DC:H2''	4:T:2:DC:H5'	1.97	0.46
5:A:401:6CF:N4	3:P:10:DA:N6	2.63	0.46
1:A:157:GLN:O	1:A:161:ILE:HG13	2.16	0.46
1:A:164:ASN:O	1:A:168:LYS:HG3	2.15	0.46
1:A:311:LEU:HD13	1:A:322:TYR:CE1	2.52	0.45
1:A:45:VAL:CG1	1:A:62:LEU:HB3	2.46	0.45
1:A:168:LYS:HE2	1:A:168:LYS:HB3	1.61	0.45
1:A:15:ILE:H	1:A:15:ILE:HD12	1.82	0.44
5:A:401:6CF:O1A	3:P:10:DA:H3'	2.17	0.44
1:A:165:GLU:OE2	1:A:221:VAL:HG11	2.18	0.44
4:T:15:DG:C2'	4:T:16:DC:H5''	2.45	0.44
1:A:135:HIS:CD2	1:A:135:HIS:C	2.91	0.44
1:A:283:LYS:HE3	4:T:6:8OG:H22	1.82	0.43
1:A:292:THR:O	1:A:292:THR:CG2	2.66	0.43
1:A:33:ILE:HG23	1:A:34:HIS:N	2.33	0.43
1:A:197:HIS:CG	1:A:198:PRO:HD2	2.54	0.42
1:A:194:LEU:HD11	1:A:260:ILE:HB	2.01	0.42
1:A:292:THR:HG22	1:A:299:ARG:HB2	2.01	0.42
1:A:194:LEU:CD1	1:A:260:ILE:HD12	2.49	0.42
1:A:178:CYS:SG	1:A:269:VAL:HG23	2.59	0.42
2:D:3:DC:H2''	2:D:4:DG:C8	2.55	0.42
1:A:300:PRO:HD2	1:A:309:GLU:N	2.34	0.41
1:A:267:CYS:SG	1:A:313:VAL:HG21	2.60	0.41
4:T:14:DA:H2'	4:T:15:DG:H8	1.78	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:LYS:HG2	1:A:293:ILE:O	2.20	0.41
1:A:280:LYS:NZ	4:T:6:8OG:OP1	2.54	0.41
1:A:142:TYR:CE1	1:A:252:HIS:CD2	3.08	0.41
1:A:18:MET:HG3	1:A:19:LEU:N	2.36	0.41
1:A:272:PHE:HA	1:A:272:PHE:HD1	1.77	0.40
1:A:197:HIS:CG	1:A:198:PRO:CD	3.05	0.40
1:A:317:LYS:HG3	1:A:327:TYR:CG	2.57	0.40
1:A:260:ILE:HD13	1:A:268:GLY:HA3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/335 (92%)	289 (94%)	18 (6%)	1 (0%)	46	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	285	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	274/295 (93%)	267 (97%)	7 (3%)	54 66

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	MET
1	A	172	GLU
1	A	199	SER
1	A	272	PHE
1	A	301	LEU
1	A	318	ASP
1	A	334	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	157	GLN
1	A	159	GLN
1	A	164	ASN
1	A	213	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	8OG	T	6	4	16,25,26	2.44	4 (25%)	21,37,40	2.56	11 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	8OG	T	6	4	-	0/3/21/22	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	T	6	8OG	O4'-C4'	-2.02	1.40	1.45
4	T	6	8OG	C8-N7	2.92	1.37	1.34
4	T	6	8OG	C2-N2	5.61	1.45	1.34
4	T	6	8OG	O6-C6	6.40	1.40	1.24

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	6	8OG	N3-C2-N1	-3.39	122.28	127.44
4	T	6	8OG	C1'-N9-C4	-3.29	122.27	127.37
4	T	6	8OG	C4'-O4'-C1'	-3.28	101.18	109.47
4	T	6	8OG	C6-C5-C4	-3.24	117.02	120.90
4	T	6	8OG	O3'-C3'-C4'	-3.08	97.58	110.05
4	T	6	8OG	C5-C6-N1	-2.05	120.78	123.59
4	T	6	8OG	O4'-C4'-C3'	2.19	111.17	105.67
4	T	6	8OG	C6-N1-C2	2.70	119.69	115.94
4	T	6	8OG	O4'-C1'-N9	2.80	110.59	108.22
4	T	6	8OG	O3'-C3'-C2'	4.09	124.30	110.74
4	T	6	8OG	O4'-C1'-C2'	5.44	117.12	106.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	T	6	8OG	3	0

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 9 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	6CF	A	401	6	26,31,31	1.97	10 (38%)	28,50,50	1.91	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	6CF	A	401	6	-	0/17/46/46	0/2/2/2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	401	6CF	F1A-C3A	-3.42	1.32	1.36
5	A	401	6CF	P2-C3A	-3.15	1.82	1.85
5	A	401	6CF	P1-C3A	-2.94	1.82	1.85
5	A	401	6CF	O4'-C4'	-2.80	1.38	1.45
5	A	401	6CF	O3'-C3'	-2.17	1.38	1.43
5	A	401	6CF	C4-N3	2.10	1.39	1.35
5	A	401	6CF	C6-N1	2.84	1.39	1.35
5	A	401	6CF	P2-O1B	2.92	1.56	1.51
5	A	401	6CF	C4-N4	2.96	1.43	1.35
5	A	401	6CF	P1-O2A	3.42	1.57	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	401	6CF	P3-O3B-P2	-4.50	117.58	132.67
5	A	401	6CF	C5-C4-N3	-3.97	116.79	121.80
5	A	401	6CF	O1G-P3-O3G	-2.21	103.45	110.58
5	A	401	6CF	O4'-C1'-N1	2.44	111.95	107.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	401	6CF	O1G-P3-O3B	2.58	116.80	105.09
5	A	401	6CF	C2-N3-C4	2.95	119.77	115.61
5	A	401	6CF	N4-C4-N3	3.40	122.70	116.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	401	6CF	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	315/335 (94%)	1.03	55 (17%) 2 2	16, 39, 71, 86	0
2	D	5/5 (100%)	0.26	0 100 100	29, 32, 47, 48	0
3	P	10/10 (100%)	0.59	1 (10%) 9 8	24, 38, 46, 52	0
4	T	15/16 (93%)	0.63	1 (6%) 21 20	29, 44, 59, 61	0
All	All	345/366 (94%)	0.99	57 (16%) 2 2	16, 39, 70, 86	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	ASP	8.0
1	A	290	GLY	7.4
1	A	287	LEU	6.9
1	A	307	ALA	6.1
1	A	300	PRO	5.5
1	A	204	SER	5.2
1	A	293	ILE	4.1
1	A	245	ASN	4.1
1	A	314	ASP	3.8
1	A	268	GLY	3.7
1	A	243	SER	3.7
1	A	331	LYS	3.7
1	A	257	ILE	3.7
1	A	203	GLU	3.6
1	A	270	LEU	3.6
1	A	325	TRP	3.6
1	A	116	ASP	3.6
1	A	319	ILE	3.5
1	A	329	GLU	3.4
1	A	79	THR	3.3
1	A	267	CYS	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	22	LEU	3.1
1	A	39	TYR	3.1
1	A	191	MET	3.1
1	A	327	TYR	2.9
1	A	255	ILE	2.9
1	A	292	THR	2.8
1	A	69	ILE	2.7
1	A	298	ILE	2.7
1	A	90	GLN	2.7
1	A	288	GLU	2.7
1	A	326	LYS	2.6
1	A	19	LEU	2.6
1	A	301	LEU	2.6
1	A	209	LYS	2.6
1	A	271	TYR	2.6
4	T	1	DC	2.5
1	A	285	HIS	2.5
1	A	202	SER	2.5
1	A	282	MET	2.5
1	A	272	PHE	2.5
3	P	5	DA	2.4
1	A	222	HIS	2.4
1	A	193	VAL	2.4
1	A	244	LYS	2.4
1	A	10	THR	2.3
1	A	291	PHE	2.3
1	A	125	LEU	2.3
1	A	73	ILE	2.3
1	A	201	THR	2.3
1	A	106	ILE	2.2
1	A	130	ASP	2.2
1	A	284	ALA	2.2
1	A	11	LEU	2.1
1	A	100	LEU	2.1
1	A	283	LYS	2.1
1	A	323	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	8OG	T	6	23/24	0.76	0.28	-	47,59,71,86	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	6CF	A	401	30/30	0.89	0.16	-1.14	24,37,44,47	0
7	NA	A	406	1/1	0.98	0.14	-1.35	18,18,18,18	0
6	MN	A	404	1/1	0.86	0.09	-	76,76,76,76	0
6	MN	P	101	1/1	0.88	0.07	-	65,65,65,65	0
6	MN	T	102	1/1	0.90	0.08	-	86,86,86,86	0
6	MN	T	101	1/1	0.65	0.11	-	68,68,68,68	0
6	MN	A	403	1/1	0.98	0.08	-	28,28,28,28	0
7	NA	D	101	1/1	0.86	0.44	-	50,50,50,50	0
6	MN	A	402	1/1	0.96	0.10	-	32,32,32,32	0
6	MN	A	405	1/1	0.33	0.10	-	79,79,79,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.