



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:23 AM GMT

PDB ID : 2GYS
Title : 2.7 Å structure of the extracellular domains of the human beta common receptor involved in IL-3, IL-5, and GM-CSF signalling
Authors : Carr, P.D.; Conlan, F.; Ford, S.; Ollis, D.L.; Young, I.G.
Deposited on : 2006-05-09
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	 4% 58% 29% 7% • 5%
1	B	419	 61% 27% 7% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	1001	X	-	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6454 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytokine receptor common beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	397	Total	C	N	O	S	0	0	0
			3098	1953	551	578	16			
1	B	406	Total	C	N	O	S	0	0	0
			3182	1995	575	595	17			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	328	GLN	ASN	ENGINEERED	GB 47678387
B	328	GLN	ASN	ENGINEERED	GB 47678387

- Molecule 2 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	4	Total	C	N	O	0	0
			49	28	2	19		
2	B	4	Total	C	N	O	0	0
			49	28	2	19		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			38	22	2	14		


- Molecule 4 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	3	Total	C	N	O	0	0
			38	22	2	14		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytokine receptor common beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	184.91Å 184.91Å 101.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.70 46.23 – 2.70	Depositor EDS
% Data completeness (in resolution range)	91.4 (50.00-2.70) 91.4 (46.23-2.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.212 , 0.269 0.209 , 0.267	Depositor DCC
R_{free} test set	1392 reflections (4.50%)	DCC
Wilson B-factor (Å ²)	52.6	Xtriage
Anisotropy	0.310	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 45.5	EDS
Estimated twinning fraction	0.022 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 32327 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6454	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, NDG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	5/3185 (0.2%)	1.02	9/4364 (0.2%)
1	B	0.95	3/3273 (0.1%)	1.02	10/4481 (0.2%)
All	All	0.96	8/6458 (0.1%)	1.02	19/8845 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
2	A	1	0
All	All	1	2

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	210	CYS	CB-SG	-7.39	1.69	1.82
1	A	80	VAL	CB-CG2	7.13	1.67	1.52
1	A	62	CYS	CB-SG	7.10	1.94	1.82
1	A	210	CYS	CB-SG	-6.57	1.71	1.82
1	A	205	TRP	CG-CD1	6.33	1.45	1.36

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	33	VAL	CB-CA-C	-8.22	95.78	111.40
1	B	394	VAL	CB-CA-C	-7.58	96.99	111.40
1	A	10	ARG	CG-CD-NE	-6.64	97.85	111.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	B	40	ARG	NE-CZ-NH2	-6.20	117.20	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1001	NAG	C1

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	76	CYS	Peptide
1	B	18	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	2866	110	0
1	B	3182	0	2953	116	0
2	A	49	0	43	3	0
2	B	49	0	43	1	0
3	A	38	0	34	1	0
4	B	38	0	34	3	0
All	All	6454	0	5973	216	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 216 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:191:ARG:HG2	1:B:191:ARG:HH11	1.08	1.16
1:B:214:GLN:NE2	1:B:214:GLN:H	1.61	0.98
1:B:348:GLU:HG2	1:B:350:ILE:HD12	1.48	0.94
1:B:155:GLN:HE21	1:B:155:GLN:H	0.94	0.92

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:174:GLY:H	1:B:177:HIS:HD2	1.12	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	387/419 (92%)	356 (92%)	23 (6%)	8 (2%)	9	23
1	B	400/419 (96%)	363 (91%)	30 (8%)	7 (2%)	11	27
All	All	787/838 (94%)	719 (91%)	53 (7%)	15 (2%)	10	25

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	264	VAL
1	B	2	GLU
1	B	386	PRO
1	A	218	GLU
1	A	366	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/379 (86%)	266 (81%)	61 (19%)	2	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	339/379 (89%)	279 (82%)	60 (18%)	2 5
All	All	666/758 (88%)	545 (82%)	121 (18%)	2 5

5 of 121 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	393	ARG
1	B	53	LEU
1	B	362	THR
1	A	403	TYR
1	B	6	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	404	ASN
1	B	42	ASN
1	B	328	GLN
1	B	18	HIS
1	B	64	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	1001	1,2	14,14,15	0.72	0	15,19,21	1.38	3 (20%)
2	FUC	A	1002	2	10,10,11	1.40	1 (10%)	14,14,16	2.38	7 (50%)
2	NAG	A	1003	2	14,14,15	0.77	1 (7%)	15,19,21	2.56	4 (26%)
2	BMA	A	1004	2	11,11,12	0.87	0	14,15,17	1.78	3 (21%)
3	NAG	A	2001	1,3	14,14,15	0.70	0	15,19,21	1.79	5 (33%)
3	FUC	A	2002	3	10,10,11	0.94	0	14,14,16	1.44	2 (14%)
3	NAG	A	2003	3	14,14,15	1.05	1 (7%)	15,19,21	2.60	4 (26%)
2	NAG	B	1001	1,2	14,14,15	0.56	0	15,19,21	1.96	4 (26%)
2	FUC	B	1002	2	10,10,11	1.18	1 (10%)	14,14,16	3.61	7 (50%)
2	NAG	B	1003	2	14,14,15	0.62	0	15,19,21	1.38	2 (13%)
2	BMA	B	1004	2	11,11,12	1.24	2 (18%)	14,15,17	2.37	3 (21%)
4	NDG	B	2001	1,4	14,14,15	1.42	2 (14%)	15,19,21	3.91	8 (53%)
4	FUC	B	2002	4	10,10,11	1.18	2 (20%)	14,14,16	2.53	5 (35%)
4	NAG	B	2003	4	14,14,15	0.89	1 (7%)	15,19,21	2.51	6 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1001	1,2	1/1/5/7	0/6/23/26	0/1/1/1
2	FUC	A	1002	2	-	0/0/17/20	0/1/1/1
2	NAG	A	1003	2	-	0/6/23/26	0/1/1/1
2	BMA	A	1004	2	-	0/2/19/22	0/1/1/1
3	NAG	A	2001	1,3	-	0/6/23/26	0/1/1/1
3	FUC	A	2002	3	-	0/0/17/20	0/1/1/1
3	NAG	A	2003	3	-	0/6/23/26	0/1/1/1
2	NAG	B	1001	1,2	-	0/6/23/26	0/1/1/1
2	FUC	B	1002	2	-	0/0/17/20	0/1/1/1
2	NAG	B	1003	2	-	0/6/23/26	0/1/1/1
2	BMA	B	1004	2	-	0/2/19/22	0/1/1/1
4	NDG	B	2001	1,4	-	0/6/23/26	0/1/1/1
4	FUC	B	2002	4	-	0/0/17/20	0/1/1/1
4	NAG	B	2003	4	-	2/6/23/26	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1003	NAG	O5-C1	-2.16	1.40	1.43
2	B	1004	BMA	C1-C2	2.05	1.57	1.52
4	B	2001	NDG	C3-C2	2.15	1.57	1.52
4	B	2003	NAG	C1-C2	2.17	1.55	1.52
4	B	2002	FUC	C1-C2	2.20	1.57	1.52

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1003	NAG	C2-N2-C7	-7.74	113.10	123.04
2	B	1002	FUC	C1-C2-C3	-7.35	100.85	109.54
2	A	1004	BMA	C1-C2-C3	-4.35	104.39	109.54
3	A	2003	NAG	C4-C3-C2	-3.96	105.07	111.23
3	A	2001	NAG	C3-C4-C5	-3.00	104.96	110.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1001	NAG	C1

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	2003	NAG	O7-C7-N2-C2
4	B	2003	NAG	C8-C7-N2-C2

There are no ring outliers.

10 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	NAG	1	0
2	A	1002	FUC	2	0
2	A	1003	NAG	1	0
3	A	2002	FUC	1	0
3	A	2003	NAG	1	0
2	B	1002	FUC	1	0
2	B	1003	NAG	1	0
4	B	2001	NDG	2	0
4	B	2002	FUC	2	0
4	B	2003	NAG	2	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	397/419 (94%)	0.34	15 (3%) 44 44	36, 50, 60, 69	0
1	B	406/419 (96%)	0.20	2 (0%) 91 93	35, 49, 58, 61	0
All	All	803/838 (95%)	0.27	17 (2%) 67 68	35, 50, 58, 69	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	295	PRO	3.8
1	A	264	VAL	3.3
1	A	416	TRP	3.3
1	A	373	LEU	3.1
1	A	325	PRO	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	B	1001	14/15	0.90	0.15	-1.42	72,77,87,87	0
2	NAG	A	1001	14/15	0.92	0.11	-1.95	42,52,64,66	0
2	BMA	A	1004	11/12	0.77	0.32	-	93,96,97,97	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	FUC	A	2002	10/11	0.74	0.25	-	95,97,99,99	0
3	NAG	A	2003	14/15	0.84	0.21	-	95,98,100,100	0
2	FUC	B	1002	10/11	0.66	0.44	-	93,95,97,97	0
2	FUC	A	1002	10/11	0.87	0.38	-	76,80,82,82	0
4	NAG	B	2003	14/15	0.82	0.17	-	94,96,97,98	0
2	NAG	A	1003	14/15	0.92	0.20	-	66,75,81,88	0
4	FUC	B	2002	10/11	0.56	0.24	-	96,100,100,101	0
4	NDG	B	2001	14/15	0.75	0.20	-	84,91,94,97	0
3	NAG	A	2001	14/15	0.89	0.18	-	84,91,96,96	0
2	BMA	B	1004	11/12	0.82	0.37	-	99,101,102,102	0
2	NAG	B	1003	14/15	0.87	0.23	-	93,95,99,99	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.