



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 18, 2017 – 07:48 AM EST

PDB ID : 4GYD  
Title : Nostoc sp Cytochrome c6  
Authors : Skubak, P.; Ubbink, M.; Cavazzini, D.; Rossi, G.L.; Pannu, N.S.  
Deposited on : 2012-09-05  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	86	<div> <div>2%</div> <div>90%</div> <div>9%</div> </div>
1	B	86	<div> <div>%</div> <div>88%</div> <div>10%</div> </div>
1	C	86	<div> <div>5%</div> <div>86%</div> <div>12%</div> </div>
1	D	86	<div> <div>9%</div> <div>85%</div> <div>15%</div> </div>
1	E	86	<div> <div>9%</div> <div>84%</div> <div>14%</div> </div>
1	F	86	<div> <div>12%</div> <div>86%</div> <div>13%</div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4598 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	86	Total	C	N	O	S	0	0	0
			640	401	112	123	4			
1	B	86	Total	C	N	O	S	0	0	0
			640	401	112	123	4			
1	C	86	Total	C	N	O	S	0	0	0
			640	401	112	123	4			
1	D	86	Total	C	N	O	S	0	0	0
			640	401	112	123	4			
1	E	86	Total	C	N	O	S	0	0	0
			640	401	112	123	4			
1	F	86	Total	C	N	O	S	0	0	0
			640	401	112	123	4			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	E	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	97	Total 97	O 97	0	0
3	B	100	Total 100	O 100	0	0
3	C	92	Total 92	O 92	0	0
3	D	66	Total 66	O 66	0	0
3	E	70	Total 70	O 70	0	0
3	F	75	Total 75	O 75	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cytochrome c6




- Molecule 1: Cytochrome c6




- Molecule 1: Cytochrome c6




- Molecule 1: Cytochrome c6



- Molecule 1: Cytochrome c6



- Molecule 1: Cytochrome c6





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.72Å 79.80Å 80.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.48 – 1.80 17.48 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (17.48-1.80) 99.7 (17.48-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.75 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.7.0027	Depositor
R, $R_{free}$	0.179 , 0.216 0.192 , 0.230	Depositor DCC
$R_{free}$ test set	2367 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.9	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.018 for -h,l,k 0.016 for -l,-k,-h 0.016 for k,h,-l 0.038 for k,l,h 0.038 for l,h,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4598	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 69.89 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.4264e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.03	1/649 (0.2%)	0.90	0/870
1	B	0.90	0/649	0.81	0/870
1	C	0.99	0/649	0.91	0/870
1	D	1.02	1/649 (0.2%)	0.87	0/870
1	E	1.03	2/649 (0.3%)	0.87	2/870 (0.2%)
1	F	1.06	1/649 (0.2%)	0.92	1/870 (0.1%)
All	All	1.01	5/3894 (0.1%)	0.88	3/5220 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	45	GLU	CD-OE2	5.70	1.31	1.25
1	E	71	GLU	CD-OE1	-5.70	1.19	1.25
1	E	68	GLU	CD-OE1	5.54	1.31	1.25
1	A	68	GLU	CD-OE1	5.54	1.31	1.25
1	F	58	MET	CG-SD	-5.43	1.67	1.81

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	64	ARG	NE-CZ-NH1	5.65	123.12	120.30
1	E	24	LEU	CA-CB-CG	5.60	128.18	115.30
1	F	84	ASP	CB-CG-OD1	-5.56	113.30	118.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	640	0	646	12	0
1	B	640	0	646	13	0
1	C	640	0	646	20	0
1	D	640	0	646	16	0
1	E	640	0	646	19	0
1	F	640	0	646	16	0
2	A	43	0	30	6	0
2	B	43	0	30	4	0
2	C	43	0	30	7	0
2	D	43	0	30	4	0
2	E	43	0	30	7	0
2	F	43	0	30	4	0
3	A	97	0	0	0	0
3	B	100	0	0	3	0
3	C	92	0	0	2	0
3	D	66	0	0	5	0
3	E	70	0	0	5	0
3	F	75	0	0	2	0
All	All	4598	0	4056	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:86:LYS:HG2	3:C:381:HOH:O	1.41	1.16
1:E:1:ALA:O	1:F:4:VAL:HG21	1.43	1.15
1:E:17:CYS:SG	2:E:200:HEM:HAC	1.99	1.00
1:C:17:CYS:SG	2:C:200:HEM:HAC	2.02	0.97
1:F:17:CYS:SG	2:F:200:HEM:HAC	2.02	0.96
1:B:17:CYS:SG	2:B:200:HEM:HAC	2.08	0.93
1:D:86:LYS:HA	3:D:360:HOH:O	1.71	0.91
1:A:17:CYS:SG	2:A:200:HEM:HAC	2.07	0.90
1:A:85:TRP:O	1:A:86:LYS:HB2	1.73	0.89

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:32:LYS:HG2	1:C:85:TRP:O	1.73	0.88
1:C:17:CYS:SG	2:C:200:HEM:CBC	2.61	0.88
1:D:17:CYS:SG	2:D:200:HEM:HAC	2.12	0.88
1:A:17:CYS:SG	2:A:200:HEM:CBC	2.63	0.85
1:B:17:CYS:SG	2:B:200:HEM:CBC	2.65	0.85
1:E:3:SER:HB2	3:E:354:HOH:O	1.75	0.85
1:C:17:CYS:SG	2:C:200:HEM:C3C	2.71	0.83
1:F:17:CYS:SG	2:F:200:HEM:C3C	2.72	0.83
1:E:17:CYS:SG	2:E:200:HEM:CBC	2.67	0.82
1:B:85:TRP:O	1:B:86:LYS:HB2	1.78	0.82
1:B:17:CYS:SG	2:B:200:HEM:C3C	2.73	0.82
1:E:17:CYS:SG	2:E:200:HEM:C3C	2.73	0.81
1:D:17:CYS:SG	2:D:200:HEM:CBC	2.70	0.79
1:C:17:CYS:HG	2:C:200:HEM:HAC	1.48	0.79
1:F:85:TRP:O	1:F:86:LYS:HB2	1.84	0.77
1:D:17:CYS:SG	2:D:200:HEM:C3C	2.78	0.77
1:A:17:CYS:SG	2:A:200:HEM:C3C	2.78	0.76
1:F:17:CYS:SG	2:F:200:HEM:CBC	2.73	0.75
1:C:17:CYS:HG	2:C:200:HEM:CAC	1.98	0.74
1:F:45:GLU:HG3	3:F:347:HOH:O	1.88	0.73
1:A:34:ALA:H	1:A:86:LYS:HG2	1.54	0.72
1:C:33:LYS:HB3	1:C:86:LYS:HD3	1.74	0.69
1:D:86:LYS:CA	3:D:360:HOH:O	2.35	0.67
1:F:26:GLN:HE21	1:F:28:GLN:HE22	1.47	0.62
1:E:1:ALA:C	1:F:4:VAL:HG21	2.20	0.62
1:B:32:LYS:HB3	1:B:86:LYS:HD2	1.83	0.60
1:F:80:LYS:HG2	1:F:85:TRP:CZ2	2.38	0.59
1:D:85:TRP:O	1:D:86:LYS:HB2	2.02	0.58
1:B:34:ALA:CB	1:B:86:LYS:HD3	2.35	0.56
1:B:68:GLU:HG3	3:B:314:HOH:O	2.07	0.55
1:C:34:ALA:H	1:C:86:LYS:NZ	2.05	0.55
1:E:85:TRP:O	1:E:86:LYS:HB2	2.06	0.55
1:A:32:LYS:HB3	1:A:86:LYS:HD3	1.87	0.55
1:B:34:ALA:HB2	1:B:86:LYS:HD3	1.89	0.55
1:B:85:TRP:O	1:B:86:LYS:CB	2.51	0.55
1:D:64:ARG:CD	3:D:353:HOH:O	2.55	0.54
1:F:32:LYS:HB3	1:F:86:LYS:HD3	1.88	0.54
1:C:40:GLY:HA3	1:D:8:LYS:HA	1.89	0.53
1:D:50:GLN:HE21	1:D:58:MET:HG2	1.73	0.52
1:B:64:ARG:NH2	3:B:346:HOH:O	2.42	0.51
1:D:80:LYS:HG2	1:D:85:TRP:CZ2	2.44	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:83:ALA:HB3	3:E:346:HOH:O	2.10	0.51
1:A:58:MET:HB2	2:A:200:HEM:C4D	2.46	0.51
1:E:50:GLN:HE21	1:E:58:MET:HG2	1.75	0.51
1:C:58:MET:HB2	2:C:200:HEM:C1D	2.46	0.50
1:E:32:LYS:HB3	1:E:86:LYS:HD3	1.93	0.50
1:D:86:LYS:N	3:D:360:HOH:O	2.41	0.50
1:A:17:CYS:HG	2:A:200:HEM:HAC	1.78	0.49
1:F:25:VAL:O	1:F:25:VAL:HG12	2.11	0.49
1:B:62:LYS:HE2	3:B:389:HOH:O	2.13	0.48
1:D:58:MET:HB2	2:D:200:HEM:C4D	2.49	0.48
1:C:34:ALA:CB	1:C:86:LYS:HZ3	2.27	0.47
1:B:58:MET:HB2	2:B:200:HEM:C1D	2.49	0.47
1:F:85:TRP:O	1:F:86:LYS:CB	2.58	0.47
1:E:17:CYS:HG	2:E:200:HEM:HAC	1.76	0.47
1:E:86:LYS:C	3:E:347:HOH:O	2.52	0.47
1:E:17:CYS:CB	2:E:200:HEM:C3C	2.98	0.47
1:D:4:VAL:HA	3:D:358:HOH:O	2.14	0.47
1:C:82:ASP:OD1	3:C:320:HOH:O	2.20	0.47
1:D:32:LYS:HB3	1:D:86:LYS:HD3	1.96	0.46
1:E:64:ARG:HD3	3:E:351:HOH:O	2.14	0.46
1:B:33:LYS:HB3	1:B:86:LYS:HG2	1.96	0.46
1:A:50:GLN:HE21	1:A:58:MET:HG2	1.81	0.45
1:E:58:MET:HB2	2:E:200:HEM:C4D	2.52	0.44
1:C:34:ALA:H	1:C:86:LYS:HZ3	1.63	0.44
1:D:34:ALA:H	1:D:86:LYS:HG2	1.83	0.44
1:F:26:GLN:NE2	1:F:28:GLN:HE22	2.14	0.44
1:E:86:LYS:HA	1:E:86:LYS:HD2	1.90	0.43
1:A:17:CYS:CB	2:A:200:HEM:C3C	3.02	0.43
1:E:1:ALA:C	3:E:363:HOH:O	2.56	0.43
1:C:86:LYS:HA	1:C:86:LYS:HD2	1.44	0.42
1:E:17:CYS:HB2	2:E:200:HEM:C3C	2.54	0.42
1:C:34:ALA:HB2	1:C:86:LYS:HZ3	1.85	0.42
1:A:34:ALA:N	1:A:86:LYS:HG2	2.29	0.41
1:D:86:LYS:HD2	1:D:86:LYS:HA	1.90	0.41
1:C:84:ASP:OD1	1:C:84:ASP:N	2.36	0.41
1:F:58:MET:HB2	2:F:200:HEM:C1D	2.55	0.41
1:C:17:CYS:CB	2:C:200:HEM:C3C	3.02	0.41
1:A:45:GLU:OE1	1:C:56:ASN:HB3	2.20	0.41
1:C:55:LYS:O	1:C:58:MET:HB3	2.21	0.41
1:E:34:ALA:H	1:E:86:LYS:HG2	1.86	0.40
1:F:86:LYS:HG3	3:F:369:HOH:O	2.20	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:80:LYS:HG3	1:F:85:TRP:CE2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	84/86 (98%)	81 (96%)	3 (4%)	0	100	100
1	B	84/86 (98%)	82 (98%)	2 (2%)	0	100	100
1	C	84/86 (98%)	80 (95%)	4 (5%)	0	100	100
1	D	84/86 (98%)	79 (94%)	5 (6%)	0	100	100
1	E	84/86 (98%)	82 (98%)	2 (2%)	0	100	100
1	F	84/86 (98%)	80 (95%)	4 (5%)	0	100	100
All	All	504/516 (98%)	484 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	63/63 (100%)	62 (98%)	1 (2%)	70	59
1	B	63/63 (100%)	62 (98%)	1 (2%)	70	59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	63/63 (100%)	61 (97%)	2 (3%)	46	29
1	D	63/63 (100%)	62 (98%)	1 (2%)	70	59
1	E	63/63 (100%)	62 (98%)	1 (2%)	70	59
1	F	63/63 (100%)	63 (100%)	0	100	100
All	All	378/378 (100%)	372 (98%)	6 (2%)	70	59

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	LYS
1	B	86	LYS
1	C	84	ASP
1	C	86	LYS
1	D	82	ASP
1	E	24	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	GLN
1	C	26	GLN
1	D	50	GLN
1	E	50	GLN
1	F	26	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	A	200	1	24,50,50	1.38	4 (16%)	16,82,82	2.04	6 (37%)
2	HEM	B	200	1	24,50,50	1.11	3 (12%)	16,82,82	2.06	4 (25%)
2	HEM	C	200	1	24,50,50	1.55	2 (8%)	16,82,82	1.77	3 (18%)
2	HEM	D	200	1	24,50,50	1.16	2 (8%)	16,82,82	2.38	7 (43%)
2	HEM	E	200	1	24,50,50	1.23	2 (8%)	16,82,82	2.28	7 (43%)
2	HEM	F	200	1	24,50,50	1.43	4 (16%)	16,82,82	2.10	6 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	200	1	-	0/6/54/54	0/0/8/8
2	HEM	B	200	1	-	0/6/54/54	0/0/8/8
2	HEM	C	200	1	-	0/6/54/54	0/0/8/8
2	HEM	D	200	1	-	0/6/54/54	0/0/8/8
2	HEM	E	200	1	-	0/6/54/54	0/0/8/8
2	HEM	F	200	1	-	0/6/54/54	0/0/8/8

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	200	HEM	C3B-C2B	-5.33	1.33	1.40
2	A	200	HEM	C3B-C2B	-4.30	1.34	1.40
2	F	200	HEM	C3C-C2C	-4.23	1.35	1.40
2	E	200	HEM	C3B-C2B	-3.91	1.35	1.40
2	C	200	HEM	C1B-NB	-3.74	1.31	1.36
2	D	200	HEM	C3B-C2B	-2.99	1.36	1.40
2	A	200	HEM	C4D-ND	-2.94	1.32	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	200	HEM	C1B-NB	-2.47	1.33	1.36
2	D	200	HEM	C1B-NB	-2.36	1.33	1.36
2	B	200	HEM	C3B-C2B	-2.28	1.37	1.40
2	A	200	HEM	CMC-C2C	-2.21	1.46	1.51
2	A	200	HEM	C4C-NC	-2.09	1.33	1.36
2	F	200	HEM	C3D-C2D	-2.09	1.31	1.37
2	B	200	HEM	CBB-CAB	2.01	1.43	1.28
2	E	200	HEM	CAD-C3D	2.12	1.55	1.52
2	F	200	HEM	CMB-C2B	2.12	1.56	1.51
2	B	200	HEM	CAD-C3D	2.34	1.55	1.52

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	200	HEM	C3B-CAB-CBB	-5.35	115.63	126.40
2	B	200	HEM	CBD-CAD-C3D	-4.92	103.84	112.47
2	F	200	HEM	C3B-CAB-CBB	-4.74	116.87	126.40
2	E	200	HEM	CMA-C3A-C4A	-4.65	120.40	128.31
2	C	200	HEM	C3B-CAB-CBB	-4.05	118.26	126.40
2	A	200	HEM	CBD-CAD-C3D	-4.00	105.46	112.47
2	B	200	HEM	C3C-CAC-CBC	-3.72	118.91	126.40
2	D	200	HEM	CMA-C3A-C4A	-3.65	122.10	128.31
2	A	200	HEM	C3B-CAB-CBB	-3.63	119.10	126.40
2	E	200	HEM	C3B-CAB-CBB	-3.63	119.10	126.40
2	C	200	HEM	CBD-CAD-C3D	-3.56	106.22	112.47
2	F	200	HEM	CBD-CAD-C3D	-3.36	106.57	112.47
2	D	200	HEM	CBD-CAD-C3D	-3.30	106.68	112.47
2	E	200	HEM	CBA-CAA-C2A	-3.16	106.94	112.49
2	B	200	HEM	CMA-C3A-C4A	-3.00	123.20	128.31
2	E	200	HEM	CBD-CAD-C3D	-2.76	107.63	112.47
2	D	200	HEM	C3C-CAC-CBC	-2.74	120.88	126.40
2	F	200	HEM	CMD-C2D-C1D	-2.60	123.89	128.31
2	A	200	HEM	CBA-CAA-C2A	-2.54	108.02	112.49
2	B	200	HEM	C3B-CAB-CBB	-2.53	121.31	126.40
2	F	200	HEM	C3C-CAC-CBC	-2.51	121.36	126.40
2	E	200	HEM	C3C-CAC-CBC	-2.29	121.79	126.40
2	A	200	HEM	CMD-C2D-C1D	-2.04	124.85	128.31
2	A	200	HEM	CAA-C2A-C3A	-2.03	123.19	129.00
2	C	200	HEM	CMD-C2D-C1D	-2.01	124.89	128.31
2	D	200	HEM	C3C-C4C-NC	-2.01	107.15	110.94
2	D	200	HEM	CMC-C2C-C3C	2.24	129.48	125.09
2	F	200	HEM	CAA-CBA-CGA	2.43	117.50	112.78

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	200	HEM	CMC-C2C-C3C	2.49	129.96	125.09
2	E	200	HEM	CMC-C2C-C3C	2.67	130.32	125.09
2	F	200	HEM	CMB-C2B-C3B	2.81	130.58	125.09
2	D	200	HEM	CAA-CBA-CGA	3.14	118.89	112.78
2	E	200	HEM	CMA-C3A-C2A	3.22	131.97	125.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 32 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	200	HEM	6	0
2	B	200	HEM	4	0
2	C	200	HEM	7	0
2	D	200	HEM	4	0
2	E	200	HEM	7	0
2	F	200	HEM	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	86/86 (100%)	-0.15	2 (2%) 64 59	8, 13, 25, 36	0
1	B	86/86 (100%)	-0.13	1 (1%) 81 78	8, 13, 23, 53	0
1	C	86/86 (100%)	0.02	4 (4%) 35 29	8, 15, 33, 37	0
1	D	86/86 (100%)	0.82	8 (9%) 11 8	12, 19, 57, 106	0
1	E	86/86 (100%)	0.36	8 (9%) 11 8	10, 17, 39, 76	0
1	F	86/86 (100%)	0.53	10 (11%) 6 5	10, 16, 38, 97	0
All	All	516/516 (100%)	0.24	33 (6%) 23 18	8, 15, 35, 106	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1	ALA	15.8
1	F	1	ALA	13.4
1	E	1	ALA	12.5
1	D	4	VAL	11.2
1	D	2	ASP	8.3
1	F	4	VAL	6.9
1	F	86	LYS	6.9
1	D	3	SER	6.7
1	F	2	ASP	6.6
1	D	5	ASN	6.1
1	E	86	LYS	6.1
1	D	86	LYS	5.5
1	F	85	TRP	5.3
1	C	63	GLY	5.1
1	E	3	SER	4.9
1	F	3	SER	4.5
1	F	84	ASP	3.9
1	F	63	GLY	3.9
1	B	86	LYS	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	82	ASP	3.6
1	E	63	GLY	3.2
1	E	4	VAL	3.1
1	E	82	ASP	2.9
1	C	1	ALA	2.9
1	D	63	GLY	2.7
1	E	85	TRP	2.4
1	E	2	ASP	2.4
1	A	86	LYS	2.3
1	C	86	LYS	2.2
1	F	62	LYS	2.2
1	C	62	LYS	2.1
1	F	83	ALA	2.0
1	A	24	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	HEM	A	200	43/43	0.95	0.11	0.23	7,9,13,17	0
2	HEM	B	200	43/43	0.96	0.09	-0.05	7,8,11,15	0
2	HEM	F	200	43/43	0.96	0.10	-0.17	8,10,11,12	0
2	HEM	D	200	43/43	0.95	0.10	-0.21	10,12,15,16	0
2	HEM	C	200	43/43	0.96	0.09	-0.22	8,9,12,20	0
2	HEM	E	200	43/43	0.96	0.10	-0.25	8,10,15,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.