



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:03 AM GMT

PDB ID : 3H10
Title : Aurora A inhibitor complex
Authors : Wiesmann, C.; Ultsch, M.H.; Cochran, A.G.
Deposited on : 2009-04-10
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	<div> <div>3%</div> <div>85%</div> <div>12%</div> <div>.</div> </div>
1	B	268	<div> <div>3%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>
1	D	268	<div> <div>%</div> <div>84%</div> <div>9%</div> <div>5%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	0	0
			2191	1407	389	389	6			
1	B	267	Total	C	N	O	S	0	0	0
			2186	1404	388	388	6			
1	D	255	Total	C	N	O	S	0	0	0
			2099	1352	368	374	5			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	ALA	LYS	ENGINEERED	UNP O14965
A	287	ALA	THR	ENGINEERED	UNP O14965
A	288	ALA	THR	ENGINEERED	UNP O14965
B	124	ALA	LYS	ENGINEERED	UNP O14965
B	287	ALA	THR	ENGINEERED	UNP O14965
B	288	ALA	THR	ENGINEERED	UNP O14965
D	124	ALA	LYS	ENGINEERED	UNP O14965
D	287	ALA	THR	ENGINEERED	UNP O14965
D	288	ALA	THR	ENGINEERED	UNP O14965

- Molecule 2 is 9-CHLORO-7-(2,6-DIFLUOROPHENYL)-N-{4-[(4-METHYLPIPERAZIN-1-YL)CARBONYL]PHENYL}-5H-PYRIMIDO[5,4-D][2]BENZAZEPIN-2-AMINE (three-letter code: 97B) (formula: C₃₀H₂₅ClF₂N₆O).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	0	0
			40	30	1	2	6	1		
2	B	1	Total	C	Cl	F	N	O	0	0
			40	30	1	2	6	1		
2	D	1	Total	C	Cl	F	N	O	0	0
			40	30	1	2	6	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	60	Total	O	0	0
			60	60		
3	B	45	Total	O	0	0
			45	45		
3	D	42	Total	O	0	0
			42	42		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase 6



- Molecule 1: Serine/threonine-protein kinase 6



- Molecule 1: Serine/threonine-protein kinase 6



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.75Å 88.98Å 122.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20 43.88 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-2.20) 99.7 (43.88-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.41 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.194 , 0.240 0.198 , 0.240	Depositor DCC
R_{free} test set	2482 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	40.4	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.6	EDS
Estimated twinning fraction	0.089 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 49153 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6743	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 97B

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2246	0.79	3/3035 (0.1%)
1	B	0.60	0/2241	0.72	1/3028 (0.0%)
1	D	0.61	0/2152	0.71	1/2909 (0.0%)
All	All	0.63	0/6639	0.74	5/8972 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	270	LEU	CA-CB-CG	7.62	132.83	115.30
1	B	270	LEU	CA-CB-CG	7.56	132.68	115.30
1	A	286	ARG	NE-CZ-NH1	5.33	122.96	120.30
1	D	232	ARG	NE-CZ-NH2	-5.29	117.65	120.30
1	A	151	ARG	NE-CZ-NH2	-5.06	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2191	0	2204	28	0
1	B	2186	0	2199	25	0
1	D	2099	0	2100	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	40	0	25	3	0
2	B	40	0	25	1	0
2	D	40	0	25	1	0
3	A	60	0	0	0	0
3	B	45	0	0	2	0
3	D	42	0	0	3	1
All	All	6743	0	6578	73	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:375:ARG:HD3	1:A:375:ARG:H	1.41	0.83
1:D:292:THR:HG23	3:D:85:HOH:O	1.80	0.82
1:D:224:LYS:HG2	1:D:225:LEU:HD13	1.67	0.78
1:D:292:THR:CG2	3:D:85:HOH:O	2.33	0.76
1:B:174:VAL:HG13	1:B:177:GLN:HE21	1.51	0.75
1:A:375:ARG:H	1:A:375:ARG:CD	1.99	0.73
1:D:174:VAL:HG11	1:D:277:TRP:CZ3	2.24	0.73
1:A:221:GLU:HG2	1:A:232:ARG:NH2	2.05	0.70
1:B:174:VAL:HG12	1:B:174:VAL:O	1.92	0.69
1:A:358:ASP:OD2	1:A:380:HIS:HE1	1.78	0.67
1:B:174:VAL:HG11	1:B:277:TRP:CE3	2.30	0.67
1:A:190:HIS:HD2	1:A:192:ASN:H	1.47	0.63
1:D:174:VAL:O	1:D:174:VAL:HG12	1.98	0.63
1:A:335:GLN:HB3	1:B:338:TYR:OH	1.98	0.62
1:B:144:PHE:HD2	1:B:168:GLN:HG3	1.64	0.62
1:A:375:ARG:HD3	1:A:375:ARG:N	2.15	0.62
1:B:262:LEU:HD11	1:B:318:LEU:HD11	1.83	0.61
2:A:1:97B:H19	2:A:1:97B:N3	2.16	0.61
1:B:190:HIS:HD2	1:B:192:ASN:H	1.49	0.60
1:B:174:VAL:HG13	1:B:177:GLN:NE2	2.15	0.59
1:A:190:HIS:CD2	1:A:192:ASN:H	2.21	0.59
1:D:339:LYS:NZ	3:D:91:HOH:O	2.37	0.58
2:D:3:97B:H19	2:D:3:97B:N3	2.19	0.58
1:A:307:ASP:O	1:A:310:VAL:HG22	2.04	0.57
1:A:135:ILE:CD1	1:A:161:LEU:HD12	2.34	0.57
2:B:2:97B:H19	2:B:2:97B:N3	2.20	0.57
1:B:190:HIS:CD2	1:B:192:ASN:H	2.23	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:ARG:HH12	2:A:1:97B:H22	1.71	0.56
1:B:217:THR:OG1	1:B:220:ARG:HG2	2.06	0.55
1:D:174:VAL:HG13	1:D:177:GLN:NE2	2.21	0.55
1:A:375:ARG:CD	1:A:375:ARG:N	2.68	0.54
1:A:221:GLU:O	1:A:225:LEU:HD13	2.07	0.54
1:B:293:LEU:HG	1:B:337:THR:HG21	1.90	0.53
1:B:174:VAL:HG11	1:B:277:TRP:CZ3	2.44	0.52
1:B:256:ASP:OD1	1:B:258:LYS:HE3	2.08	0.52
1:B:256:ASP:CG	1:B:261:ASN:HD22	2.13	0.52
1:B:389:LYS:HE3	3:B:104:HOH:O	2.09	0.51
1:A:224:LYS:HB3	1:A:225:LEU:HD12	1.93	0.51
1:B:380:HIS:CD2	1:B:382:TRP:H	2.28	0.51
1:B:181:GLU:HB3	1:B:275:PHE:CE2	2.46	0.51
1:A:135:ILE:HD12	1:A:161:LEU:HD12	1.93	0.49
1:B:144:PHE:CD2	1:B:168:GLN:HG3	2.47	0.49
1:A:221:GLU:HG2	1:A:232:ARG:HH22	1.78	0.48
1:A:240:LEU:CD1	1:A:318:LEU:CD1	2.91	0.48
1:A:274:ASP:HB2	1:A:275:PHE:CD1	2.47	0.48
1:D:240:LEU:HG	1:D:270:LEU:HD21	1.95	0.48
1:B:282:PRO:O	1:B:284:SER:N	2.46	0.48
1:A:317:VAL:HG13	1:A:328:PRO:HD2	1.95	0.48
1:B:335:GLN:HG2	3:B:113:HOH:O	2.12	0.48
1:B:307:ASP:O	1:B:310:VAL:HG22	2.13	0.48
1:D:380:HIS:CD2	1:D:382:TRP:H	2.33	0.47
1:B:231:GLN:NE2	1:B:386:ASN:OD1	2.47	0.47
1:D:190:HIS:HD2	1:D:192:ASN:H	1.61	0.47
1:D:307:ASP:O	1:D:310:VAL:HG22	2.16	0.46
1:D:190:HIS:CD2	1:D:192:ASN:H	2.34	0.46
1:A:338:TYR:OH	1:B:335:GLN:HA	2.16	0.46
1:B:174:VAL:O	1:B:174:VAL:CG1	2.61	0.45
1:D:295:TYR:OH	1:D:318:LEU:HD23	2.18	0.43
1:D:161:LEU:HD13	1:D:163:VAL:HG22	2.00	0.43
1:A:233:THR:HG21	1:A:323:LEU:HG	2.01	0.42
1:B:353:THR:O	1:B:357:ARG:HG3	2.19	0.42
1:A:289:LEU:HD23	1:A:289:LEU:C	2.40	0.42
1:D:221:GLU:HG2	1:D:232:ARG:NH2	2.35	0.42
1:A:380:HIS:CD2	1:A:382:TRP:H	2.38	0.42
1:D:240:LEU:HD13	1:D:318:LEU:HD12	2.03	0.41
1:A:260:GLU:OE2	1:A:286:ARG:CD	2.68	0.41
1:A:137:ARG:NH1	2:A:1:97B:H22	2.33	0.41
1:D:318:LEU:HD13	1:D:322:PHE:CE2	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:174:VAL:O	1:D:174:VAL:CG1	2.68	0.41
1:D:190:HIS:CG	1:D:191:PRO:HD2	2.56	0.41
1:A:165:PHE:CZ	1:D:303:GLY:HA2	2.55	0.40
1:A:130:LEU:HD13	1:A:207:TYR:CE2	2.56	0.40
1:A:240:LEU:HD13	1:A:318:LEU:HD12	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:5:HOH:O	3:D:103:HOH:O 4_544	2.03	0.17

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/268 (99%)	254 (96%)	9 (3%)	3 (1%)	17	14
1	B	265/268 (99%)	255 (96%)	9 (3%)	1 (0%)	39	42
1	D	251/268 (94%)	247 (98%)	4 (2%)	0	100	100
All	All	782/804 (97%)	756 (97%)	22 (3%)	4 (0%)	34	35

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	283	SER
1	B	283	SER
1	A	284	SER
1	A	186	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/234 (100%)	225 (96%)	9 (4%)	40	49
1	B	234/234 (100%)	220 (94%)	14 (6%)	24	26
1	D	225/234 (96%)	215 (96%)	10 (4%)	35	42
All	All	693/702 (99%)	660 (95%)	33 (5%)	31	37

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	GLN
1	A	161	LEU
1	A	253	ILE
1	A	270	LEU
1	A	285	ARG
1	A	286	ARG
1	A	318	LEU
1	A	335	GLN
1	A	375	ARG
1	B	161	LEU
1	B	175	GLU
1	B	181	GLU
1	B	185	GLN
1	B	189	ARG
1	B	225	LEU
1	B	270	LEU
1	B	283	SER
1	B	304	ARG
1	B	318	LEU
1	B	326	LYS
1	B	369	SER
1	B	375	ARG
1	B	388	SER
1	D	137	ARG
1	D	161	LEU
1	D	168	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	175	GLU
1	D	189	ARG
1	D	225	LEU
1	D	270	LEU
1	D	308	GLU
1	D	318	LEU
1	D	375	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	GLN
1	A	190	HIS
1	A	192	ASN
1	A	242	ASN
1	A	254	HIS
1	A	306	HIS
1	A	332	ASN
1	A	366	HIS
1	A	380	HIS
1	B	177	GLN
1	B	190	HIS
1	B	192	ASN
1	B	231	GLN
1	B	242	ASN
1	B	261	ASN
1	B	306	HIS
1	B	366	HIS
1	B	380	HIS
1	B	386	ASN
1	D	177	GLN
1	D	190	HIS
1	D	192	ASN
1	D	242	ASN
1	D	306	HIS
1	D	332	ASN
1	D	366	HIS
1	D	380	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	97B	A	1	-	43,45,45	0.81	2 (4%)	61,65,65	2.03	11 (18%)
2	97B	B	2	-	43,45,45	0.79	2 (4%)	61,65,65	2.17	13 (21%)
2	97B	D	3	-	43,45,45	0.82	1 (2%)	61,65,65	2.17	13 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	97B	A	1	-	-	0/16/39/39	0/5/6/6
2	97B	B	2	-	-	0/16/39/39	0/5/6/6
2	97B	D	3	-	-	0/16/39/39	0/5/6/6

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	97B	C10-C9	-2.38	1.38	1.41
2	B	2	97B	C14-N4	-2.19	1.36	1.40
2	A	1	97B	C10-C9	-2.19	1.38	1.41
2	D	3	97B	C10-C9	-2.14	1.38	1.41
2	A	1	97B	C14-N4	-2.03	1.36	1.40

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	97B	N2-C13-N3	-7.55	118.62	126.67
2	D	3	97B	N2-C13-N3	-7.50	118.68	126.67
2	A	1	97B	N2-C13-N3	-6.35	119.91	126.67
2	D	3	97B	C26-C11-C7	-5.35	116.93	122.69
2	B	2	97B	C9-C10-N3	-3.46	119.21	122.86
2	A	1	97B	C9-C10-N3	-3.45	119.23	122.86
2	B	2	97B	C26-C11-C7	-3.43	119.00	122.69
2	A	1	97B	C26-C11-C7	-3.26	119.18	122.69
2	B	2	97B	C4-C3-C10	-2.72	115.30	119.12
2	A	1	97B	C4-C3-C10	-2.63	115.43	119.12
2	D	3	97B	C24-C23-N6	-2.51	108.13	110.79
2	D	3	97B	C9-C12-N2	-2.45	119.60	123.86
2	D	3	97B	C4-C3-C10	-2.35	115.83	119.12
2	D	3	97B	C27-C26-C11	-2.31	119.42	123.42
2	B	2	97B	C29-C30-C11	-2.20	119.61	123.42
2	D	3	97B	C9-C10-N3	-2.18	120.56	122.86
2	A	1	97B	C2-C7-C11	-2.13	115.60	118.08
2	B	2	97B	O1-C20-N5	-2.13	119.05	122.42
2	A	1	97B	O1-C20-N5	-2.13	119.05	122.42
2	D	3	97B	C29-C30-C11	-2.03	119.90	123.42
2	A	1	97B	C12-N2-C13	2.14	120.04	115.95
2	D	3	97B	F1-C30-C11	2.18	121.38	118.09
2	B	2	97B	C4-C3-C2	2.26	120.62	118.03
2	B	2	97B	C21-N5-C24	2.27	116.77	112.56
2	B	2	97B	C11-C7-N1	2.47	120.73	116.86
2	B	2	97B	C12-N2-C13	2.66	121.03	115.95
2	A	1	97B	C30-C11-C26	2.87	119.13	115.08
2	A	1	97B	C11-C7-N1	2.97	121.52	116.86
2	D	3	97B	C12-N2-C13	3.23	122.13	115.95
2	B	2	97B	C30-C11-C26	3.32	119.78	115.08
2	D	3	97B	C30-C11-C26	3.83	120.50	115.08
2	A	1	97B	C10-N3-C13	5.53	121.33	116.66
2	D	3	97B	C10-N3-C13	5.64	121.43	116.66
2	B	2	97B	C10-N3-C13	6.59	122.22	116.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	97B	C8-N1-C7	8.50	127.82	117.54
2	D	3	97B	C8-N1-C7	8.54	127.86	117.54
2	B	2	97B	C8-N1-C7	8.59	127.92	117.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	97B	3	0
2	B	2	97B	1	0
2	D	3	97B	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/268 (100%)	-0.00	8 (2%) 54 53	26, 42, 65, 97	0
1	B	267/268 (99%)	-0.03	7 (2%) 59 58	31, 44, 64, 102	0
1	D	255/268 (95%)	-0.18	2 (0%) 87 87	29, 39, 55, 90	0
All	All	790/804 (98%)	-0.07	17 (2%) 65 64	26, 42, 64, 102	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	283	SER	5.9
1	A	290	CYS	5.5
1	A	289	LEU	4.5
1	A	284	SER	4.2
1	A	283	SER	3.9
1	B	338	TYR	3.7
1	B	284	SER	3.6
1	B	289	LEU	3.2
1	A	338	TYR	3.1
1	B	290	CYS	3.0
1	D	127	GLN	3.0
1	A	179	ARG	2.6
1	A	291	GLY	2.6
1	A	304	ARG	2.3
1	B	301	ILE	2.2
1	D	126	ARG	2.1
1	B	285	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	97B	B	2	40/40	0.95	0.15	0.92	28,37,78,79	0
2	97B	D	3	40/40	0.94	0.12	-0.08	26,36,46,47	0
2	97B	A	1	40/40	0.95	0.12	-0.42	18,29,77,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.