



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:25 AM GMT

PDB ID : 2H2R  
Title : Crystal structure of the human CD23 Lectin domain, apo form  
Authors : Wurzburg, B.A.  
Deposited on : 2006-05-19  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	175	
1	B	175	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2553 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Low affinity immunoglobulin epsilon Fc receptor (Lymphocyte IgE receptor) (Fc-epsilon-RII)(Immunoglobulin E-binding factor) (CD23 antigen).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	135	Total	C	N	O	S	0	2	0
			1096	690	194	200	12			
1	B	134	Total	C	N	O	S	0	3	0
			1097	691	194	200	12			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	SER	-	CLONING ARTIFACT	UNP P06734
A	148	PRO	-	CLONING ARTIFACT	UNP P06734
A	149	GLY	-	CLONING ARTIFACT	UNP P06734
A	213	ARG	HIS	ENGINEERED	UNP P06734
A	256	SER	GLY	ENGINEERED	UNP P06734
B	147	SER	-	CLONING ARTIFACT	UNP P06734
B	148	PRO	-	CLONING ARTIFACT	UNP P06734
B	149	GLY	-	CLONING ARTIFACT	UNP P06734
B	213	ARG	HIS	ENGINEERED	UNP P06734
B	256	SER	GLY	ENGINEERED	UNP P06734


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	171	Total	O	0	0
			171	171		
2	B	189	Total	O	0	0
			189	189		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Low affinity immunoglobulin epsilon Fc receptor (Lymphocyte IgE receptor) (Fc-epsilon-RII)(Immunoglobulin E-binding factor) (CD23 antigen)



- Molecule 1: Low affinity immunoglobulin epsilon Fc receptor (Lymphocyte IgE receptor) (Fc-epsilon-RII)(Immunoglobulin E-binding factor) (CD23 antigen)



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.35Å 50.67Å 75.06Å 90.00° 127.09° 90.00°	Depositor
Resolution (Å)	59.87 – 1.50 28.55 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (59.87-1.50) 99.9 (28.55-1.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.09 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.124 , 0.165 0.125 , 0.167	Depositor DCC
$R_{free}$ test set	2727 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.7	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 52.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 53946 reflections	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	2553	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.11	5/1135 (0.4%)	0.82	2/1537 (0.1%)
1	B	0.96	0/1139	0.76	1/1542 (0.1%)
All	All	1.04	5/2274 (0.2%)	0.79	3/3079 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	280	TRP	CZ3-CH2	8.73	1.54	1.40
1	A	280	TRP	CE3-CZ3	-8.54	1.24	1.38
1	A	224	ARG	CB-CG	-8.24	1.30	1.52
1	A	270	ASP	CB-CG	-5.41	1.40	1.51
1	A	235	VAL	CB-CG2	-5.05	1.42	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	267	ARG	NE-CZ-NH1	8.17	124.38	120.30
1	A	192	ASP	CB-CG-OD1	6.34	124.00	118.30
1	B	213	ARG	NE-CZ-NH1	6.15	123.38	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1096	0	1024	17	1
1	B	1097	0	1027	12	1
2	A	171	0	0	3	0
2	B	189	0	0	9	0
All	All	2553	0	2051	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:ASP:HB2	2:B:461:HOH:O	1.78	0.83
1:A:230:GLY:O	2:A:492:HOH:O	2.04	0.75
1:A:226:LEU:HD23	1:A:228:LEU:HB2	1.71	0.72
1:A:229:LYS:CD	1:A:233:ILE:HD12	2.22	0.69
1:A:230:GLY:HA2	2:A:492:HOH:O	1.93	0.68
1:B:227:ASP:OD2	2:B:478:HOH:O	2.13	0.65
1:B:197[A]:GLN:HG2	2:B:501:HOH:O	2.00	0.62
1:B:197[A]:GLN:HG3	2:B:351:HOH:O	2.00	0.61
1:A:165:GLU:HG2	1:A:166:LYS:HG3	1.85	0.58
1:A:229:LYS:HD2	1:A:233:ILE:HD12	1.85	0.57
1:B:168[A]:ILE:HD13	2:B:381:HOH:O	2.03	0.56
1:B:200:SER:HA	1:B:234:TRP:CE3	2.44	0.53
1:A:229:LYS:HD3	1:A:233:ILE:HD12	1.90	0.52
1:B:252:SER:OG	2:B:440:HOH:O	2.19	0.51
1:A:225:ASN:HB2	1:A:232:PHE:CE2	2.45	0.51
1:B:208:ASP:CB	2:B:461:HOH:O	2.48	0.50
1:A:229:LYS:HD3	1:A:233:ILE:CD1	2.43	0.49
1:A:254:SER:HB3	1:A:258:ASP:OD1	2.13	0.48
1:A:258:ASP:OD2	1:A:270:ASP:OD2	2.31	0.47
1:B:194[A]:MET:HB2	1:B:194[A]:MET:HE2	1.78	0.46
1:B:195:GLU:HG2	2:B:484:HOH:O	2.15	0.45
1:A:188:ARG:NE	2:A:365:HOH:O	2.48	0.45
1:A:227:ASP:OD1	1:A:228:LEU:N	2.50	0.45
1:B:158:PHE:N	2:B:410:HOH:O	2.49	0.45
1:B:225:ASN:HB2	1:B:232:PHE:CE2	2.51	0.45
1:A:254:SER:OG	1:A:256:SER:N	2.42	0.44
1:A:176:TYR:CD2	1:A:194[A]:MET:HE1	2.53	0.44
1:A:225:ASN:HB2	1:A:232:PHE:CD2	2.54	0.42
1:A:200:SER:HA	1:A:234:TRP:CE3	2.55	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:ARG:NH1	1:B:257:GLU:OE1[2_657]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/175 (77%)	132 (98%)	3 (2%)	0	100	100
1	B	135/175 (77%)	130 (96%)	5 (4%)	0	100	100
All	All	270/350 (77%)	262 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/149 (79%)	113 (96%)	5 (4%)	36	7
1	B	119/149 (80%)	117 (98%)	2 (2%)	68	37
All	All	237/298 (80%)	230 (97%)	7 (3%)	48	15

All (7) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	A	159	VAL
1	A	224	ARG
1	A	226	LEU
1	A	254	SER
1	A	258	ASP
1	B	225	ASN
1	B	241	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	171	GLN
1	B	225	ASN
1	B	255	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	135/175 (77%)	0.35	12 (8%) 12 11	13, 22, 50, 74	0
1	B	134/175 (76%)	0.08	7 (5%) 31 32	13, 19, 39, 55	0
All	All	269/350 (76%)	0.21	19 (7%) 19 19	13, 20, 41, 74	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	292	ALA	8.7
1	B	291	PRO	7.2
1	B	158	PHE	7.2
1	A	158	PHE	6.5
1	A	228	LEU	6.4
1	A	291	PRO	6.0
1	A	229	LYS	5.5
1	B	228	LEU	5.4
1	A	230	GLY	4.4
1	B	230	GLY	4.3
1	A	226	LEU	4.0
1	B	255	GLN	3.8
1	A	255	GLN	3.6
1	A	216	HIS	3.4
1	A	227	ASP	3.1
1	B	229	LYS	2.9
1	A	221	ILE	2.5
1	B	216	HIS	2.1
1	A	159	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

### 6.5 Other polymers

There are no such residues in this entry.