



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:27 AM GMT

PDB ID : 2H7O
Title : Crystal structure of the Rho-GTPase binding domain of YpkA
Authors : Prehna, G.; Ivanov, M.; Bliska, J.B.; Stebbins, C.E.
Deposited on : 2006-06-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	<div> <div>11%</div> <div> <div></div> <div>74%</div> <div>12%</div> <div>••</div> <div>11%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein kinase ypkA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	0	0
			2169	1356	389	417	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	430	GLY	-	CLONING ARTIFACT	UNP Q05608
A	431	PRO	-	CLONING ARTIFACT	UNP Q05608
A	432	VAL	-	CLONING ARTIFACT	UNP Q05608
A	433	ASP	-	CLONING ARTIFACT	UNP Q05608


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	276	Total	O	0	0
			276	276		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein kinase ypkA



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	60.01Å 60.01Å 402.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.88 – 2.00 33.88 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.5 (33.88-2.00) 98.5 (33.88-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.208 , 0.237 0.272 , 0.298	Depositor DCC
R_{free} test set	1485 reflections (5.20%)	DCC
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 30024 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2445	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	6/2194 (0.3%)	0.92	6/2953 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	725	ASN	CB-CG	7.49	1.68	1.51
1	A	443	LYS	CD-CE	6.50	1.67	1.51
1	A	709	GLU	CG-CD	6.05	1.61	1.51
1	A	589	GLU	CB-CG	-5.26	1.42	1.52
1	A	678	GLU	CD-OE2	5.13	1.31	1.25
1	A	621	GLN	CG-CD	5.02	1.62	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	715	ARG	NE-CZ-NH1	6.11	123.36	120.30
1	A	642	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	597	LEU	CA-CB-CG	5.60	128.17	115.30
1	A	579	LEU	CA-CB-CG	5.30	127.49	115.30
1	A	684	ARG	NE-CZ-NH2	-5.04	117.78	120.30
1	A	453	ARG	NE-CZ-NH2	-5.01	117.80	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	703	MET	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2169	0	2212	36	0
2	A	276	0	0	17	2
All	All	2445	0	2212	36	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:702:LEU:O	1:A:704:ASP:N	2.09	0.85
1:A:628:ARG:HD2	2:A:801:HOH:O	1.81	0.78
1:A:682:ASP:OD2	2:A:996:HOH:O	2.06	0.74
1:A:598:ALA:O	1:A:602:ILE:HG23	1.88	0.74
1:A:612:GLN:NE2	2:A:992:HOH:O	2.08	0.73
1:A:709:GLU:HG3	2:A:896:HOH:O	1.92	0.69
1:A:619:LYS:NZ	2:A:915:HOH:O	2.31	0.62
1:A:495:LYS:HG3	2:A:899:HOH:O	1.99	0.62
1:A:491:LYS:HD2	2:A:899:HOH:O	1.99	0.61
1:A:646:THR:OG1	1:A:668:HIS:NE2	2.25	0.61
1:A:621:GLN:CG	2:A:871:HOH:O	2.49	0.60
1:A:503:LYS:NZ	2:A:998:HOH:O	2.36	0.56
1:A:719:THR:HG22	2:A:999:HOH:O	2.06	0.56
1:A:647:ARG:HA	1:A:668:HIS:CE1	2.42	0.53
1:A:702:LEU:HB2	1:A:708:VAL:CG2	2.38	0.53
1:A:621:GLN:HG3	2:A:871:HOH:O	2.09	0.52
1:A:702:LEU:HB2	1:A:708:VAL:HG23	1.92	0.51
1:A:661:HIS:HB3	2:A:827:HOH:O	2.13	0.49
1:A:445:LEU:HD22	1:A:489:VAL:HG11	1.95	0.48
1:A:617:SER:O	1:A:621:GLN:HG3	2.16	0.46
1:A:715:ARG:O	1:A:719:THR:HG23	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:727:LEU:HD22	1:A:731:TRP:CD1	2.51	0.46
1:A:709:GLU:CG	2:A:896:HOH:O	2.58	0.46
1:A:663:GLN:HG2	2:A:815:HOH:O	2.15	0.46
1:A:495:LYS:CG	2:A:899:HOH:O	2.60	0.45
1:A:663:GLN:CG	2:A:815:HOH:O	2.65	0.45
1:A:674:GLN:HA	1:A:674:GLN:HE21	1.82	0.44
1:A:702:LEU:HD13	1:A:707:LEU:HD23	2.00	0.44
1:A:576:LEU:HD12	1:A:579:LEU:HD22	2.00	0.44
1:A:602:ILE:HG13	1:A:603:THR:N	2.29	0.43
1:A:674:GLN:NE2	2:A:890:HOH:O	2.51	0.43
1:A:489:VAL:HG11	1:A:494:LEU:HD13	1.99	0.43
1:A:702:LEU:HD13	1:A:707:LEU:CD2	2.50	0.42
1:A:703:MET:SD	1:A:703:MET:N	2.93	0.41
1:A:727:LEU:HD22	1:A:731:TRP:HD1	1.86	0.41
1:A:674:GLN:NE2	1:A:674:GLN:HA	2.36	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:991:HOH:O	2:A:997:HOH:O[12_564]	1.93	0.27
2:A:753:HOH:O	2:A:1002:HOH:O[8_665]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	262/303 (86%)	261 (100%)	0	1 (0%)	39 33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	703	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	245/273 (90%)	235 (96%)	10 (4%)	37 32

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	458	SER
1	A	462	LYS
1	A	491	LYS
1	A	494	LEU
1	A	506	ARG
1	A	561	SER
1	A	579	LEU
1	A	602	ILE
1	A	703	MET
1	A	727	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	493	GLN
1	A	533	ASN
1	A	606	GLN
1	A	674	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/303 (89%)	0.78	33 (12%) 5 6	32, 45, 59, 76	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	703	MET	6.3
1	A	514	GLY	5.5
1	A	512	VAL	5.3
1	A	492	ASP	4.7
1	A	585	PRO	4.7
1	A	502	LEU	4.5
1	A	581	GLN	4.2
1	A	546	ILE	4.1
1	A	510	ASP	3.7
1	A	511	TYR	3.6
1	A	495	LYS	3.3
1	A	513	LYS	3.3
1	A	506	ARG	3.2
1	A	491	LYS	3.2
1	A	569	LEU	3.2
1	A	489	VAL	3.0
1	A	509	GLU	3.0
1	A	543	LEU	2.9
1	A	451	LEU	2.8
1	A	562	LEU	2.6
1	A	678	GLU	2.6
1	A	597	LEU	2.5
1	A	572	LEU	2.5
1	A	565	ALA	2.5
1	A	490	ASP	2.4
1	A	680	THR	2.3
1	A	681	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	473	LEU	2.3
1	A	697	LEU	2.3
1	A	568	HIS	2.2
1	A	729	ARG	2.2
1	A	452	LEU	2.1
1	A	664	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.