



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 3, 2017 – 01:18 PM EST

PDB ID : 5H8U  
Title : Crystal structure of mycobacterium tuberculosis wild-type malate synthase in complex with product malate  
Authors : Krieger, I.V.; Huang, H.-L.; Sacchettini, J.C.  
Deposited on : 2015-12-23  
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2228 (2.90-2.82)
Clashscore	102246	2499 (2.90-2.82)
Ramachandran outliers	100387	2439 (2.90-2.82)
Sidechain outliers	100360	2442 (2.90-2.82)
RSRZ outliers	91569	2236 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	741	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>••</div> </div> </div>
1	B	741	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>16%</div> <div>••</div> </div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 11065 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Malate synthase G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	720	Total	C	N	O	S	0	0	0
			5505	3458	969	1056	22			
1	B	716	Total	C	N	O	S	0	0	0
			5483	3446	965	1050	22			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is (2S)-2-hydroxybutanedioic acid (three-letter code: LMR) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			9	4	5		

- Molecule 4 is GLYOXYLIC ACID (three-letter code: GLV) (formula:  $C_2H_2O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			5	2	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	31	Total	O	0	0
			31	31		
5	B	29	Total	O	0	0
			29	29		


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Malate synthase G



#### • Molecule 1: Malate synthase G





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.66Å 120.66Å 232.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.36 – 2.85 48.93 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.0 (37.36-2.85) 85.5 (48.93-2.29)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.89 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.226 , 0.276 0.226 , 0.276	Depositor DCC
$R_{free}$ test set	2045 reflections (5.11%)	DCC
Wilson B-factor (Å <sup>2</sup> )	77.4	Xtriage
Anisotropy	0.256	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 37.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	11065	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LMR, MG, GLV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.21	0/5610	0.39	0/7631
1	B	0.20	0/5587	0.39	0/7598
All	All	0.21	0/11197	0.39	0/15229

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5505	0	5450	61	0
1	B	5483	0	5432	67	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
3	A	9	0	3	2	0
4	B	5	0	1	0	0
5	A	31	0	0	4	0
5	B	29	0	0	3	0
All	All	11065	0	10886	128	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.



All (128) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:LYS:HG2	1:B:158:PRO:HD2	1.71	0.73
1:B:347:ASN:ND2	5:B:904:HOH:O	2.27	0.68
1:B:286:LEU:O	1:B:290:ASN:ND2	2.28	0.67
1:B:150:GLU:HG3	1:B:155:GLU:HB3	1.77	0.66
1:A:483:MET:SD	5:A:909:HOH:O	2.54	0.65
1:B:78:MET:HG2	1:B:576:GLU:HG3	1.80	0.64
1:A:66:LYS:HG3	1:A:69:ARG:HH21	1.63	0.63
1:B:6:SER:OG	1:B:11:ARG:NH1	2.31	0.62
1:B:145:THR:O	1:B:164:ARG:NH2	2.33	0.61
1:A:317:ASP:OD2	1:A:333:ARG:NH1	2.34	0.61
1:B:240:GLU:OE1	1:B:318:ARG:NH1	2.34	0.60
1:A:638:ARG:NH2	1:A:707:ASN:O	2.34	0.60
1:A:400:MET:O	1:A:438:THR:OG1	2.18	0.60
1:B:482:THR:HG21	1:B:585:LYS:HE3	1.83	0.59
1:B:230:VAL:HG13	1:B:241:ILE:HB	1.85	0.59
1:A:622:VAL:HB	1:A:632:GLU:HG3	1.85	0.58
1:A:658:VAL:HG21	1:A:699:ILE:HG21	1.85	0.58
1:B:533:PRO:HB2	1:B:558:VAL:HG21	1.86	0.58
1:B:398:PRO:HA	1:B:432:MET:HB2	1.86	0.58
1:B:53:LEU:HD13	1:B:405:GLU:HG2	1.86	0.58
1:B:129:ASN:OD1	1:B:312:ARG:NH2	2.37	0.57
1:A:269:ILE:HD13	1:A:337:PHE:HB2	1.86	0.57
1:B:210:GLY:N	5:B:907:HOH:O	2.36	0.56
1:A:348:ASP:O	1:A:350:ILE:N	2.37	0.56
1:B:379:SER:OG	1:B:380:ASP:N	2.38	0.56
1:A:50:ASN:OD1	5:A:901:HOH:O	2.18	0.56
1:A:339:ARG:HH12	3:A:802:LMR:H3	1.71	0.56
1:A:433:ASP:O	1:A:492:TYR:OH	2.15	0.55
1:B:247:SER:HB2	1:B:258:LYS:HD3	1.89	0.55
1:A:155:GLU:O	1:A:161:ASN:ND2	2.41	0.53
1:B:690:ILE:HD13	1:B:690:ILE:H	1.74	0.53
1:B:116:GLN:HG3	1:B:267:THR:HG23	1.90	0.52
1:A:142:LEU:HD12	1:A:148:ILE:HG12	1.90	0.52
1:B:291:TRP:O	1:B:295:ASN:ND2	2.40	0.51
1:B:123:ASN:HB3	1:B:126:PHE:HB2	1.92	0.51
1:B:439:THR:HG21	1:B:463:ARG:HE	1.75	0.51
1:B:622:VAL:HB	1:B:632:GLU:HG3	1.92	0.50
1:A:608:TYR:HD2	1:A:637:LEU:HD13	1.78	0.49
1:B:193:PHE:HD1	1:B:243:ILE:HD11	1.77	0.49
1:A:7:VAL:HG11	1:A:36:TRP:HB3	1.96	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:7:VAL:HG11	1:B:36:TRP:HB3	1.96	0.48
1:B:538:SER:HA	1:B:560:VAL:HG11	1.95	0.48
1:B:259:ASP:OD1	1:B:260:VAL:N	2.45	0.48
1:B:180:VAL:HG11	1:B:232:LEU:HD13	1.94	0.48
1:B:612:TRP:CD1	1:B:618:GLY:HA2	2.49	0.48
1:A:698:LEU:HD23	1:A:713:ILE:HD12	1.96	0.48
1:A:344:LEU:HD22	1:A:709:TYR:CD1	2.48	0.48
1:A:392:SER:HB3	1:A:428:LYS:HD3	1.96	0.48
1:A:463:ARG:HG3	1:A:488:TRP:HZ3	1.79	0.48
1:A:529:LYS:NZ	1:A:532:GLN:OE1	2.43	0.48
1:A:516:TRP:NE1	1:A:518:MET:O	2.47	0.47
1:A:96:ASP:N	1:A:96:ASP:OD2	2.47	0.47
1:B:431:ILE:HB	1:B:456:ILE:HG22	1.97	0.47
1:A:273:GLU:OE1	1:A:399:LYS:NZ	2.37	0.46
1:B:182:LEU:HA	1:B:211:LEU:H	1.80	0.46
1:A:437:ARG:O	1:A:441:ASN:ND2	2.47	0.46
1:A:431:ILE:HD11	1:A:449:ALA:HB3	1.96	0.46
1:A:609:VAL:HG11	1:A:695:ALA:HB2	1.96	0.46
1:A:344:LEU:HD22	1:A:709:TYR:HD1	1.80	0.46
1:A:11:ARG:HB2	1:A:351:VAL:HG23	1.98	0.46
1:B:104:GLY:O	1:B:392:SER:OG	2.27	0.46
1:B:119:VAL:HG21	1:B:130:ALA:HB2	1.97	0.46
1:B:128:LEU:HD23	1:B:299:LEU:HD23	1.96	0.46
1:A:108:GLU:OE2	1:A:134:ARG:NH2	2.49	0.45
1:A:108:GLU:HG2	1:A:108:GLU:H	1.52	0.45
1:B:417:GLU:OE2	1:B:426:THR:N	2.49	0.45
1:B:401:HIS:NE2	1:B:435:GLU:OE1	2.49	0.45
1:A:680:ARG:NE	5:A:910:HOH:O	2.48	0.45
1:A:512:GLY:HA2	1:A:539:THR:O	2.16	0.45
1:A:317:ASP:OD1	1:A:389:ARG:NH2	2.50	0.45
1:B:217:PHE:CE2	1:B:323:PRO:HB3	2.52	0.45
1:A:585:LYS:HB3	1:A:586:GLU:H	1.59	0.45
1:A:659:ARG:HA	1:A:659:ARG:HD2	1.82	0.44
1:B:219:GLY:HA2	1:B:323:PRO:HD3	1.97	0.44
1:A:393:ILE:O	1:A:428:LYS:N	2.45	0.44
1:A:154:ALA:HB1	1:A:161:ASN:HD21	1.82	0.44
1:A:314:LEU:HB3	1:A:333:ARG:HD3	1.98	0.44
1:A:177:ASP:OD1	1:A:186:SER:OG	2.30	0.44
1:B:292:LEU:HD11	1:B:296:LYS:HE3	1.99	0.44
1:B:362:ILE:HG23	1:B:412:LEU:HD11	1.98	0.44
1:B:318:ARG:NH1	1:B:320:TYR:OH	2.51	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:134:ARG:O	1:B:264:SER:N	2.42	0.44
1:B:338:VAL:HG13	1:B:395:ILE:HG12	1.99	0.44
1:A:12:ILE:HG23	1:A:350:ILE:HG22	2.00	0.44
1:A:154:ALA:HB1	1:A:161:ASN:ND2	2.33	0.44
1:B:697:GLU:O	1:B:701:SER:OG	2.30	0.44
1:A:364:ASP:O	1:A:368:THR:HB	2.18	0.43
1:B:119:VAL:HG11	1:B:127:ALA:HA	1.99	0.43
1:A:518:MET:N	1:A:518:MET:SD	2.92	0.43
1:A:39:VAL:HG22	1:A:366:LEU:HD13	2.00	0.43
1:B:193:PHE:HE2	1:B:202:VAL:HG22	1.84	0.43
1:A:12:ILE:HG12	1:A:350:ILE:HG22	2.00	0.43
1:A:53:LEU:HD13	1:A:405:GLU:HG2	2.00	0.43
1:A:407:ALA:O	1:A:411:GLU:HG2	2.19	0.43
1:B:349:ALA:HB3	1:B:363:MET:HE1	2.01	0.43
1:B:413:PHE:CE2	1:B:429:ILE:HG12	2.54	0.43
1:A:605:ILE:HD13	1:A:637:LEU:HD12	2.01	0.42
1:B:206:ASP:OD2	1:B:206:ASP:N	2.52	0.42
1:B:613:VAL:O	1:B:718:ARG:NH1	2.42	0.42
1:B:479:ARG:NE	5:B:912:HOH:O	2.51	0.42
1:A:397:LYS:HD3	1:A:398:PRO:HD2	2.01	0.42
1:A:358:VAL:HB	1:A:363:MET:HE2	2.01	0.42
1:B:200:LEU:HD23	1:B:211:LEU:HD11	2.00	0.42
1:A:173:ARG:NH2	5:A:911:HOH:O	2.48	0.42
1:A:259:ASP:OD1	1:A:260:VAL:N	2.52	0.42
1:B:463:ARG:NH1	1:B:467:GLU:HB2	2.35	0.42
1:B:265:ALA:O	1:B:334:SER:OG	2.30	0.41
1:B:240:GLU:OE1	1:B:320:TYR:OH	2.33	0.41
1:A:386:ILE:HG23	1:A:389:ARG:HH12	1.84	0.41
1:A:395:ILE:O	1:A:429:ILE:HA	2.20	0.41
1:A:541:TRP:HZ3	3:A:802:LMR:H3A	1.85	0.41
1:B:12:ILE:HD13	1:B:12:ILE:H	1.85	0.41
1:B:231:LEU:HD11	1:B:238:HIS:HB3	2.02	0.41
1:B:316:ARG:HA	1:B:316:ARG:HD3	1.83	0.41
1:A:116:GLN:OE1	1:A:541:TRP:NE1	2.51	0.41
1:A:612:TRP:CD1	1:A:618:GLY:HA2	2.55	0.41
1:A:649:ARG:NH1	1:A:649:ARG:O	2.54	0.41
1:B:158:PRO:HB2	1:B:159:THR:H	1.78	0.41
1:B:12:ILE:HG22	1:B:350:ILE:HG23	2.02	0.41
1:B:269:ILE:HG12	1:B:337:PHE:HB2	2.03	0.41
1:A:292:LEU:HD11	1:A:296:LYS:HE2	2.01	0.41
1:B:10:LEU:HD21	1:B:40:ASP:HA	2.03	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:273:GLU:O	1:A:634:ARG:NH2	2.54	0.41
1:B:14:ARG:HB3	1:B:14:ARG:HH11	1.85	0.41
1:B:31:ASP:HA	1:B:32:PRO:HD3	1.86	0.41
1:B:443:LYS:HE2	1:B:443:LYS:HB2	1.91	0.40
1:B:245:PRO:HG3	1:B:256:GLY:HA3	2.02	0.40
1:B:25:LEU:N	1:B:26:PRO:HD2	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	716/741 (97%)	668 (93%)	41 (6%)	7 (1%)	19	49
1	B	710/741 (96%)	654 (92%)	45 (6%)	11 (2%)	13	38
All	All	1426/1482 (96%)	1322 (93%)	86 (6%)	18 (1%)	15	42

All (18) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	348	ASP
1	A	349	ALA
1	B	158	PRO
1	B	585	LYS
1	A	4	ARG
1	A	151	THR
1	B	209	THR
1	B	211	LEU
1	B	265	ALA
1	B	352	ASP
1	B	387	ASN
1	A	355	GLY

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	680	ARG
1	B	355	GLY
1	B	378	ALA
1	B	72	VAL
1	A	325	GLY
1	A	514	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	579/594 (98%)	560 (97%)	19 (3%)	45	78
1	B	578/594 (97%)	552 (96%)	26 (4%)	34	67
All	All	1157/1188 (97%)	1112 (96%)	45 (4%)	39	73

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	96	ASP
1	A	108	GLU
1	A	213	ASN
1	A	225	GLU
1	A	289	ARG
1	A	310	PHE
1	A	319	ASN
1	A	338	VAL
1	A	386	ILE
1	A	396	VAL
1	A	527	GLU
1	A	572	ARG
1	A	575	ILE
1	A	586	GLU
1	A	595	ARG
1	A	627	ASP
1	A	648	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	688	ASP
1	A	724	ARG
1	B	10	LEU
1	B	12	ILE
1	B	14	ARG
1	B	29	ASP
1	B	33	ASP
1	B	54	LEU
1	B	72	VAL
1	B	97	ASP
1	B	117	LEU
1	B	119	VAL
1	B	155	GLU
1	B	161	ASN
1	B	162	LYS
1	B	197	ASP
1	B	230	VAL
1	B	299	LEU
1	B	316	ARG
1	B	317	ASP
1	B	439	THR
1	B	454	VAL
1	B	575	ILE
1	B	595	ARG
1	B	648	LEU
1	B	668	LEU
1	B	690	ILE
1	B	715	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	347	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	LMR	A	802	2	1,8,8	0.40	0	2,10,10	2.96	1 (50%)
4	GLV	B	803	2	1,4,4	5.67	1 (100%)	0,4,4	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LMR	A	802	2	-	0/2/8/8	0/0/0/0
4	GLV	B	803	2	-	0/0/2/2	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	803	GLV	O1-C1	5.67	1.41	1.21

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	LMR	C4-C3-C2	-4.13	109.95	114.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	LMR	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	720/741 (97%)	-0.06	12 (1%)	73 70	40, 70, 104, 138	0
1	B	716/741 (96%)	0.05	35 (4%)	33 27	51, 79, 119, 140	0
All	All	1436/1482 (96%)	-0.00	47 (3%)	50 43	40, 74, 114, 140	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	184	SER	5.5
1	B	157	GLY	5.1
1	B	614	ASP	4.0
1	B	247	SER	3.9
1	B	195	VAL	3.8
1	A	64	ILE	3.6
1	B	258	LYS	3.5
1	B	46	LEU	3.5
1	B	249	VAL	3.5
1	A	11	ARG	3.4
1	B	163	VAL	3.4
1	A	185	GLY	3.2
1	B	149	PRO	3.1
1	B	251	THR	3.0
1	A	643	LEU	3.0
1	B	165	GLY	2.9
1	B	167	LYS	2.9
1	B	200	LEU	2.9
1	B	250	GLY	2.9
1	A	10	LEU	2.8
1	A	152	ASP	2.8
1	B	253	ASP	2.8
1	A	73	ILE	2.7
1	B	669	VAL	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	606	LEU	2.6
1	B	348	ASP	2.6
1	A	12	ILE	2.5
1	A	625	ILE	2.5
1	B	536	GLY	2.5
1	B	698	LEU	2.5
1	B	172	ALA	2.4
1	B	151	THR	2.4
1	B	193	PHE	2.4
1	B	679	TYR	2.3
1	B	171	TYR	2.3
1	B	683	ALA	2.3
1	B	246	GLU	2.2
1	B	217	PHE	2.2
1	B	686	PHE	2.1
1	B	571	ARG	2.1
1	B	192	GLY	2.1
1	A	355	GLY	2.0
1	B	219	GLY	2.0
1	B	158	PRO	2.0
1	B	328	PHE	2.0
1	A	60	LEU	2.0
1	B	208	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	LMR	A	802	9/9	0.89	0.21	-0.87	67,78,93,98	0
2	MG	B	801	1/1	0.98	0.12	-1.35	67,67,67,67	0
4	GLV	B	803	5/5	0.96	0.11	-1.66	56,62,72,72	0
2	MG	A	801	1/1	0.96	0.17	-2.88	76,76,76,76	0
2	MG	B	802	1/1	0.93	0.22	-	55,55,55,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.