



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:49 PM GMT

PDB ID : 1HBX
Title : TERNARY COMPLEX OF SAP-1 AND SRF WITH SPECIFIC SRE DNA
Authors : Hassler, M.; Richmond, T.J.
Deposited on : 2001-04-20
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1249 (3.20-3.12)
Ramachandran outliers	100387	1222 (3.20-3.12)
Sidechain outliers	100360	1221 (3.20-3.12)




The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	92	
1	B	92	
1	D	92	
1	E	92	
2	C	26	
2	F	26	
3	G	157	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	157	 17% 46% 6% 31%
4	W	26	 85% 15%
4	X	26	 100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6682 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERUM RESPONSE FACTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	87	Total	C	N	O	S	0	0	0
			681	430	121	126	4			
1	B	86	Total	C	N	O	S	0	0	0
			685	434	122	125	4			
1	D	86	Total	C	N	O	S	0	0	0
			677	428	120	125	4			
1	E	89	Total	C	N	O	S	0	0	0
			691	436	123	128	4			

- Molecule 2 is a DNA chain called 5'-D(*GP*AP*TP*GP*GP*CP*CP*TP*AP*AP*TP*T
P*AP* GP*GP*AP*CP*TP*TP*CP*CP*GP*GP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	26	Total	C	N	O	P	0	0	0
			533	255	96	157	25			
2	F	26	Total	C	N	O	P	0	0	0
			533	255	96	157	25			

- Molecule 3 is a protein called ETS-DOMAIN PROTEIN ELK-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	112	Total	C	N	O	S	0	0	0
			925	598	163	160	4			
3	H	109	Total	C	N	O	S	0	0	0
			903	587	156	156	4			

- Molecule 4 is a DNA chain called 5'-D(*CP*AP*CP*AP*CP*CP*GP*GP*AP*AP*GP*T
P*CP* CP*TP*AP*AP*TP*TP*AP*GP*GP*CP*CP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	W	26	Total	C	N	O	P	0	0	0
			527	252	99	151	25			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	X	26	Total	C	N	O	P	0	0	0
			527	252	99	151	25			

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: SERUM RESPONSE FACTOR




• Molecule 1: SERUM RESPONSE FACTOR




• Molecule 1: SERUM RESPONSE FACTOR




• Molecule 1: SERUM RESPONSE FACTOR






• Molecule 2: 5'-D(*GP*AP*TP*GP*GP*CP*CP*TP*AP*AP*TP*TP*AP* GP*GP*AP*CP*TP*TP*CP*CP*GP*GP*TP*G)-3'

Chain C: 92% 8%





• Molecule 2: 5'-D(*GP*AP*TP*GP*GP*CP*CP*TP*AP*AP*TP*TP*AP* GP*GP*AP*CP*TP*TP*CP*CP*GP*GP*TP*G)-3'

Chain F: 96%





• Molecule 3: ETS-DOMAIN PROTEIN ELK-4

Chain G: 17% 43% 11% 29%



• Molecule 3: ETS-DOMAIN PROTEIN ELK-4

Chain H: 17% 46% 6% 31%



• Molecule 4: 5'-D(*CP*AP*CP*AP*CP*CP*GP*GP*AP*AP*GP*TP*CP* CP*TP*AP*AP*TP*TP*AP*GP*GP*CP*CP*AP*T)-3'

Chain W: 85% 15%

C-17	A-16	C-15	A-14	C-13	C-12	G-11	G-10	A-9	A-8	G-7	T-6	C-5	C-4	T-3	A-2	A-1	T1	T2	A3	G4	G5	C6	C7	A8	T9
------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	----	----	----	----	----	----	----	----

● Molecule 4: 5'-D(*CP*AP*CP*AP*CP*CP*GP*GP*AP*AP*GP*TP*CP* CP*TP*AP*AP*TP*TP*AP*GP*GP*CP*CP*AP*T)-3'

Chain X:

100%

C-17	A-16	C-15	A-14	C-13	C-12	G-11	G-10	A-9	A-8	G-7	T-6	C-5	C-4	T-3	A-2	A-1	T1	T2	A3	G4	G5	C6	C7	A8	T9
------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	----	----	----	----	----	----	----	----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	142.68Å 144.39Å 75.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.00 – 3.15	Depositor
% Data completeness (in resolution range)	98.8 (51.00-3.15)	Depositor
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.252 , 0.285	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6682	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/689	0.71	1/923 (0.1%)
1	B	0.48	0/693	0.66	0/926
1	D	0.44	0/685	0.67	0/918
1	E	0.43	0/700	0.65	0/938
2	C	0.71	0/597	0.88	1/921 (0.1%)
2	F	0.60	0/597	0.82	0/921
3	G	0.35	0/947	0.59	0/1274
3	H	0.32	0/925	0.52	0/1247
4	W	0.76	0/591	1.00	4/909 (0.4%)
4	X	0.59	0/591	0.85	0/909
All	All	0.51	0/7015	0.74	6/9886 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	W	5	DG	C5'-C4'-C3'	-6.96	101.57	114.10
1	A	178	LEU	CA-CB-CG	5.88	128.83	115.30
4	W	4	DG	C5'-C4'-C3'	-5.85	103.57	114.10
4	W	-8	DA	N9-C1'-C2'	5.36	122.78	112.60
2	C	-8	DT	C5'-C4'-C3'	-5.32	104.53	114.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	-6	DA	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	681	0	717	47	0
1	B	685	0	736	51	0
1	D	677	0	714	61	0
1	E	691	0	724	73	0
2	C	533	0	296	56	0
2	F	533	0	296	78	0
3	G	925	0	909	136	0
3	H	903	0	896	115	0
4	W	527	0	293	80	0
4	X	527	0	293	80	0
All	All	6682	0	5874	683	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 54.

The worst 5 of 683 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:X:-14:DA:C2'	4:X:-13:DC:H5''	1.66	1.25
2:F:4:DC:C2'	2:F:3:DT:H5''	1.69	1.23
2:F:-10:DC:C2'	2:F:-11:DC:H5''	1.71	1.20
2:F:4:DC:H2''	2:F:3:DT:C5'	1.76	1.15
2:C:-10:DC:C2'	2:C:-11:DC:H5''	1.76	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/92 (92%)	64 (75%)	17 (20%)	4 (5%)	3	21
1	B	84/92 (91%)	73 (87%)	8 (10%)	3 (4%)	4	28
1	D	84/92 (91%)	73 (87%)	9 (11%)	2 (2%)	7	41
1	E	87/92 (95%)	76 (87%)	6 (7%)	5 (6%)	2	16
3	G	108/157 (69%)	67 (62%)	30 (28%)	11 (10%)	1	3
3	H	105/157 (67%)	67 (64%)	27 (26%)	11 (10%)	1	3
All	All	553/682 (81%)	420 (76%)	97 (18%)	36 (6%)	1	12

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	3	SER
3	G	50	ASN
3	G	78	GLN
3	H	50	ASN
3	H	80	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	75/80 (94%)	71 (95%)	4 (5%)	28	67
1	B	77/80 (96%)	72 (94%)	5 (6%)	21	60
1	D	75/80 (94%)	72 (96%)	3 (4%)	38	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	76/80 (95%)	70 (92%)	6 (8%)	15	51
3	G	97/142 (68%)	87 (90%)	10 (10%)	9	34
3	H	97/142 (68%)	92 (95%)	5 (5%)	29	68
All	All	497/604 (82%)	464 (93%)	33 (7%)	21	59

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	175	LEU
3	G	15	LEU
3	H	45	TRP
1	E	182	GLN
1	E	214	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	16	GLN
3	G	20	ASN
3	G	92	ASN
3	G	9	GLN
3	H	13	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.