



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 11, 2017 – 03:03 PM EST

PDB ID : 5HC5
Title : The structure of esterase Est22 mutant-S188A
Authors : Li, J.; Huang, J.
Deposited on : 2016-01-04
Resolution : 2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442

i

X-RAY DIFFRACTION

A.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	1003 (2.46-2.42)
Clashscore	102246	1071 (2.46-2.42)
Ramachandran outliers	100387	1065 (2.46-2.42)
Sidechain outliers	100360	1065 (2.46-2.42)
RSRZ outliers	91569	1005 (2.46-2.42)

Mol	Chain	Length	Quality of chain
1	A	365	<div><div></div><div>88%</div><div>5%</div><div>7%</div></div>
1	B	365	<div><div></div><div>86%</div><div>7%</div><div>7%</div></div>
1	C	365	<div><div></div><div>88%</div><div>•</div><div>7%</div></div>
1	D	365	<div><div></div><div>87%</div><div>5%</div><div>7%</div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lipolytic enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	S	0	7	0
			2576	1614	434	511	17			
1	B	341	Total	C	N	O	S	0	9	0
			2600	1629	438	516	17			
1	C	338	Total	C	N	O	S	0	9	0
			2580	1617	435	511	17			
1	D	338	Total	C	N	O	S	0	9	0
			2580	1617	435	511	17			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP H6BDX1
A	-19	GLY	-	expression tag	UNP H6BDX1
A	-18	SER	-	expression tag	UNP H6BDX1
A	-17	SER	-	expression tag	UNP H6BDX1
A	-16	HIS	-	expression tag	UNP H6BDX1
A	-15	HIS	-	expression tag	UNP H6BDX1
A	-14	HIS	-	expression tag	UNP H6BDX1
A	-13	HIS	-	expression tag	UNP H6BDX1
A	-12	HIS	-	expression tag	UNP H6BDX1
A	-11	HIS	-	expression tag	UNP H6BDX1
A	-10	HIS	-	expression tag	UNP H6BDX1
A	-9	SER	-	expression tag	UNP H6BDX1
A	-8	SER	-	expression tag	UNP H6BDX1
A	-7	GLY	-	expression tag	UNP H6BDX1
A	-6	LEU	-	expression tag	UNP H6BDX1
A	-5	VAL	-	expression tag	UNP H6BDX1
A	-4	PRO	-	expression tag	UNP H6BDX1
A	-3	ARG	-	expression tag	UNP H6BDX1
A	-2	GLY	-	expression tag	UNP H6BDX1
A	-1	SER	-	expression tag	UNP H6BDX1
A	0	HIS	-	expression tag	UNP H6BDX1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	188	ALA	SER	engineered mutation	UNP H6BDX1
B	-20	MET	-	expression tag	UNP H6BDX1
B	-19	GLY	-	expression tag	UNP H6BDX1
B	-18	SER	-	expression tag	UNP H6BDX1
B	-17	SER	-	expression tag	UNP H6BDX1
B	-16	HIS	-	expression tag	UNP H6BDX1
B	-15	HIS	-	expression tag	UNP H6BDX1
B	-14	HIS	-	expression tag	UNP H6BDX1
B	-13	HIS	-	expression tag	UNP H6BDX1
B	-12	HIS	-	expression tag	UNP H6BDX1
B	-11	HIS	-	expression tag	UNP H6BDX1
B	-10	HIS	-	expression tag	UNP H6BDX1
B	-9	SER	-	expression tag	UNP H6BDX1
B	-8	SER	-	expression tag	UNP H6BDX1
B	-7	GLY	-	expression tag	UNP H6BDX1
B	-6	LEU	-	expression tag	UNP H6BDX1
B	-5	VAL	-	expression tag	UNP H6BDX1
B	-4	PRO	-	expression tag	UNP H6BDX1
B	-3	ARG	-	expression tag	UNP H6BDX1
B	-2	GLY	-	expression tag	UNP H6BDX1
B	-1	SER	-	expression tag	UNP H6BDX1
B	0	HIS	-	expression tag	UNP H6BDX1
B	188	ALA	SER	engineered mutation	UNP H6BDX1
C	-20	MET	-	expression tag	UNP H6BDX1
C	-19	GLY	-	expression tag	UNP H6BDX1
C	-18	SER	-	expression tag	UNP H6BDX1
C	-17	SER	-	expression tag	UNP H6BDX1
C	-16	HIS	-	expression tag	UNP H6BDX1
C	-15	HIS	-	expression tag	UNP H6BDX1
C	-14	HIS	-	expression tag	UNP H6BDX1
C	-13	HIS	-	expression tag	UNP H6BDX1
C	-12	HIS	-	expression tag	UNP H6BDX1
C	-11	HIS	-	expression tag	UNP H6BDX1
C	-10	HIS	-	expression tag	UNP H6BDX1
C	-9	SER	-	expression tag	UNP H6BDX1
C	-8	SER	-	expression tag	UNP H6BDX1
C	-7	GLY	-	expression tag	UNP H6BDX1
C	-6	LEU	-	expression tag	UNP H6BDX1
C	-5	VAL	-	expression tag	UNP H6BDX1
C	-4	PRO	-	expression tag	UNP H6BDX1
C	-3	ARG	-	expression tag	UNP H6BDX1
C	-2	GLY	-	expression tag	UNP H6BDX1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	SER	-	expression tag	UNP H6BDX1
C	0	HIS	-	expression tag	UNP H6BDX1
C	188	ALA	SER	engineered mutation	UNP H6BDX1
D	-20	MET	-	expression tag	UNP H6BDX1
D	-19	GLY	-	expression tag	UNP H6BDX1
D	-18	SER	-	expression tag	UNP H6BDX1
D	-17	SER	-	expression tag	UNP H6BDX1
D	-16	HIS	-	expression tag	UNP H6BDX1
D	-15	HIS	-	expression tag	UNP H6BDX1
D	-14	HIS	-	expression tag	UNP H6BDX1
D	-13	HIS	-	expression tag	UNP H6BDX1
D	-12	HIS	-	expression tag	UNP H6BDX1
D	-11	HIS	-	expression tag	UNP H6BDX1
D	-10	HIS	-	expression tag	UNP H6BDX1
D	-9	SER	-	expression tag	UNP H6BDX1
D	-8	SER	-	expression tag	UNP H6BDX1
D	-7	GLY	-	expression tag	UNP H6BDX1
D	-6	LEU	-	expression tag	UNP H6BDX1
D	-5	VAL	-	expression tag	UNP H6BDX1
D	-4	PRO	-	expression tag	UNP H6BDX1
D	-3	ARG	-	expression tag	UNP H6BDX1
D	-2	GLY	-	expression tag	UNP H6BDX1
D	-1	SER	-	expression tag	UNP H6BDX1
D	0	HIS	-	expression tag	UNP H6BDX1
D	188	ALA	SER	engineered mutation	UNP H6BDX1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	52	Total O 52 52	0	0
2	B	50	Total O 50 50	0	0
2	C	47	Total O 47 47	0	0
2	D	49	Total O 49 49	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lipolytic enzyme




- Molecule 1: Lipolytic enzyme



- Molecule 1: Lipolytic enzyme



- Molecule 1: Lipolytic enzyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.47Å 121.80Å 150.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.43 47.28 – 2.43	Depositor EDS
% Data completeness (in resolution range)	87.7 (50.00-2.43) 87.7 (47.28-2.43)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.91 (at 2.42Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.161 , 0.226 0.168 , 0.226	Depositor DCC
R_{free} test set	2476 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10534	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/2640	0.79	2/3582 (0.1%)
1	B	0.62	0/2665	0.77	1/3616 (0.0%)
1	C	0.61	0/2644	0.77	0/3586
1	D	0.60	0/2644	0.76	0/3586
All	All	0.61	0/10593	0.77	3/14370 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	290	ARG	NE-CZ-NH1	5.83	123.22	120.30
1	A	331	ARG	NE-CZ-NH1	5.71	123.16	120.30
1	B	290	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2576	0	2492	5	0
1	B	2600	0	2523	11	0
1	C	2580	0	2504	10	0
1	D	2580	0	2504	6	0
2	A	52	0	0	0	0
2	B	50	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	47	0	0	2	0
2	D	49	0	0	1	0
All	All	10534	0	10023	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:72:ILE:HD12	2:B:450:HOH:O	2.00	0.61
1:C:18:ILE:HD12	1:C:325:ALA:HA	1.86	0.58
1:B:123:TRP:CD1	1:B:321:ILE:HG22	2.42	0.54
1:B:207:GLN:OE1	1:B:276[B]:LEU:HD23	2.07	0.54
1:D:281:ILE:O	1:D:309:CYS:HA	2.09	0.52
1:B:17[A]:ALA:HB1	1:C:324:ILE:HD13	1.93	0.51
1:D:259:ARG:HG2	1:D:267:PHE:CE1	2.47	0.49
1:C:64:ILE:CG2	1:C:323:PRO:HB2	2.43	0.49
1:B:17[B]:ALA:HB1	1:C:324:ILE:HD13	1.95	0.48
1:A:162:VAL:HG21	1:A:200:ARG:HG2	1.95	0.47
1:C:4:LYS:N	2:C:401:HOH:O	2.47	0.46
1:A:217:CYS:N	1:A:282:SER:O	2.49	0.46
1:B:281:ILE:O	1:B:309:CYS:HA	2.15	0.45
1:B:59:CYS:HB3	1:B:323:PRO:HD2	1.97	0.45
1:D:207:GLN:OE1	1:D:276[A]:LEU:HD12	2.18	0.44
1:A:185:ALA:HA	1:A:214:TYR:O	2.17	0.44
1:B:30:VAL:HG23	1:B:249:MET:HG3	1.98	0.43
1:B:235:ASN:HB2	1:B:288:PRO:HA	1.99	0.43
1:B:98[A]:LYS:HB3	1:B:98[A]:LYS:HE2	1.85	0.43
1:D:30:VAL:HG21	1:D:36:MET:SD	2.58	0.43
1:C:18:ILE:HD13	1:C:324:ILE:HD12	2.01	0.42
1:A:281:ILE:O	1:A:309:CYS:HA	2.20	0.42
1:D:218:PRO:HD2	2:D:413:HOH:O	2.20	0.42
1:B:18:ILE:CD1	1:B:325:ALA:HA	2.49	0.42
1:C:281:ILE:O	1:C:309:CYS:HA	2.21	0.41
1:C:114:SER:HB3	1:C:117:TYR:CE1	2.55	0.41
1:A:285:GLU:HB2	1:A:313:MET:HA	2.02	0.41
1:C:72:ILE:HG13	2:C:446:HOH:O	2.20	0.41
1:C:10:ARG:O	1:C:314:GLY:HA2	2.22	0.40
1:D:216:LEU:O	1:D:217:CYS:C	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/365 (94%)	323 (94%)	20 (6%)	0	100	100
1	B	348/365 (95%)	331 (95%)	16 (5%)	1 (0%)	46	56
1	C	343/365 (94%)	329 (96%)	12 (4%)	2 (1%)	30	36
1	D	343/365 (94%)	324 (94%)	18 (5%)	1 (0%)	46	56
All	All	1377/1460 (94%)	1307 (95%)	66 (5%)	4 (0%)	46	56

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	239	LEU
1	B	113	LEU
1	C	113	LEU
1	D	343	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/287 (94%)	266 (98%)	5 (2%)	66	79
1	B	274/287 (96%)	266 (97%)	8 (3%)	50	66
1	C	272/287 (95%)	269 (99%)	3 (1%)	80	87

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	272/287 (95%)	264 (97%)	8 (3%)	50 66
All	All	1089/1148 (95%)	1065 (98%)	24 (2%)	58 74

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	MET
1	A	44	GLU
1	A	96	THR
1	A	227	GLU
1	A	280	PHE
1	B	22	MET
1	B	53	ARG
1	B	74	ASP
1	B	112	SER
1	B	208	ASP
1	B	262	LEU
1	B	274	SER
1	B	280	PHE
1	C	48	VAL
1	C	280	PHE
1	C	324	ILE
1	D	44	GLU
1	D	74	ASP
1	D	203	GLN
1	D	214	TYR
1	D	226	SER
1	D	227	GLU
1	D	271	GLU
1	D	280	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/365 (93%)	-0.37	0 100 100	26, 37, 60, 98	0
1	B	341/365 (93%)	-0.51	4 (1%) 81 82	25, 34, 59, 100	0
1	C	338/365 (92%)	-0.32	8 (2%) 62 61	25, 37, 61, 88	0
1	D	338/365 (92%)	-0.52	4 (1%) 81 82	26, 38, 60, 83	0
All	All	1357/1460 (92%)	-0.43	16 (1%) 81 82	25, 37, 61, 100	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	43	GLU	4.1
1	B	43	GLU	3.4
1	C	27	GLY	3.3
1	C	26	GLY	3.1
1	C	174	GLU	3.1
1	D	98[A]	LYS	2.8
1	C	98[A]	LYS	2.7
1	C	170[A]	SER	2.6
1	D	96	THR	2.5
1	C	44	GLU	2.5
1	B	44	GLU	2.3
1	B	96	THR	2.2
1	C	41	SER	2.2
1	D	75	TYR	2.1
1	D	170[A]	SER	2.1
1	B	26	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.