



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:08 AM GMT

PDB ID : 3HD0  
Title : Crystal structure of Tm1865, an Endonuclease V from Thermotoga Maritima  
Authors : Utepbergenov, D.; Cooper, D.R.; Derewenda, U.; Derewenda, Z.S.; Integrated Center for Structure and Function Innovation (ISFI)  
Deposited on : 2009-05-06  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	 69% 21% • 7%
1	B	237	 70% 19% • 9%
1	D	237	 70% 21% • 6%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5512 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endonuclease V.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	Se	0	0	0
			1767	1150	299	312	2	4			
1	B	216	Total	C	N	O	S	Se	0	0	0
			1728	1127	292	304	2	3			
1	D	223	Total	C	N	O	S	Se	0	0	0
			1796	1168	307	315	2	4			

There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MSE	-	expression tag	UNP Q9X2H9
A	-10	GLY	-	expression tag	UNP Q9X2H9
A	-9	SER	-	expression tag	UNP Q9X2H9
A	-8	ASP	-	expression tag	UNP Q9X2H9
A	-7	LYS	-	expression tag	UNP Q9X2H9
A	-6	ILE	-	expression tag	UNP Q9X2H9
A	-5	HIS	-	expression tag	UNP Q9X2H9
A	-4	HIS	-	expression tag	UNP Q9X2H9
A	-3	HIS	-	expression tag	UNP Q9X2H9
A	-2	HIS	-	expression tag	UNP Q9X2H9
A	-1	HIS	-	expression tag	UNP Q9X2H9
A	0	HIS	-	expression tag	UNP Q9X2H9
A	223	GLY	-	expression tag	UNP Q9X2H9
A	224	LEU	-	expression tag	UNP Q9X2H9
A	225	PHE	-	expression tag	UNP Q9X2H9
B	-11	MSE	-	expression tag	UNP Q9X2H9
B	-10	GLY	-	expression tag	UNP Q9X2H9
B	-9	SER	-	expression tag	UNP Q9X2H9
B	-8	ASP	-	expression tag	UNP Q9X2H9
B	-7	LYS	-	expression tag	UNP Q9X2H9
B	-6	ILE	-	expression tag	UNP Q9X2H9
B	-5	HIS	-	expression tag	UNP Q9X2H9
B	-4	HIS	-	expression tag	UNP Q9X2H9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	HIS	-	expression tag	UNP Q9X2H9
B	-2	HIS	-	expression tag	UNP Q9X2H9
B	-1	HIS	-	expression tag	UNP Q9X2H9
B	0	HIS	-	expression tag	UNP Q9X2H9
B	223	GLY	-	expression tag	UNP Q9X2H9
B	224	LEU	-	expression tag	UNP Q9X2H9
B	225	PHE	-	expression tag	UNP Q9X2H9
D	-11	MSE	-	expression tag	UNP Q9X2H9
D	-10	GLY	-	expression tag	UNP Q9X2H9
D	-9	SER	-	expression tag	UNP Q9X2H9
D	-8	ASP	-	expression tag	UNP Q9X2H9
D	-7	LYS	-	expression tag	UNP Q9X2H9
D	-6	ILE	-	expression tag	UNP Q9X2H9
D	-5	HIS	-	expression tag	UNP Q9X2H9
D	-4	HIS	-	expression tag	UNP Q9X2H9
D	-3	HIS	-	expression tag	UNP Q9X2H9
D	-2	HIS	-	expression tag	UNP Q9X2H9
D	-1	HIS	-	expression tag	UNP Q9X2H9
D	0	HIS	-	expression tag	UNP Q9X2H9
D	223	GLY	-	expression tag	UNP Q9X2H9
D	224	LEU	-	expression tag	UNP Q9X2H9
D	225	PHE	-	expression tag	UNP Q9X2H9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	85	Total O 85 85	0	0
2	B	58	Total O 58 58	0	0
2	D	78	Total O 78 78	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Endonuclease V



#### • Molecule 1: Endonuclease V



#### • Molecule 1: Endonuclease V



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.57Å 71.97Å 120.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 46.18 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.70) 98.5 (46.18-2.70)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.30 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.163 , 0.255 0.176 , 0.251	Depositor DCC
$R_{free}$ test set	879 reflections (5.47%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.5	Xtriage
Anisotropy	0.318	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.3	EDS
Estimated twinning fraction	0.022 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 17001 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5512	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.76	0/1811	0.79	0/2446
1	B	0.69	0/1771	0.79	1/2392 (0.0%)
1	D	0.70	0/1841	0.77	1/2484 (0.0%)
All	All	0.72	0/5423	0.78	2/7322 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	D	120	LEU	CA-CB-CG	5.82	128.68	115.30
1	B	94	LEU	CA-CB-CG	5.81	128.66	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1767	0	1811	37	0
1	B	1728	0	1778	25	0
1	D	1796	0	1838	32	0
2	A	85	0	0	5	0
2	B	58	0	0	1	0
2	D	78	0	0	3	0
All	All	5512	0	5427	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:4:ARG:HH12	1:D:131:GLU:HB2	1.12	1.06
1:D:33:GLU:HA	1:D:33:GLU:OE1	1.80	0.81
1:D:4:ARG:NH1	1:D:131:GLU:HB2	1.94	0.81
1:A:76:ILE:HD11	1:A:92:LEU:HD12	1.70	0.74
1:A:172:THR:HG21	1:A:205:ARG:O	1.92	0.69

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	218/237 (92%)	212 (97%)	6 (3%)	0	100	100
1	B	214/237 (90%)	203 (95%)	11 (5%)	0	100	100
1	D	221/237 (93%)	215 (97%)	6 (3%)	0	100	100
All	All	653/711 (92%)	630 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/201 (95%)	175 (92%)	16 (8%)	14	30
1	B	187/201 (93%)	172 (92%)	15 (8%)	15	33
1	D	194/201 (96%)	173 (89%)	21 (11%)	8	18
All	All	572/603 (95%)	520 (91%)	52 (9%)	12	26

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	101	ARG
1	B	190	GLU
1	D	157	SER
1	B	118	ARG
1	B	128	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	218	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/237 (91%)	-0.72	0 100 100	14, 23, 41, 56	0
1	B	213/237 (89%)	-0.57	0 100 100	17, 28, 51, 69	0
1	D	219/237 (92%)	-0.70	0 100 100	16, 25, 44, 73	0
All	All	648/711 (91%)	-0.66	0 100 100	14, 25, 45, 73	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.