



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 11, 2017 – 05:27 AM EST

PDB ID : 5HD8
Title : Crystal structure of disulfide cross-linked D417C ClC-ec1
Authors : Mathews, I.I.; Khantwal, C.M.; Maduke, M.
Deposited on : 2016-01-05
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1112 (3.20-3.12)
Clashscore	102246	1249 (3.20-3.12)
Ramachandran outliers	100387	1222 (3.20-3.12)
Sidechain outliers	100360	1221 (3.20-3.12)
RSRZ outliers	91569	1117 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>15%</div> <div>• •</div> </div> </div>
1	B	450	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>14%</div> <div>• •</div> </div> </div>
2	C	222	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>17%</div> <div>•</div> </div> </div>
2	E	222	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>12%</div> <div>• •</div> </div> </div>
3	D	211	<div> <div>6%</div> <div> <div></div> <div>87%</div> <div>11%</div> <div>•</div> </div> </div>
3	F	211	<div> <div>4%</div> <div> <div></div> <div>81%</div> <div>18%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	A	501	-	-	-	X
4	CL	A	502	-	-	X	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H(+)/Cl(-) exchange transporter ClcA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	433	Total	C	N	O	S	0	0	0
			3248	2137	547	543	21			
1	B	432	Total	C	N	O	S	0	0	0
			3236	2131	542	542	21			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	16	MET	-	initiating methionine	UNP P37019
A	417	CYS	ASP	engineered mutation	UNP P37019
B	16	MET	-	initiating methionine	UNP P37019
B	417	CYS	ASP	engineered mutation	UNP P37019

- Molecule 2 is a protein called FAB FRAGMENT (HEAVY CHAIN).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	221	Total	C	N	O	S	0	0	0
			1672	1077	274	315	6			
2	E	220	Total	C	N	O	S	0	0	0
			1666	1074	273	313	6			

- Molecule 3 is a protein called FAB FRAGMENT (LIGHT CHAIN).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	211	Total	C	N	O	S	0	0	0
			1621	1008	271	334	8			
3	F	211	Total	C	N	O	S	0	0	0
			1621	1008	271	334	8			

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total 2	Cl 2	0	0
4	A	2	Total 2	Cl 2	0	0



• Molecule 3: FAB FRAGMENT (LIGHT CHAIN)



• Molecule 3: FAB FRAGMENT (LIGHT CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	231.74Å 96.06Å 169.95Å 90.00° 131.58° 90.00°	Depositor
Resolution (Å)	39.20 – 3.15 39.21 – 3.15	Depositor EDS
% Data completeness (in resolution range)	90.7 (39.20-3.15) 90.7 (39.21-3.15)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.205 , 0.257 0.202 , 0.256	Depositor DCC
R_{free} test set	2202 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	95.9	Xtriage
Anisotropy	0.368	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.015 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	13068	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3319	0.56	0/4503
1	B	0.37	0/3306	0.53	0/4485
2	C	0.38	0/1721	0.61	0/2355
2	E	0.41	0/1715	0.61	0/2348
3	D	0.35	0/1660	0.54	0/2257
3	F	0.42	0/1660	0.62	0/2257
All	All	0.38	0/13381	0.57	0/18205

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3248	0	3398	52	0
1	B	3236	0	3391	43	0
2	C	1672	0	1654	21	0
2	E	1666	0	1649	15	0
3	D	1621	0	1546	16	0
3	F	1621	0	1546	16	0
4	A	2	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	2	0	0	1	0
All	All	13068	0	13184	143	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 143 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:95:GLN:N	3:F:95:GLN:OE1	2.09	0.86
3:D:95:GLN:OE1	3:D:95:GLN:N	2.20	0.73
1:A:413:LEU:HD22	1:A:422:ILE:HD11	1.72	0.71
2:C:17:SER:HB3	2:C:84:SER:HA	1.73	0.69
1:B:191:ASN:OD1	1:B:230:ARG:NH1	2.29	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	429/450 (95%)	405 (94%)	18 (4%)	6 (1%)	14	55
1	B	428/450 (95%)	402 (94%)	24 (6%)	2 (0%)	34	76
2	C	219/222 (99%)	201 (92%)	15 (7%)	3 (1%)	14	55
2	E	218/222 (98%)	194 (89%)	21 (10%)	3 (1%)	14	55
3	D	209/211 (99%)	189 (90%)	18 (9%)	2 (1%)	19	63
3	F	209/211 (99%)	188 (90%)	17 (8%)	4 (2%)	10	48
All	All	1712/1766 (97%)	1579 (92%)	113 (7%)	20 (1%)	16	58

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	167	ARG
1	A	420	GLN
1	B	417	CYS
2	C	65	LYS
2	E	140	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/340 (96%)	310 (95%)	16 (5%)	31	71
1	B	325/340 (96%)	307 (94%)	18 (6%)	27	66
2	C	181/182 (100%)	170 (94%)	11 (6%)	23	62
2	E	181/182 (100%)	167 (92%)	14 (8%)	16	52
3	D	185/185 (100%)	179 (97%)	6 (3%)	46	81
3	F	185/185 (100%)	171 (92%)	14 (8%)	16	53
All	All	1383/1414 (98%)	1304 (94%)	79 (6%)	25	65

5 of 79 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	18	LEU
2	C	216	LYS
3	F	96	THR
2	C	22	CYS
2	C	188	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	ASN
1	A	420	GLN
3	F	75	ASN
3	F	144	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	433/450 (96%)	-0.16	7 (1%) 74 61	77, 101, 140, 186	0
1	B	432/450 (96%)	-0.08	19 (4%) 38 22	81, 112, 155, 216	0
2	C	221/222 (99%)	-0.26	8 (3%) 46 29	64, 100, 145, 171	0
2	E	220/222 (99%)	-0.47	2 (0%) 85 77	67, 95, 139, 184	0
3	D	211/211 (100%)	-0.11	13 (6%) 24 12	79, 116, 148, 159	0
3	F	211/211 (100%)	-0.18	9 (4%) 39 23	59, 89, 141, 162	0
All	All	1728/1766 (97%)	-0.19	58 (3%) 49 32	59, 104, 147, 216	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	7	SER	4.6
3	F	153	GLU	4.5
1	B	70	HIS	4.5
3	F	155	GLN	4.2
1	B	73	ASP	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	CL	A	502	1/1	0.87	0.67	3.26	140,140,140,140	0
4	CL	A	501	1/1	0.90	0.46	1.70	115,115,115,115	0
4	CL	B	502	1/1	0.76	0.38	0.81	113,113,113,113	0
4	CL	B	501	1/1	0.94	0.21	-2.06	104,104,104,104	0

6.5 Other polymers [i](#)

There are no such residues in this entry.