



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 14, 2016 – 07:38 PM EDT

PDB ID : 5HFA  
Title : Crystal structure of human acetylcholinesterase in complex with paraoxon and 2-PAM  
Authors : Franklin, M.F.; Rudolph, M.J.; Ginter, C.; Cassidy, M.S.; Cheung, J.  
Deposited on : 2016-01-06  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	<div> <div>2%</div> <div>90%</div> <div>8%</div> <div>..</div> </div>
1	B	542	<div> <div>2%</div> <div>92%</div> <div>6%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	DEP	A	604[A]	-	-	X	-
4	DEP	B	605[A]	-	-	-	X
5	EDO	A	606	-	-	-	X
5	EDO	B	606	-	-	-	X
5	EDO	B	607	-	-	-	X
6	FP1	A	608	-	-	-	X
7	NO3	B	608	-	-	-	X

## 2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 9173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	533	Total	C	N	O	S	0	6	0
			4207	2700	737	757	13			
1	B	538	Total	C	N	O	S	0	3	0
			4214	2704	739	758	13			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		
3	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is DIETHYL PHOSPHONATE (three-letter code: DEP) (formula: C<sub>4</sub>H<sub>11</sub>O<sub>3</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	1
			8	4	3	1		
4	B	1	Total	C	O	P	0	1
			8	4	3	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is N-hydroxy-1-(1-methylpyridin-2(1H)-ylidene)methanamine (three-letter code: FP1) (formula: C<sub>7</sub>H<sub>10</sub>N<sub>2</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			10	7	2	1		
6	A	1	Total	C	N	O	0	0
			10	7	2	1		
6	B	1	Total	C	N	O	0	0
			10	7	2	1		

- Molecule 7 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	N	O	0	0
			4	1	3		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	329	Total	O	0	0
			329	329		
8	B	267	Total	O	0	0
			267	267		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Acetylcholinesterase



#### • Molecule 1: Acetylcholinesterase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.80Å 104.80Å 323.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.13 – 2.20 47.13 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.13-2.20) 99.3 (47.13-2.20)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.50 (at 2.20Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.181 , 0.224 0.181 , 0.224	Depositor DCC
$R_{free}$ test set	5282 reflections (5.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	40.3	Xtriage
Anisotropy	0.595	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 34.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9173	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, DEP, EDO, FP1, FUC, NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/4339	0.59	0/5930
1	B	0.50	0/4344	0.60	0/5937
All	All	0.50	0/8683	0.59	0/11867

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4207	0	4079	30	0
1	B	4214	0	4085	18	0
2	A	28	0	24	1	0
2	B	42	0	37	1	0
3	A	10	0	10	0	0
3	B	10	0	10	0	0
4	A	8	0	10	5	0
4	B	8	0	10	1	0
5	A	8	0	12	0	0
5	B	8	0	12	0	0
6	A	20	0	20	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	10	0	10	0	0
7	B	4	0	0	0	0
8	A	329	0	0	8	0
8	B	267	0	0	2	0
All	All	9173	0	8319	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:527:GLN:NE2	8:A:701:HOH:O	2.09	0.83
1:B:527:GLN:NE2	8:B:701:HOH:O	2.12	0.83
1:B:4:GLU:HG2	1:B:9:LEU:HD11	1.67	0.75
1:A:295[A]:PHE:HE1	4:A:604[A]:DEP:H21	1.53	0.74
1:A:18[B]:ARG:NH1	8:A:704:HOH:O	2.20	0.73
1:A:30:SER:HB2	1:A:103:THR:HG22	1.74	0.68
1:A:258:PRO:O	8:A:702:HOH:O	2.13	0.65
1:A:295[A]:PHE:CE1	4:A:604[A]:DEP:H21	2.30	0.65
1:A:265:ASN:N	8:A:708:HOH:O	2.35	0.59
1:B:113:PRO:HG2	1:B:485:ARG:HG2	1.82	0.59
1:A:492:PRO:O	1:A:493:ARG:HG3	2.07	0.54
1:A:295[A]:PHE:HE1	4:A:604[A]:DEP:C2	2.21	0.53
1:B:492:PRO:O	1:B:493:ARG:HB2	2.08	0.52
1:B:25:PRO:HG3	1:B:461:PRO:HD3	1.95	0.49
1:B:295[A]:PHE:CE1	4:B:605[A]:DEP:H21	2.48	0.49
1:A:304:ASP:OD1	8:A:703:HOH:O	2.18	0.48
1:B:51:GLU:HG2	1:B:52:PRO:HD2	1.95	0.48
1:B:24:THR:OG1	1:B:140:GLN:HG3	2.13	0.48
1:A:200:PHE:CB	1:A:226:VAL:HB	2.43	0.48
1:A:305:GLY:HA2	1:A:309:SER:HA	1.96	0.48
1:B:347:SER:HB2	2:B:601:NAG:H62	1.96	0.47
1:B:252:ALA:HA	1:B:273:LEU:HD21	1.97	0.46
1:B:328:VAL:O	1:B:427:ALA:HA	2.15	0.46
1:A:103:THR:HG23	1:A:104:PRO:O	2.15	0.46
1:B:200:PHE:CB	1:B:226:VAL:HB	2.45	0.45
1:A:453:PHE:HB3	1:A:476:LEU:HD12	1.99	0.45
1:B:332:LYS:HE3	8:B:755:HOH:O	2.16	0.45
1:B:491:GLU:HB2	1:B:494:ASP:HB3	1.99	0.45
1:B:24:THR:HG22	1:B:136:ARG:HD3	1.99	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:LEU:HD23	1:A:60:VAL:HB	1.99	0.44
1:A:177:ARG:CZ	1:A:217:PRO:HB2	2.48	0.44
1:A:66:GLN:HG3	1:A:98:TYR:CD2	2.54	0.42
1:B:210:GLY:O	1:B:213:LEU:HB2	2.19	0.42
1:A:236:TRP:CZ2	4:A:604[A]:DEP:H23	2.55	0.42
1:A:286:TRP:HB2	6:A:608:FP1:O1	2.19	0.42
1:B:294:VAL:HG23	1:B:295[A]:PHE:HB2	2.02	0.42
6:A:607:FP1:HB	8:A:712:HOH:O	2.19	0.41
1:A:24:THR:HG22	1:A:136:ARG:HD3	2.03	0.41
1:A:295[A]:PHE:HE1	4:A:604[A]:DEP:C1	2.33	0.41
1:A:21:ARG:NH2	8:A:715:HOH:O	2.43	0.41
1:A:161:LEU:HD11	1:A:269:LEU:HD22	2.03	0.41
1:A:322:HIS:HA	1:A:422:GLY:O	2.20	0.41
1:B:294:VAL:HG12	1:B:341:TYR:O	2.21	0.41
1:A:328:VAL:O	1:A:427:ALA:HA	2.21	0.40
1:A:386:LEU:HA	1:A:386:LEU:HD23	1.91	0.40
1:A:141:ALA:HB2	8:A:928:HOH:O	2.21	0.40
1:A:458:PRO:HA	1:A:465:TYR:CD1	2.56	0.40
1:A:202:GLU:HA	1:A:228:GLN:O	2.22	0.40
1:A:347:SER:HB2	2:A:601:NAG:H62	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	535/542 (99%)	515 (96%)	18 (3%)	2 (0%)	39	42
1	B	537/542 (99%)	519 (97%)	16 (3%)	2 (0%)	39	42
All	All	1072/1084 (99%)	1034 (96%)	34 (3%)	4 (0%)	39	42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	493	ARG
1	B	493	ARG
1	B	495	PRO
1	A	495	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	439/437 (100%)	431 (98%)	8 (2%)	66	79
1	B	439/437 (100%)	428 (98%)	11 (2%)	55	67
All	All	878/874 (100%)	859 (98%)	19 (2%)	63	72

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	VAL
1	A	103	THR
1	A	200	PHE
1	A	295[A]	PHE
1	A	295[B]	PHE
1	A	396	GLU
1	A	493	ARG
1	A	541	SER
1	B	21	ARG
1	B	81	GLU
1	B	146	LEU
1	B	200	PHE
1	B	292	GLU
1	B	295[A]	PHE
1	B	295[B]	PHE
1	B	313	GLU
1	B	396	GLU
1	B	413	GLN
1	B	493	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	601	1,3,2	14,14,15	0.61	0	15,19,21	0.38	0
3	FUC	A	602	2	10,10,11	1.14	1 (10%)	13,14,16	0.87	0
2	NAG	A	603	2	14,14,15	0.29	0	15,19,21	0.22	0
4	DEP	A	604[A]	1	2,7,7	0.91	0	2,7,7	0.55	0
5	EDO	A	605	-	3,3,3	0.50	0	2,2,2	0.38	0
5	EDO	A	606	-	3,3,3	0.62	0	2,2,2	0.07	0
6	FP1	A	607	-	9,10,10	4.50	4 (44%)	9,12,12	1.60	3 (33%)
6	FP1	A	608	-	9,10,10	5.07	5 (55%)	9,12,12	2.18	3 (33%)
2	NAG	B	601	1,3,2	14,14,15	0.30	0	15,19,21	0.89	1 (6%)
3	FUC	B	602	2	10,10,11	1.19	2 (20%)	13,14,16	1.27	2 (15%)
2	NAG	B	603	2	14,14,15	0.50	0	15,19,21	0.26	0
2	NAG	B	604	1	14,14,15	0.51	0	15,19,21	0.32	0
4	DEP	B	605[A]	1	2,7,7	0.94	0	2,7,7	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	B	606	-	3,3,3	0.52	0	2,2,2	0.54	0
5	EDO	B	607	-	3,3,3	0.43	0	2,2,2	0.23	0
7	NO3	B	608	-	1,3,3	1.04	0	0,3,3	0.00	-
6	FP1	B	609	-	9,10,10	4.67	5 (55%)	9,12,12	1.37	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1,3,2	-	0/6/23/26	0/1/1/1
3	FUC	A	602	2	-	0/0/17/20	0/1/1/1
2	NAG	A	603	2	-	0/6/23/26	0/1/1/1
4	DEP	A	604[A]	1	-	0/2/6/6	0/0/0/0
5	EDO	A	605	-	-	0/1/1/1	0/0/0/0
5	EDO	A	606	-	-	0/1/1/1	0/0/0/0
6	FP1	A	607	-	-	0/0/3/3	0/1/1/1
6	FP1	A	608	-	-	0/0/3/3	0/1/1/1
2	NAG	B	601	1,3,2	-	0/6/23/26	0/1/1/1
3	FUC	B	602	2	-	0/0/17/20	0/1/1/1
2	NAG	B	603	2	-	0/6/23/26	0/1/1/1
2	NAG	B	604	1	-	0/6/23/26	0/1/1/1
4	DEP	B	605[A]	1	-	0/2/6/6	0/0/0/0
5	EDO	B	606	-	-	0/1/1/1	0/0/0/0
5	EDO	B	607	-	-	0/1/1/1	0/0/0/0
7	NO3	B	608	-	-	0/0/0/0	0/0/0/0
6	FP1	B	609	-	-	0/0/3/3	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	607	FP1	C3-C4	-2.25	1.34	1.38
6	B	609	FP1	C3-C4	-2.19	1.34	1.38
6	A	608	FP1	C3-C4	-2.06	1.34	1.38
6	B	609	FP1	C1-N1	2.03	1.38	1.33
6	A	607	FP1	C2-C3	2.08	1.43	1.38
6	B	609	FP1	C2-C3	2.29	1.43	1.38
3	B	602	FUC	C2-C3	2.31	1.55	1.52
3	A	602	FUC	C2-C3	2.36	1.55	1.52
6	A	608	FP1	C2-C3	2.43	1.44	1.38
3	B	602	FUC	C1-C2	2.62	1.58	1.52

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	608	FP1	C1-N1	2.80	1.40	1.33
6	A	607	FP1	C4-C5	5.68	1.48	1.39
6	B	609	FP1	C4-C5	5.83	1.48	1.39
6	A	608	FP1	C4-C5	6.38	1.49	1.39
6	A	607	FP1	C6-N2	11.45	1.44	1.30
6	B	609	FP1	C6-N2	12.00	1.45	1.30
6	A	608	FP1	C6-N2	13.01	1.46	1.30

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	608	FP1	C4-C5-N1	-3.16	115.02	117.37
6	B	609	FP1	C4-C5-N1	-2.68	115.38	117.37
6	A	607	FP1	C7-N1-C5	-2.62	118.72	120.73
6	A	607	FP1	C4-C5-N1	-2.37	115.61	117.37
6	B	609	FP1	C7-N1-C5	-2.21	119.03	120.73
3	B	602	FUC	O5-C5-C4	2.35	113.65	109.58
6	A	607	FP1	C5-C6-N2	2.70	131.53	127.26
6	A	608	FP1	C3-C4-C5	2.84	122.65	121.26
2	B	601	NAG	C1-O5-C5	2.95	116.47	112.14
3	B	602	FUC	C1-C2-C3	3.08	113.28	109.55
6	A	608	FP1	C5-C6-N2	4.44	134.27	127.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	NAG	1	0
4	A	604[A]	DEP	5	0
6	A	607	FP1	1	0
6	A	608	FP1	1	0
2	B	601	NAG	1	0
4	B	605[A]	DEP	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	533/542 (98%)	-0.39	9 (1%) 73 72	30, 43, 64, 106	0
1	B	538/542 (99%)	-0.29	12 (2%) 65 64	31, 45, 68, 115	0
All	All	1071/1084 (98%)	-0.34	21 (1%) 68 67	30, 44, 66, 115	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	493	ARG	9.4
1	B	495	PRO	7.1
1	B	262	THR	6.4
1	A	497	ALA	5.2
1	A	496	LYS	4.9
1	B	259	PRO	4.6
1	B	492	PRO	4.5
1	B	496	LYS	4.4
1	B	290	PRO	4.1
1	A	495	PRO	4.0
1	A	493	ARG	3.8
1	B	497	ALA	3.5
1	B	494	ASP	3.5
1	A	295[A]	PHE	3.0
1	B	264	GLY	3.0
1	A	492	PRO	2.9
1	A	542	ALA	2.9
1	B	291	GLN	2.7
1	B	295[A]	PHE	2.5
1	A	4	GLU	2.5
1	A	59	VAL	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	A	606	4/4	0.93	0.16	7.66	41,45,48,50	0
5	EDO	B	607	4/4	0.96	0.13	5.41	40,42,44,47	4
6	FP1	A	608	10/10	0.90	0.17	3.33	56,61,65,68	0
5	EDO	B	606	4/4	0.98	0.15	2.53	43,44,44,45	0
7	NO3	B	608	4/4	0.97	0.16	2.29	43,45,47,47	4
4	DEP	B	605[A]	8/8	0.97	0.28	2.11	34,42,48,54	8
6	FP1	A	607	10/10	0.92	0.19	1.81	33,38,45,55	10
4	DEP	A	604[A]	8/8	0.93	0.22	0.99	40,46,51,51	8
5	EDO	A	605	4/4	0.98	0.12	0.88	37,39,40,52	0
6	FP1	B	609	10/10	0.95	0.15	-0.02	35,40,45,49	0
2	NAG	A	603	14/15	0.85	0.38	-	69,94,103,106	0
3	FUC	A	602	10/11	0.92	0.39	-	60,84,89,90	0
2	NAG	B	601	14/15	0.90	0.24	-	60,74,84,85	0
2	NAG	B	603	14/15	0.85	0.32	-	62,84,93,97	0
2	NAG	B	604	14/15	0.88	0.28	-	85,91,95,97	0
3	FUC	B	602	10/11	0.94	0.33	-	70,82,91,93	0
2	NAG	A	601	14/15	0.93	0.23	-	60,69,83,84	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.