



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 22, 2016 – 12:35 AM EDT

PDB ID : 5HJ8  
Title : Bacterial sodium channel neck 3G mutant  
Authors : Rohaim, A.; Minor, D.L.  
Deposited on : 2016-01-13  
Resolution : 3.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0122  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107



ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CA	A	301	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ion transport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	124	Total	C	N	O	S	0	0	0
			948	636	141	165	6			
1	B	118	Total	C	N	O	S	0	0	0
			925	622	139	158	6			
1	C	125	Total	C	N	O	S	0	0	0
			933	626	140	161	6			
1	D	125	Total	C	N	O	S	0	0	0
			933	629	141	157	6			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	137	GLY	-	expression tag	UNP Q0ABW0
A	138	PRO	-	expression tag	UNP Q0ABW0
A	139	SER	-	expression tag	UNP Q0ABW0
A	140	SER	-	expression tag	UNP Q0ABW0
A	141	PRO	-	expression tag	UNP Q0ABW0
A	142	SER	-	expression tag	UNP Q0ABW0
A	248	GLY	ALA	engineered mutation	UNP Q0ABW0
A	249	GLY	GLU	engineered mutation	UNP Q0ABW0
A	250	GLY	ASP	engineered mutation	UNP Q0ABW0
B	137	GLY	-	expression tag	UNP Q0ABW0
B	138	PRO	-	expression tag	UNP Q0ABW0
B	139	SER	-	expression tag	UNP Q0ABW0
B	140	SER	-	expression tag	UNP Q0ABW0
B	141	PRO	-	expression tag	UNP Q0ABW0
B	142	SER	-	expression tag	UNP Q0ABW0
B	248	GLY	ALA	engineered mutation	UNP Q0ABW0
B	249	GLY	GLU	engineered mutation	UNP Q0ABW0
B	250	GLY	ASP	engineered mutation	UNP Q0ABW0
C	137	GLY	-	expression tag	UNP Q0ABW0
C	138	PRO	-	expression tag	UNP Q0ABW0
C	139	SER	-	expression tag	UNP Q0ABW0

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	140	SER	-	expression tag	UNP Q0ABW0
C	141	PRO	-	expression tag	UNP Q0ABW0
C	142	SER	-	expression tag	UNP Q0ABW0
C	248	GLY	ALA	engineered mutation	UNP Q0ABW0
C	249	GLY	GLU	engineered mutation	UNP Q0ABW0
C	250	GLY	ASP	engineered mutation	UNP Q0ABW0
D	137	GLY	-	expression tag	UNP Q0ABW0
D	138	PRO	-	expression tag	UNP Q0ABW0
D	139	SER	-	expression tag	UNP Q0ABW0
D	140	SER	-	expression tag	UNP Q0ABW0
D	141	PRO	-	expression tag	UNP Q0ABW0
D	142	SER	-	expression tag	UNP Q0ABW0
D	248	GLY	ALA	engineered mutation	UNP Q0ABW0
D	249	GLY	GLU	engineered mutation	UNP Q0ABW0
D	250	GLY	ASP	engineered mutation	UNP Q0ABW0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ca 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O 1 1	0	0
3	C	1	Total O 1 1	0	0
3	D	1	Total O 1 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ion transport protein




- Molecule 1: Ion transport protein




- Molecule 1: Ion transport protein



- Molecule 1: Ion transport protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	138.31Å 153.35Å 165.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 3.70 15.00 – 3.70	Depositor EDS
% Data completeness (in resolution range)	98.2 (15.00-3.70) 99.9 (15.00-3.70)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 3.67Å)	Xtriage
Refinement program	REFMAC 5.8.0071	Depositor
R, $R_{free}$	0.278 , 0.304 0.272 , 0.296	Depositor DCC
$R_{free}$ test set	958 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	164.1	Xtriage
Anisotropy	0.659	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 272.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.34$ , $\langle L^2 \rangle = 0.18$	Xtriage
Outliers	2 of 18852 reflections (0.011%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3743	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	316.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	0/972	0.75	0/1327
1	B	0.60	0/948	0.75	0/1290
1	C	0.58	1/954 (0.1%)	0.76	1/1300 (0.1%)
1	D	0.57	0/956	0.73	0/1305
All	All	0.57	1/3830 (0.0%)	0.75	1/5222 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2
1	C	0	2
1	D	0	1
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	173	ALA	C-N	-5.67	1.21	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	173	ALA	O-C-N	-7.57	110.60	122.70

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	276	SER	Peptide
1	B	278	VAL	Peptide
1	C	173	ALA	Mainchain
1	C	281	LEU	Peptide
1	D	284	ARG	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	948	0	901	47	0
1	B	925	0	903	37	0
1	C	933	0	884	37	0
1	D	933	0	883	46	0
2	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	4	0
3	D	1	0	0	0	0
All	All	3743	0	3571	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:273:ASP:O	3:C:301:HOH:O	1.54	1.23
1:D:281:LEU:HA	1:D:284:ARG:CB	1.95	0.97
1:C:277:LYS:N	3:C:301:HOH:O	2.02	0.90
1:C:276:SER:N	3:C:301:HOH:O	2.05	0.88
1:A:234:ILE:HD11	1:D:237:ILE:HA	1.59	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/152 (79%)	101 (84%)	17 (14%)	2 (2%)	11	57
1	B	112/152 (74%)	95 (85%)	14 (12%)	3 (3%)	6	48
1	C	119/152 (78%)	98 (82%)	19 (16%)	2 (2%)	11	57
1	D	119/152 (78%)	101 (85%)	16 (13%)	2 (2%)	11	57
All	All	470/608 (77%)	395 (84%)	66 (14%)	9 (2%)	10	55

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173	ALA
1	C	173	ALA
1	D	173	ALA
1	D	215	TRP
1	B	173	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/129 (72%)	83 (89%)	10 (11%)	8	41
1	B	94/129 (73%)	86 (92%)	8 (8%)	13	53
1	C	89/129 (69%)	83 (93%)	6 (7%)	20	63
1	D	88/129 (68%)	79 (90%)	9 (10%)	9	43
All	All	364/516 (70%)	331 (91%)	33 (9%)	12	49

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	258	GLN
1	C	215	TRP
1	D	266	GLU
1	B	259	ARG
1	B	269	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	269	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	124/152 (81%)	-0.08	6 (4%) 34 22	237, 315, 407, 434	0
1	B	118/152 (77%)	-0.21	2 (1%) 73 58	232, 306, 398, 442	0
1	C	125/152 (82%)	0.18	7 (5%) 28 18	243, 325, 404, 452	0
1	D	125/152 (82%)	-0.05	5 (4%) 42 28	226, 298, 392, 446	0
All	All	492/608 (80%)	-0.04	20 (4%) 41 27	226, 310, 404, 452	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	181	GLY	6.6
1	C	182	THR	5.8
1	C	239	GLU	5.5
1	C	202	GLY	4.4
1	C	201	MET	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	301	1/1	-0.24	0.86	9.51	306,306,306,306	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.