



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:51 PM GMT

PDB ID : 1HM2  
Title : ACTIVE SITE OF CHONDROITINASE AC LYASE REVEALED BY THE  
STRUCTURE OF ENZYME-OLIGOSACCHARIDE COMPLEXES AND  
MUTAGENESIS  
Authors : Huang, W.; Boju, L.; Tkalec, L.; Su, H.; Yang, H.O.; Gunay, N.S.; Linhardt,  
R.J.; Kim, Y.S.; Matte, A.; Cygler, M.  
Deposited on : 2000-12-04  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	700	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	RAM	A	702	X	-	-	-
2	MXY	A	705	X	-	-	-
3	MAN	A	706	-	-	-	X
4	IDR	A	711	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ASG	A	712	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5843 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHONDROITINASE AC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	674	Total	C	N	O	S	0	0	0
			5382	3445	919	1004	14			

- Molecule 2 is a polymer of unknown type called SUGAR (RAM-MAN-GCU-XYS-MXY).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	5	Total	C	O	0	0
			53	30	23		

- Molecule 3 is a polymer of unknown type called SUGAR (RAM-MAN-GCU).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	3	Total	C	O	0	0
			33	18	15		

- Molecule 4 is a polymer of unknown type called SUGAR (DERMATAN SULFATE TETRASACCHARIDE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	4	Total	C	N	O	S	0	0
			61	28	2	29	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	313	Total 313	O 313	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CHONDROITINASE AC



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.90 Å 86.90 Å 192.30 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 20.03 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.00) 92.9 (20.03-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.65 (at 2.01 Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.233 , 0.270 0.239 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtriage
Anisotropy	0.700	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 48566 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5843	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XYP, MXY, CA, RAM, IDR, GCU, ASG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/5520	0.61	1/7487 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	2	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	405	GLU	N-CA-C	-5.17	97.05	111.00

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	702	RAM	C1
2	A	705	MXY	C1

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5382	0	5240	96	0
2	A	53	0	43	1	0
3	A	33	0	25	0	0
4	A	61	0	37	6	0
5	A	1	0	0	0	0
6	A	313	0	0	0	0
All	All	5843	0	5345	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:TYR:H	1:A:217:GLN:HE22	1.11	0.96
1:A:247:ASN:HB2	1:A:311:ILE:HD11	1.64	0.80
1:A:212:TYR:N	1:A:217:GLN:HE22	1.79	0.78
1:A:690:GLN:HB3	1:A:691:GLU:OE2	1.87	0.74
1:A:499:SER:HB3	1:A:520:LEU:HD23	1.69	0.74
1:A:106:PHE:CE2	1:A:149:LYS:HE3	2.24	0.73
1:A:544:GLY:O	1:A:558:VAL:HG22	1.89	0.71
1:A:203:LEU:HD13	1:A:245:LEU:HD22	1.71	0.71
1:A:212:TYR:H	1:A:217:GLN:NE2	1.88	0.69
1:A:127:TRP:HB2	4:A:711:IDR:O3	1.99	0.62
1:A:156:VAL:O	1:A:160:THR:HG23	2.01	0.60
1:A:205:TYR:HB3	1:A:206:PRO:HD3	1.83	0.60
1:A:56:THR:HG23	1:A:63:TRP:CD1	2.38	0.59
1:A:149:LYS:HB2	1:A:149:LYS:NZ	2.19	0.58
1:A:288:ARG:NH2	4:A:711:IDR:O2	2.36	0.58
1:A:114:LYS:NZ	1:A:118:ASP:OD1	2.36	0.58
1:A:509:ILE:HD13	1:A:510:THR:N	2.20	0.57
1:A:368:ARG:HD3	1:A:430:GLN:HA	1.85	0.57
1:A:54:LEU:HD21	1:A:105:VAL:HG22	1.87	0.57
1:A:247:ASN:CB	1:A:311:ILE:HD11	2.35	0.56
1:A:509:ILE:C	1:A:509:ILE:HD13	2.25	0.56
1:A:47:ASP:HB3	1:A:98:HIS:NE2	2.19	0.56
1:A:356:TYR:CD2	1:A:584:PRO:HB2	2.40	0.56
1:A:71:ASP:O	1:A:124:ARG:HG2	2.06	0.56
1:A:181:LEU:O	1:A:184:PHE:HB3	2.06	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:ILE:HB	1:A:285:VAL:HB	1.89	0.55
1:A:499:SER:HB3	1:A:520:LEU:CD2	2.36	0.55
1:A:105:VAL:HA	1:A:108:GLN:HE21	1.72	0.54
4:A:709:ASG:O6	4:A:709:ASG:OSA	2.26	0.54
1:A:56:THR:HG23	1:A:63:TRP:NE1	2.23	0.53
1:A:601:LYS:HE3	1:A:603:LEU:CD2	2.39	0.53
1:A:506:ARG:HG3	1:A:561:ASP:OD1	2.08	0.53
1:A:102:ASP:OD2	1:A:104:LYS:HB3	2.09	0.53
1:A:415:ARG:HD3	1:A:480:ALA:CB	2.38	0.53
1:A:647:ILE:HG12	1:A:656:ILE:HG12	1.91	0.52
1:A:593:LYS:HD3	1:A:593:LYS:O	2.09	0.52
1:A:224:GLN:O	1:A:225:HIS:HB2	2.11	0.51
1:A:415:ARG:HD3	1:A:480:ALA:HB3	1.92	0.51
1:A:506:ARG:HG2	1:A:507:GLY:N	2.26	0.51
1:A:261:LEU:HD13	1:A:261:LEU:C	2.31	0.51
1:A:56:THR:HG23	1:A:63:TRP:HE1	1.75	0.50
1:A:32:MET:HE3	1:A:188:LEU:HB3	1.92	0.50
1:A:228:GLN:O	1:A:230:GLN:HG2	2.12	0.49
1:A:547:PHE:CG	1:A:555:LYS:HG2	2.48	0.49
1:A:133:THR:HB	1:A:134:PRO:HD3	1.95	0.49
1:A:76:TRP:CD1	1:A:123:SER:HB2	2.48	0.48
1:A:403:VAL:CG1	1:A:558:VAL:HG23	2.43	0.48
1:A:426:LEU:HD22	1:A:426:LEU:N	2.28	0.48
1:A:363:VAL:HG22	1:A:364:SER:N	2.29	0.48
1:A:601:LYS:HE3	1:A:603:LEU:HD23	1.96	0.47
1:A:104:LYS:O	1:A:108:GLN:HG3	2.14	0.47
1:A:509:ILE:HB	1:A:540:GLN:NE2	2.29	0.47
1:A:210:VAL:HB	1:A:213:GLU:HG2	1.96	0.47
1:A:106:PHE:CE1	1:A:149:LYS:HG3	2.49	0.47
1:A:145:ARG:HD2	1:A:189:LEU:HD23	1.96	0.47
1:A:106:PHE:CD1	1:A:149:LYS:HG3	2.50	0.47
1:A:506:ARG:HD2	1:A:561:ASP:OD1	2.15	0.47
1:A:127:TRP:CE3	4:A:709:ASG:H5	2.49	0.47
1:A:245:LEU:O	1:A:249:VAL:HG22	2.15	0.47
1:A:57:LEU:HD23	1:A:108:GLN:HB3	1.98	0.46
1:A:637:ILE:HD13	1:A:647:ILE:HD12	1.97	0.46
1:A:297:ASN:HB2	1:A:398:TYR:CE1	2.51	0.46
1:A:169:GLU:CD	1:A:169:GLU:H	2.18	0.46
1:A:603:LEU:HD11	1:A:613:TYR:HB2	1.97	0.45
1:A:261:LEU:HD13	1:A:261:LEU:O	2.15	0.45
1:A:372:SER:HB2	1:A:423:LEU:HD13	1.99	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:MET:CE	1:A:188:LEU:HB3	2.47	0.44
1:A:228:GLN:OE1	1:A:371:GLU:HG2	2.17	0.44
1:A:149:LYS:HB2	1:A:149:LYS:HZ2	1.82	0.44
1:A:529:PHE:CE2	1:A:535:LEU:HD21	2.52	0.44
1:A:252:THR:HB	1:A:253:PRO:HD2	1.99	0.44
1:A:219:ASP:OD2	1:A:343:PHE:HB3	2.18	0.44
1:A:127:TRP:CD1	1:A:131:ILE:HD12	2.53	0.44
1:A:400:ILE:HA	1:A:492:TRP:CZ2	2.54	0.43
1:A:532:GLY:O	1:A:533:ALA:HB2	2.19	0.42
1:A:615:GLN:CD	1:A:615:GLN:C	2.77	0.42
1:A:234:TYR:HE2	4:A:711:IDR:C6	2.32	0.42
4:A:709:ASG:O5	4:A:711:IDR:H3	2.18	0.42
1:A:368:ARG:HD2	1:A:428:GLY:O	2.20	0.42
1:A:473:GLY:HA3	1:A:578:TYR:CE2	2.55	0.42
1:A:204:PHE:CE2	1:A:245:LEU:HD23	2.55	0.42
1:A:32:MET:HE3	1:A:188:LEU:C	2.40	0.42
1:A:120:ASP:N	1:A:121:PRO:HD3	2.34	0.42
1:A:363:VAL:HG22	1:A:364:SER:H	1.85	0.42
1:A:149:LYS:HA	1:A:150:PRO:HD3	1.80	0.41
1:A:105:VAL:HA	1:A:108:GLN:NE2	2.33	0.41
1:A:613:TYR:CE1	1:A:650:ILE:HD11	2.54	0.41
1:A:36:MET:HG3	1:A:146:TYR:CD1	2.55	0.41
1:A:131:ILE:HG23	1:A:178:ASP:HB3	2.02	0.41
1:A:629:LYS:HD3	1:A:636:GLU:OE2	2.20	0.41
1:A:650:ILE:HG13	1:A:651:ASN:N	2.36	0.41
1:A:635:ILE:CA	1:A:675:LEU:HD13	2.51	0.41
1:A:647:ILE:HG22	1:A:650:ILE:HG22	2.02	0.41
1:A:479:ASN:OD1	2:A:701:MAN:H5	2.20	0.40
1:A:293:PRO:HA	1:A:546:TRP:CH2	2.56	0.40
1:A:219:ASP:HB3	1:A:344:TRP:CE2	2.57	0.40
1:A:126:TRP:CH2	1:A:171:LYS:HE2	2.56	0.40
1:A:32:MET:SD	1:A:145:ARG:NE	2.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	672/700 (96%)	630 (94%)	40 (6%)	2 (0%)	46	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	TYR
1	A	148	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	563/598 (94%)	552 (98%)	11 (2%)	63	65

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	HIS
1	A	137	LEU
1	A	149	LYS
1	A	195	LEU
1	A	306	LEU
1	A	401	MET
1	A	404	TRP
1	A	468	GLU
1	A	506	ARG
1	A	509	ILE
1	A	543	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	108	GLN
1	A	208	GLN
1	A	217	GLN
1	A	374	ASN
1	A	540	GLN
1	A	551	ASN
1	A	616	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

12 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MAN	A	701	1,2	11,11,12	0.46	0	14,15,17	0.57	0
2	RAM	A	702	2	10,10,11	1.50	1 (10%)	14,14,16	1.09	1 (7%)
2	GCU	A	703	2	9,12,13	1.32	1 (11%)	13,17,19	0.94	1 (7%)
2	XYP	A	704	2	9,9,10	1.43	1 (11%)	12,12,14	1.31	2 (16%)
2	MXY	A	705	2	11,11,12	1.10	1 (9%)	14,15,17	0.94	0
3	MAN	A	706	1,3	11,11,12	0.50	0	14,15,17	0.64	0
3	RAM	A	707	3	10,10,11	1.21	1 (10%)	14,14,16	1.15	2 (14%)
3	GCU	A	708	3	9,12,13	1.34	1 (11%)	13,17,19	0.99	0
4	ASG	A	709	4	18,18,19	1.26	2 (11%)	21,26,28	1.35	2 (9%)
4	IDR	A	710	4	9,12,13	3.08	6 (66%)	13,17,19	2.04	3 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	IDR	A	711	4	9,12,13	3.08	4 (44%)	13,17,19	2.10	4 (30%)
4	ASG	A	712	4	19,19,19	1.27	2 (10%)	21,28,28	2.01	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	A	701	1,2	-	0/2/19/22	0/1/1/1
2	RAM	A	702	2	1/1/4/5	0/0/17/20	0/1/1/1
2	GCU	A	703	2	-	0/0/21/24	0/1/1/1
2	XYP	A	704	2	-	0/0/14/17	0/1/1/1
2	MXY	A	705	2	1/1/4/5	0/2/19/22	0/1/1/1
3	MAN	A	706	1,3	-	0/2/19/22	0/1/1/1
3	RAM	A	707	3	-	0/0/17/20	0/1/1/1
3	GCU	A	708	3	-	0/0/21/24	0/1/1/1
4	ASG	A	709	4	-	0/11/28/31	0/1/1/1
4	IDR	A	710	4	-	0/0/21/24	0/1/1/1
4	IDR	A	711	4	-	0/0/21/24	0/1/1/1
4	ASG	A	712	4	-	0/11/31/31	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	712	ASG	O4-S	-4.11	1.43	1.57
4	A	709	ASG	O4-S	-3.52	1.45	1.57
3	A	708	GCU	O5-C5	2.37	1.46	1.43
2	A	703	GCU	C4-C5	2.50	1.58	1.53
2	A	705	MXY	C1-C2	2.50	1.55	1.51
3	A	707	RAM	C4-C5	2.50	1.58	1.52
4	A	712	ASG	C1-C2	2.52	1.56	1.53
4	A	709	ASG	O5-C1	2.60	1.48	1.43
4	A	710	IDR	O4-C4	2.64	1.49	1.43
4	A	710	IDR	O3-C3	2.73	1.49	1.43
4	A	711	IDR	O2-C2	2.87	1.49	1.43
4	A	711	IDR	O3-C3	2.91	1.49	1.43
4	A	710	IDR	C2-C3	3.20	1.56	1.52
2	A	702	RAM	C4-C5	3.21	1.59	1.52
4	A	710	IDR	O5-C1	3.68	1.49	1.43
2	A	704	XYP	O5B-C1B	3.80	1.50	1.42
4	A	710	IDR	O2-C2	4.10	1.52	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	711	IDR	O5-C1	5.15	1.52	1.43
4	A	710	IDR	O5-C5	5.33	1.48	1.43
4	A	711	IDR	O5-C5	6.13	1.49	1.43

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	711	IDR	O2-C2-C1	-4.39	100.42	109.21
3	A	707	RAM	C1-C2-C3	-2.87	106.15	109.54
4	A	709	ASG	O7-C7-C8	-2.20	118.02	122.06
2	A	703	GCU	C1-C2-C3	-2.04	107.13	109.54
4	A	711	IDR	O4-C4-C5	2.09	114.34	110.42
3	A	707	RAM	O5-C5-C6	2.13	109.64	106.13
2	A	704	XYP	C1B-C2B-C3B	2.34	112.31	109.54
2	A	702	RAM	C1-O5-C5	2.64	116.45	112.38
4	A	710	IDR	C2-C3-C4	3.04	116.20	111.04
2	A	704	XYP	C4B-C3B-C2B	3.13	114.33	111.24
4	A	711	IDR	C2-C3-C4	3.44	116.89	111.04
4	A	711	IDR	C1-C2-C3	4.13	114.43	109.54
4	A	710	IDR	O2-C2-C1	4.22	117.67	109.21
4	A	709	ASG	C4-O4-S	4.56	127.45	118.77
4	A	710	IDR	C1-C2-C3	4.78	115.19	109.54
4	A	712	ASG	C4-O4-S	8.22	134.43	118.77

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	702	RAM	C1
2	A	705	MXV	C1

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	MAN	1	0
4	A	709	ASG	3	0
4	A	711	IDR	4	0

## 5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	674/700 (96%)	0.29	27 (4%) 42 44	15, 27, 41, 48	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	ALA	7.6
1	A	43	LEU	4.2
1	A	193	GLU	4.1
1	A	593	LYS	3.7
1	A	51	GLU	3.7
1	A	597	GLY	3.6
1	A	652	GLY	3.6
1	A	212	TYR	3.5
1	A	615	GLN	3.4
1	A	26	GLY	3.1
1	A	598	THR	3.0
1	A	109	ILE	2.9
1	A	74	THR	2.8
1	A	96	ASP	2.8
1	A	98	HIS	2.7
1	A	59	PRO	2.6
1	A	95	LYS	2.6
1	A	500	THR	2.6
1	A	509	ILE	2.5
1	A	124	ARG	2.4
1	A	42	PRO	2.4
1	A	502	GLY	2.4
1	A	681	ASN	2.3
1	A	418	LEU	2.3
1	A	253	PRO	2.1
1	A	676	LYS	2.1
1	A	88	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	IDR	A	711	12/13	0.64	0.45	15.74	44,50,51,51	0
4	ASG	A	712	19/19	0.76	0.35	3.43	52,54,57,57	0
3	MAN	A	706	11/12	0.94	0.18	2.69	27,29,31,32	0
4	ASG	A	709	18/19	0.94	0.13	-0.27	33,37,40,40	0
2	GCU	A	703	12/13	0.90	0.13	-	41,42,43,45	0
3	RAM	A	707	10/11	0.92	0.17	-	31,32,33,33	0
4	IDR	A	710	12/13	0.77	0.24	-	42,43,44,45	0
2	XYP	A	704	9/10	0.70	0.17	-	47,49,50,53	0
2	MXY	A	705	11/12	0.58	0.38	-	54,56,56,57	0
2	RAM	A	702	10/11	0.50	0.57	-	52,54,54,55	0
3	GCU	A	708	12/13	0.87	0.23	-	33,34,36,36	0
2	MAN	A	701	11/12	0.80	0.22	-	40,44,45,49	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	CA	A	1801	1/1	0.99	0.03	-2.09	23,23,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.