



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:18 AM GMT

PDB ID : 3HOW  
Title : Complete RNA polymerase II elongation complex III with a T-U mismatch and a frayed RNA 3'-uridine  
Authors : Sydow, J.F.; Brueckner, F.; Cheung, A.C.M.; Damsma, G.E.; Dengl, S.; Lehmann, E.; Vassylyev, D.; Cramer, P.  
Deposited on : 2009-06-03  
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1408 (3.80-3.40)
Clashscore	102246	1010 (3.74-3.46)
Ramachandran outliers	100387	1007 (3.76-3.44)
Sidechain outliers	100360	1007 (3.76-3.44)
RSRZ outliers	91569	1003 (3.78-3.42)
RNA backbone	2183	1058 (4.40-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 29%, yellow 41%, orange 10%, grey 18%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>29%</span> <span>41%</span> <span>10%</span> <span>18%</span> </div> </div>
2	B	1224	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 28%, yellow 49%, orange 13%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>28%</span> <span>49%</span> <span>13%</span> <span>9%</span> </div> </div>
3	C	347	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 20%, yellow 44%, orange 12%, grey 23%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>20%</span> <span>44%</span> <span>12%</span> <span>23%</span> </div> </div>
4	D	221	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 25%, yellow 41%, orange 12%, grey 19%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span></span> <span>25%</span> <span>41%</span> <span>12%</span> <span>19%</span> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	E	215	
6	F	155	
7	G	171	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	2	12	
14	1	26	
15	3	18	

## 2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 31876 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1416	Total	C	N	O	S	0	0	0
			11143	7021	1949	2111	62			

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1112	Total	C	N	O	S	0	0	0
			8838	5594	1550	1639	55			

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	266	Total	C	N	O	S	0	0	0
			2095	1317	348	417	13			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-28	MET	-	EXPRESSION TAG	UNP P16370
C	-27	GLY	-	EXPRESSION TAG	UNP P16370
C	-26	SER	-	EXPRESSION TAG	UNP P16370
C	-25	HIS	-	EXPRESSION TAG	UNP P16370
C	-24	HIS	-	EXPRESSION TAG	UNP P16370
C	-23	HIS	-	EXPRESSION TAG	UNP P16370
C	-22	HIS	-	EXPRESSION TAG	UNP P16370
C	-21	HIS	-	EXPRESSION TAG	UNP P16370
C	-20	HIS	-	EXPRESSION TAG	UNP P16370
C	-19	SER	-	EXPRESSION TAG	UNP P16370
C	-18	ASN	-	EXPRESSION TAG	UNP P16370
C	-17	SER	-	EXPRESSION TAG	UNP P16370
C	-16	GLY	-	EXPRESSION TAG	UNP P16370
C	-15	LEU	-	EXPRESSION TAG	UNP P16370

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-14	ASN	-	EXPRESSION TAG	UNP P16370
C	-13	ASP	-	EXPRESSION TAG	UNP P16370
C	-12	ILE	-	EXPRESSION TAG	UNP P16370
C	-11	PHE	-	EXPRESSION TAG	UNP P16370
C	-10	GLU	-	EXPRESSION TAG	UNP P16370
C	-9	ALA	-	EXPRESSION TAG	UNP P16370
C	-8	GLN	-	EXPRESSION TAG	UNP P16370
C	-7	LYS	-	EXPRESSION TAG	UNP P16370
C	-6	ILE	-	EXPRESSION TAG	UNP P16370
C	-5	GLU	-	EXPRESSION TAG	UNP P16370
C	-4	TRP	-	EXPRESSION TAG	UNP P16370
C	-3	HIS	-	EXPRESSION TAG	UNP P16370
C	-2	GLU	-	EXPRESSION TAG	UNP P16370
C	-1	ASP	-	EXPRESSION TAG	UNP P16370
C	0	THR	-	EXPRESSION TAG	UNP P16370
C	1	GLY	-	EXPRESSION TAG	UNP P16370

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	178	Total	C	N	O	S	0	0	0
			1434	887	257	288	2			

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	87	Total	C	N	O	S	0	0	0
			705	451	119	132	3			

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	171	Total	C	N	O	S	0	0	0
			1340	861	222	249	8			

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	134	Total	C	N	O	S	0	0	0
			1076	677	182	213	4			

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	114	Total	C	N	O	S	0	0	0
			927	571	168	178	10			

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	65	Total	C	N	O	S	0	0	0
			532	339	93	94	6			

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	114	Total	C	N	O	S	0	0	0
			919	590	156	171	2			

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	44	Total	C	N	O	S	0	0	0
			351	217	70	60	4			

- Molecule 13 is a DNA chain called 5'-D(\*AP\*CP\*TP\*AP\*CP\*TP\*TP\*GP\*AP\*GP\*CP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	2	7	Total	C	N	O	P	0	0	0
			137	68	22	41	6			

- Molecule 14 is a DNA chain called 5'-D(\*AP\*GP\*CP\*TP\*C\*AP\*AP\*GP\*TP\*AP\*GP\*TP\*TP\*AP\*TP\*GP\*CP\*CP\*(BRU)P\*GP\*GP\*TP\*CP\*AP\*TP\*T)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
14	1	19	Total	Br	C	N	O	P	0	0	0
			389	1	186	71	113	18			

- Molecule 15 is a RNA chain called 5'-R(\*UP\*GP\*CP\*AP\*UP\*UP\*U\*CP\*AP\*AP\*CP\*CP

\*AP\*GP\*GP\*CP\*UP\*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	3	11	Total	C	N	O	P	0	0	0
			229	104	41	74	10			

- Molecule 16 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	J	1	Total	Zn	0	0
			1	1		
16	B	1	Total	Zn	0	0
			1	1		
16	I	2	Total	Zn	0	0
			2	2		
16	C	1	Total	Zn	0	0
			1	1		
16	A	2	Total	Zn	0	0
			2	2		
16	L	1	Total	Zn	0	0
			1	1		


- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	1	Total	Mg	0	0
			1	1		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed RNA polymerase II subunit RPB1






PRO	LYS	PRO	LEU	L1409	I1333	M1267	D1206	H1140	E1074	G1002	S915	V842	Q767
ALA	GLN	VAL	VAL	F1410	D1334	L1288	L1207	T1141	P1075	K1003	G916	K843	Q769
TYR	ASN	ASN	GLY		I1335	E1289	M1209	T1142	A1076	M1004	E1919	A844	K773
SER	ALA	PHE	ASP	A1414	M1336		M1208	S1145	M1079	E1005	E1920	E846	R774
PRO	GLU	SER	ASP	S1415	V1337	T1272	Q1211	T1146	T1080	E1006	E1921		I775
THR	THR	PRO	THR	A1416	E1338	L11273	Q1212	T1147	L1081	M1009	D922		A776
SER	TYR	THR	ASP		L1339	R1274	V1212	T1148	ASN	A1010	E1923		F777
PRO	VAL	SER	VAL	D1419	G1340	E1277	G1213	I1148	THR	Q1011	E1924	H851	F778
ASN	PRO	PRO	GLY	D1420	I1341	E1278	E1214	A1149		E1016	E1925		F779
GLU	TYR	THR	GLU	C1421	E1342	M1278	R1215	S1150	PHE	Q1011	E1926	D853	V780
SER	ASN	THR	TYR	R1422	A1343	I1279	I1216	E1151	HIS		Q926	D854	D781
GLU	PRO	PRO	LEU		G1344	E1280	K1217	I1152	PHE	V1015	L929	T855	R782
ASN	THR	PRO	MET	S1425	R1345	R1281	Q1218	I1153	ALA	T1016		T856	T783
SER	SER	THR	PHE	E1426		V1282	T1219	T1154	GLY	L1017		R857	T785
PRO	PRO	PRO	SER	L1348		V1283	F1220	D1155	VAL	F1018	K934	R858	L784
SER	PRO	PRO	PRO	I1349		M1284	K1221	P1158	ALA	C1019	Q935	P785	P785
TYR	THR	ALA	LEU	K1350		M1285	M1222	P1159	SER	C1020	E1937	S859	H786
SER	VAL	VAL	VAL	E1351		D1286	D1223	R1160	K1092	L1021	L860	L860	G861
PRO	TYR	TYR	ASP	L1352		Y1287	L1224	T1161	K1093	L1022	G861	G862	K789
THR	PRO	PRO	PRO	Y1353		D1288	F1225	T1162	V1094	R1023	D939	F866	S793
SER	THR	THR	GLY	V1355		R1289	V1226	V1163	T1095	R1024	E1941	Y868	E795
PRO	SER	SER	SER			K1290	I1227	I1163		A1027	F942	Y868	S796
TYR	ASP	ASP	ASP	Y1362		P1292	S1229	P1164	V1098	T1028	E1943	G869	K797
ALA	ALA	TYR	ALA	L1363		S1293	D1230	D1166	R1100	R1030	Y946	E870	G798
MET	MET	SER	PRO	M1364		P1294	D1231	E1167		R1031	F947	D871	F799
PRO	ALA	PRO	ALA	Y1365		T1295	M1232	E1168	T1104	V1031		G872	W800
GLY	TYR	THR	GLY	G1366		G1296	D1233	T1169	L1105	E1034	A952	M873	E801
SER	GLY	SER	SER	H1367		E1297	L1236	I1170	M1106	Y1035	N953	D874	N802
PRO	PHE	PRO	PHE	M1368		Y1298	I1237	Q1171	V1107	R1036	Y954	A875	S803
THR	THR	THR	THR	A1369		V1299	I1237	L1172		L1037	P955		Y804
SER	TYR	SER	TYR	L1370		K1300	I1238		M1110	T1038	Y958	E879	L805
ALA	ALA	TYR	ALA	L1371		E1301	R1239	S1175	M1111	T1039	Y958	L883	R806
PRO	TYR	SER	TYR	D1372		K1303	C1240	LEU	K1112	Q1040	R961	D884	T809
GLY	GLY	PRO	GLY	Y1373		E1304	R1241	LEU	T1113	A1041	R962	T885	
PRO	ALA	SER	ALA	V1374		V1305	V1242	ASP	S1115	F1042	Y963	I886	E812
ASN	ASN	PRO	ASP	M1375		L1306	V1243	GLU	L1116		Q964	G887	F815
TYR	TYR	SER	SER	T1376		E1387	R1244	GLU	T1117	L1046	Y965	G888	H816
SER	TYR	TYR	TYR	T1377			PRO	ALA	Y1118	S1047	N1048	S889	D890
PRO	GLY	SER	GLY	T1381		D1309	LYS	GLU	T1118	N1049	A967	D890	A891
THR	ILE	PRO	ALA			G1310	SER	GLN	Y1119	E1050	Q968		K821
THR	THR	THR	THR			V1311	LEU	SER	L1120		Q969		D826
SER	GLU	SER	SER	T1385		M1312	ASP	PHE	E1121	F1053	R896	Y897	T827
PRO	PRO	PRO	PRO	H1386		S1314	ALA	ASP	E1123	L1054	R898	R898	A828
TYR	GLU	SER	PHE	H1387		E1315	GLU	Q1187	G1123	R1056	V899	V899	X820
GLY	ASP	TYR	GLY			V1316	THR	S1189	H1124	S1057	D900	X820	X820
ALA	GLY	PRO	ALA	R1391			GLU			V1058	L901	L901	T831
THR	GLY	PRO	THR			V1319	A1254	L1192	D1127	P1090	L902	L902	A832
PRO	GLY	THR	GLY	T1394		P1320	E1255	L1193	Q1128	P1099	N903	N903	T834
GLY	GLY	PRO	SER	G1395		G1321	E1256	R1194	E1129	P1099	T982	T982	E833
VAL	VAL	PRO	VAL	A1396		D1323	H1288	L1195	Q1130	P1099	K984	T904	T834
THR	THR	THR	THR	L1397		M1259	M1259	L1195	K1132	M1063	D985	D905	G835
PRO	PRO	PRO	PRO	G1400		K1324	K1260	D1198	L1133	V1064	H906	H906	Y836
TYR	TYR	PRO	PRO	S1401		T1325	K1261	R1199	T1134		Y987	T907	L837
GLY	GLY	TYR	TYR	F1402		K1262	K1262	R1135	R1135	L1087	L908	L908	Q838
SER	SER	THR	PHE	E1403		T1329	I1263	S1136	S1136		D992	L913	R840
PRO	ALA	PRO	PHE	E1404		M1330	E1264	A1137	I1138	I1072	R840	R840	R840
TYR	TYR	THR	VAL	S1331		M1265	M1265	D1204	I1138	G1073	L993	L914	L841
SER	SER	TYR	SER	T1405		F1332	T1266	K1205	E1139				

• Molecule 2: DNA-directed RNA polymerase II subunit RPB2









• Molecule 3: DNA-directed RNA polymerase II subunit RPB3

Chain C: 20% 44% 12% 23%




• Molecule 4: DNA-directed RNA polymerase II subunit RPB4

Chain D: 25% 41% 12% 19%




• Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1

Chain E: 24% 62% 11%



• Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2

Chain F: 17% 30% 8% 44%




• Molecule 7: DNA-directed RNA polymerase II subunit RPB7

Chain G: 40% 51% 8%




• Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3


Chain H: 21% 50% 18% 8%




- Molecule 9: DNA-directed RNA polymerase II subunit RPB9




- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5




- Molecule 11: DNA-directed RNA polymerase II subunit RPB11



- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4




- Molecule 13: 5'-D(\*AP\*CP\*TP\*AP\*CP\*TP\*TP\*GP\*AP\*GP\*CP\*T)-3'




- Molecule 14: 5'-D(\*AP\*GP\*CP\*TP\*C\*AP\*AP\*GP\*TP\*AP\*GP\*TP\*TP\*AP\*TP\*GP\*CP\*C P\*(BRU)P\*GP\*GP\*TP\*CP\*AP\*TP\*T)-3'





● Molecule 15: 5'-R(\*UP\*GP\*CP\*AP\*UP\*UP\*U\*CP\*AP\*AP\*CP\*CP\*AP\*GP\*GP\*CP\*UP\*U)-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	222.65Å 395.96Å 283.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.60 48.95 – 3.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-3.60) 99.9 (48.95-3.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.74 (at 3.57Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.214 , 0.254 0.212 , 0.249	Depositor DCC
$R_{free}$ test set	2843 reflections (2.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	91.4	Xtriage
Anisotropy	0.061	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 91.3	EDS
Estimated twinning fraction	0.011 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.018 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 144004 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	31876	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, BRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.53	0/11342	0.80	4/15337 (0.0%)
2	B	0.51	0/9009	0.75	2/12146 (0.0%)
3	C	0.53	0/2133	0.78	1/2891 (0.0%)
4	D	0.48	0/1444	0.76	2/1935 (0.1%)
5	E	0.50	0/1788	0.70	0/2406
6	F	0.60	0/717	0.85	0/967
7	G	0.54	0/1368	0.81	0/1844
8	H	0.48	0/1094	0.77	1/1481 (0.1%)
9	I	0.46	0/945	0.74	0/1273
10	J	0.53	0/541	0.88	0/727
11	K	0.54	0/937	0.72	0/1265
12	L	0.45	0/353	0.77	0/468
13	2	0.68	0/152	0.89	0/232
14	1	0.59	0/413	0.80	0/634
15	3	0.60	0/255	0.89	0/395
All	All	0.52	0/32491	0.78	10/44001 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	C	0	1
14	1	0	3
All	All	0	4

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	26	THR	N-CA-C	-6.09	94.56	111.00
1	A	56	PRO	N-CA-C	-6.08	96.28	112.10
1	A	55	ASP	N-CA-CB	6.07	121.52	110.60
1	A	311	GLN	N-CA-C	5.98	127.14	111.00
2	B	43	LEU	CA-CB-CG	-5.82	101.91	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
14	1	18	DA	Sidechain
14	1	20	DG	Sidechain
14	1	21	DC	Sidechain
3	C	229	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11143	0	11217	1115	0
2	B	8838	0	8870	1052	0
3	C	2095	0	2051	258	0
4	D	1434	0	1460	178	0
5	E	1752	0	1776	175	0
6	F	705	0	731	64	0
7	G	1340	0	1357	142	0
8	H	1076	0	1046	152	0
9	I	927	0	882	122	0
10	J	532	0	542	98	0
11	K	919	0	929	98	0
12	L	351	0	376	59	0
13	2	137	0	82	12	0
14	1	389	0	214	16	0
15	3	229	0	121	12	0
16	A	2	0	0	0	0
16	B	1	0	0	0	0
16	C	1	0	0	0	0
16	I	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	J	1	0	0	0	0
16	L	1	0	0	0	0
17	A	1	0	0	0	0
All	All	31876	0	31654	3273	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 52.

The worst 5 of 3273 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:882:THR:HG21	2:B:935:ARG:HA	1.24	1.19
1:A:53:LEU:HD23	1:A:54:ASN:N	1.59	1.18
4:D:71:LYS:HA	4:D:74:GLN:HB2	1.23	1.13
7:G:26:LEU:HD12	7:G:56:ILE:HD11	1.22	1.11
1:A:567:LYS:CD	1:A:568:PRO:HD2	1.80	1.11

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1406/1733 (81%)	1004 (71%)	279 (20%)	123 (9%)	1	14
2	B	1094/1224 (89%)	774 (71%)	202 (18%)	118 (11%)	0	10
3	C	264/347 (76%)	181 (69%)	48 (18%)	35 (13%)	0	6
4	D	174/221 (79%)	124 (71%)	33 (19%)	17 (10%)	1	12
5	E	212/215 (99%)	144 (68%)	46 (22%)	22 (10%)	1	11
6	F	85/155 (55%)	72 (85%)	8 (9%)	5 (6%)	2	24
7	G	169/171 (99%)	137 (81%)	24 (14%)	8 (5%)	3	30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
8	H	130/146 (89%)	84 (65%)	27 (21%)	19 (15%)	0	5
9	I	112/122 (92%)	77 (69%)	26 (23%)	9 (8%)	1	16
10	J	63/70 (90%)	38 (60%)	14 (22%)	11 (18%)	0	3
11	K	112/120 (93%)	86 (77%)	22 (20%)	4 (4%)	4	39
12	L	42/70 (60%)	20 (48%)	13 (31%)	9 (21%)	0	2
All	All	3863/4594 (84%)	2741 (71%)	742 (19%)	380 (10%)	1	12

5 of 380 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	48	ALA
1	A	57	ARG
1	A	58	LEU
1	A	67	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1239/1520 (82%)	1091 (88%)	148 (12%)	6	34
2	B	964/1061 (91%)	846 (88%)	118 (12%)	6	32
3	C	234/299 (78%)	207 (88%)	27 (12%)	7	36
4	D	160/200 (80%)	136 (85%)	24 (15%)	3	25
5	E	196/197 (100%)	176 (90%)	20 (10%)	9	42
6	F	77/137 (56%)	67 (87%)	10 (13%)	5	30
7	G	152/152 (100%)	141 (93%)	11 (7%)	18	58
8	H	118/128 (92%)	99 (84%)	19 (16%)	3	21
9	I	108/116 (93%)	96 (89%)	12 (11%)	8	38
10	J	60/65 (92%)	54 (90%)	6 (10%)	9	43
11	K	99/102 (97%)	86 (87%)	13 (13%)	5	30

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
12	L	39/57 (68%)	35 (90%)	4 (10%)	9	42
All	All	3446/4034 (85%)	3034 (88%)	412 (12%)	6	33

5 of 412 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	466	TRP
2	B	894	ASP
9	I	34	TYR
2	B	505	ASP
2	B	714	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 104 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	821	GLN
2	B	1084	GLN
8	H	133	ASN
2	B	842	ASN
2	B	957	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
15	3	10/18 (55%)	2 (20%)	1 (10%)

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
15	3	3	A
15	3	11	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
15	3	2	A

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
14	BRU	1	23	15,14	13,21,22	4.68	4 (30%)	16,30,33	4.09	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	BRU	1	23	15,14	-	0/3/21/22	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	1	23	BRU	BR-C5	-15.06	1.50	1.90
14	1	23	BRU	C6-N1	3.19	1.39	1.35
14	1	23	BRU	C4-N3	3.53	1.39	1.33
14	1	23	BRU	C4-C5	5.75	1.45	1.38

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	1	23	BRU	C5-C4-N3	-8.15	115.30	124.00
14	1	23	BRU	C5-C6-N1	2.03	123.78	119.79
14	1	23	BRU	C4-N3-C2	13.74	127.13	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
14	1	23	BRU	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1416/1733 (81%)	-0.17	17 (1%) 81 69	15, 63, 100, 128	0
2	B	1112/1224 (90%)	-0.14	16 (1%) 78 65	15, 74, 111, 122	0
3	C	266/347 (76%)	-0.19	1 (0%) 93 88	31, 63, 93, 111	0
4	D	178/221 (80%)	-0.02	1 (0%) 90 83	40, 75, 106, 115	0
5	E	214/215 (99%)	0.13	1 (0%) 91 86	41, 87, 115, 124	0
6	F	87/155 (56%)	-0.48	0 100 100	17, 44, 73, 84	0
7	G	171/171 (100%)	-0.13	1 (0%) 90 83	44, 64, 92, 100	0
8	H	134/146 (91%)	0.40	1 (0%) 89 81	67, 95, 112, 117	0
9	I	114/122 (93%)	0.09	2 (1%) 71 58	62, 93, 109, 114	0
10	J	65/70 (92%)	-0.37	0 100 100	41, 60, 85, 92	0
11	K	114/120 (95%)	-0.14	0 100 100	28, 66, 83, 92	0
12	L	44/70 (62%)	0.14	0 100 100	42, 100, 111, 114	0
13	2	7/12 (58%)	1.47	1 (14%) 4 3	112, 116, 123, 124	0
14	1	18/26 (69%)	0.83	4 (22%) 1 1	68, 109, 126, 131	0
15	3	11/18 (61%)	0.48	1 (9%) 11 8	92, 97, 123, 126	0
All	All	3951/4650 (84%)	-0.11	46 (1%) 81 69	15, 70, 109, 131	0

The worst 5 of 46 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	253	ASN	3.9
1	A	1455	PRO	3.6
1	A	255	SER	3.5
1	A	149	GLU	3.2
15	3	1	C	3.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
14	BRU	1	23	20/21	0.80	0.26	-	81,90,95,98	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
16	ZN	A	2457	1/1	1.00	0.11	-1.45	35,35,35,35	0
17	MG	A	2458	1/1	0.96	0.13	-1.88	52,52,52,52	0
16	ZN	J	1066	1/1	0.99	0.17	-1.94	39,39,39,39	0
16	ZN	C	1269	1/1	1.00	0.07	-2.31	27,27,27,27	0
16	ZN	I	1121	1/1	0.99	0.09	-2.37	74,74,74,74	0
16	ZN	I	1122	1/1	0.97	0.11	-2.80	124,124,124,124	0
16	ZN	L	1071	1/1	0.97	0.07	-2.90	86,86,86,86	0
16	ZN	A	2456	1/1	0.96	0.06	-3.60	70,70,70,70	0
16	ZN	B	2225	1/1	0.99	0.16	-	32,32,32,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.