



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:56 PM GMT

PDB ID : 1HXS
Title : CRYSTAL STRUCTURE OF MAHONEY STRAIN OF POLIOVIRUS AT 2.2A RESOLUTION
Authors : Miller, S.T.; Hogle, J.M.; Filman, D.J.
Deposited on : 2001-01-16
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	1	302	
2	2	272	
3	3	237	
4	4	68	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7256 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GENOME POLYPROTEIN, COAT PROTEIN VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1	288	Total	C	N	O	S	0	0	0
			2250	1431	383	431	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	6	GLY	LEU	SEE REMARK 999	UNP P03300
1	7	SER	GLU	SEE REMARK 999	UNP P03300
1	9	SER	MET	SEE REMARK 999	UNP P03300
1	10	THR	ILE	SEE REMARK 999	UNP P03300

- Molecule 2 is a protein called GENOME POLYPROTEIN, COAT PROTEIN VP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	2	267	Total	C	N	O	S	0	0	0
			2075	1312	357	392	14			

- Molecule 3 is a protein called GENOME POLYPROTEIN, COAT PROTEIN VP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	3	235	Total	C	N	O	S	0	0	0
			1834	1169	299	349	17			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
3	123	SER	PHE	SEE REMARK 999	UNP P03300

- Molecule 4 is a protein called GENOME POLYPROTEIN, COAT PROTEIN VP4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	4	68	Total	C	N	O	S	0	0	0
			519	319	91	108	1			

- Molecule 5 is MYRISTIC ACID (three-letter code: MYR) (formula: $C_{14}H_{28}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	4	1	Total	C	O	0	0
			15	14	1		

- Molecule 6 is PALMITIC ACID (three-letter code: PLM) (formula: $C_{16}H_{32}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	1	1	Total	C	O	0	0
			18	16	2		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	1	213	Total	O	0	0
			213	213		
7	2	136	Total	O	0	0
			136	136		
7	3	144	Total	O	0	0
			144	144		
7	4	52	Total	O	0	0
			52	52		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: GENOME POLYPROTEIN, COAT PROTEIN VP1




• Molecule 2: GENOME POLYPROTEIN, COAT PROTEIN VP2



• Molecule 3: GENOME POLYPROTEIN, COAT PROTEIN VP3



• Molecule 4: GENOME POLYPROTEIN, COAT PROTEIN VP4



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	320.50Å 355.25Å 377.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.20)	Depositor
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.268 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7256	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLM, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1	0.78	0/2312	1.49	26/3160 (0.8%)
2	2	0.79	0/2132	1.46	29/2916 (1.0%)
3	3	0.82	2/1881 (0.1%)	1.33	18/2562 (0.7%)
4	4	0.91	0/528	1.57	7/714 (1.0%)
All	All	0.81	2/6853 (0.0%)	1.44	80/9352 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	2	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	3	102	GLU	CB-CG	8.28	1.67	1.52
3	3	102	GLU	CD-OE2	6.29	1.32	1.25

All (80) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1	83	ARG	NE-CZ-NH2	-20.74	109.93	120.30
1	1	83	ARG	NE-CZ-NH1	18.22	129.41	120.30
2	2	264	ARG	NE-CZ-NH2	-14.08	113.26	120.30
3	3	149	MET	CA-CB-CG	9.73	129.84	113.30
2	2	264	ARG	NE-CZ-NH1	8.97	124.78	120.30
2	2	38	TRP	CD1-CG-CD2	8.15	112.82	106.30
2	2	201	ARG	NE-CZ-NH1	8.09	124.34	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	2	227	TRP	CD1-CG-CD2	8.07	112.75	106.30
3	3	170	TRP	CD1-CG-CD2	7.97	112.67	106.30
3	3	170	TRP	CE2-CD2-CG	-7.91	100.97	107.30
2	2	80	TRP	CD1-CG-CD2	7.79	112.53	106.30
1	1	170	TRP	CD1-CG-CD2	7.79	112.53	106.30
1	1	269	TRP	CD1-CG-CD2	7.69	112.45	106.30
1	1	175	TRP	CE2-CD2-CG	-7.62	101.20	107.30
1	1	269	TRP	CE2-CD2-CG	-7.57	101.25	107.30
1	1	175	TRP	CD1-CG-CD2	7.54	112.33	106.30
2	2	78	TRP	CD1-CG-CD2	7.53	112.33	106.30
2	2	227	TRP	CE2-CD2-CG	-7.53	101.28	107.30
2	2	201	ARG	NE-CZ-NH2	-7.50	116.55	120.30
2	2	80	TRP	CE2-CD2-CG	-7.46	101.33	107.30
1	1	170	TRP	CE2-CD2-CG	-7.46	101.33	107.30
2	2	78	TRP	CE2-CD2-CG	-7.46	101.34	107.30
4	4	67	MET	CG-SD-CE	-7.42	88.33	100.20
1	1	108	TRP	CD1-CG-CD2	7.30	112.14	106.30
2	2	141	MET	CG-SD-CE	-7.28	88.55	100.20
3	3	197	ARG	NE-CZ-NH1	7.26	123.93	120.30
1	1	108	TRP	CE2-CD2-CG	-7.18	101.55	107.30
3	3	156	TRP	CD1-CG-CD2	7.13	112.01	106.30
2	2	79	TRP	CE2-CD2-CG	-7.12	101.60	107.30
3	3	156	TRP	CE2-CD2-CG	-7.11	101.61	107.30
2	2	71	TRP	CE2-CD2-CG	-7.09	101.63	107.30
1	1	258	ARG	NE-CZ-NH1	7.08	123.84	120.30
2	2	38	TRP	CE2-CD2-CG	-7.00	101.70	107.30
2	2	79	TRP	CD1-CG-CD2	7.00	111.90	106.30
2	2	71	TRP	CD1-CG-CD2	6.94	111.86	106.30
3	3	110	TRP	CD1-CG-CD2	6.84	111.78	106.30
3	3	110	TRP	CE2-CD2-CG	-6.75	101.90	107.30
1	1	83	ARG	CG-CD-NE	-6.69	97.75	111.80
1	1	129	ARG	NE-CZ-NH1	6.50	123.55	120.30
3	3	102	GLU	CG-CD-OE1	-6.50	105.31	118.30
4	4	34	ARG	NE-CZ-NH1	6.41	123.51	120.30
2	2	221	MET	CG-SD-CE	-6.37	90.01	100.20
4	4	61	LEU	CA-CB-CG	6.25	129.68	115.30
3	3	170	TRP	CG-CD2-CE3	6.21	139.49	133.90
3	3	170	TRP	CB-CG-CD1	-6.10	119.07	127.00
3	3	102	GLU	CG-CD-OE2	6.07	130.44	118.30
1	1	286	TYR	CB-CG-CD2	-5.94	117.44	121.00
4	4	18	ARG	NE-CZ-NH2	-5.93	117.34	120.30
3	3	177	ARG	NE-CZ-NH1	5.87	123.23	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	1	175	TRP	CG-CD2-CE3	5.76	139.09	133.90
2	2	100	TYR	CB-CG-CD2	-5.75	117.55	121.00
2	2	167	GLN	CA-CB-CG	-5.72	100.81	113.40
2	2	87	ARG	NE-CZ-NH1	5.68	123.14	120.30
3	3	149	MET	N-CA-CB	-5.63	100.47	110.60
2	2	246	GLU	CA-CB-CG	5.58	125.68	113.40
3	3	85	LEU	CA-CB-CG	5.55	128.07	115.30
1	1	269	TRP	CB-CG-CD1	-5.53	119.81	127.00
2	2	222	VAL	CG1-CB-CG2	-5.53	102.05	110.90
2	2	38	TRP	CG-CD1-NE1	-5.49	104.61	110.10
3	3	149	MET	CG-SD-CE	5.48	108.96	100.20
4	4	18	ARG	CA-CB-CG	5.46	125.40	113.40
2	2	227	TRP	CG-CD2-CE3	5.43	138.79	133.90
2	2	78	TRP	CG-CD2-CE3	5.42	138.78	133.90
1	1	108	TRP	CG-CD2-CE3	5.35	138.71	133.90
2	2	78	TRP	CB-CG-CD1	-5.31	120.09	127.00
1	1	272	ARG	NE-CZ-NH1	5.31	122.96	120.30
2	2	62	ARG	NE-CZ-NH1	5.30	122.95	120.30
1	1	24	ARG	NE-CZ-NH1	5.29	122.94	120.30
3	3	223	ARG	NE-CZ-NH1	5.28	122.94	120.30
4	4	34	ARG	NE-CZ-NH2	-5.27	117.66	120.30
1	1	170	TRP	CG-CD2-CE3	5.20	138.58	133.90
1	1	70	ARG	NE-CZ-NH1	5.19	122.89	120.30
1	1	83	ARG	CA-CB-CG	5.16	124.75	113.40
1	1	83	ARG	CD-NE-CZ	5.15	130.81	123.60
4	4	33	TYR	CB-CG-CD2	-5.10	117.94	121.00
2	2	227	TRP	CG-CD1-NE1	-5.06	105.04	110.10
3	3	110	TRP	CG-CD2-CE3	5.05	138.45	133.90
1	1	127	TYR	CB-CG-CD2	-5.02	117.99	121.00
1	1	232	ALA	O-C-N	-5.02	114.67	122.70
1	1	256	LYS	CA-CB-CG	-5.02	102.36	113.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	2	11	ASP	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2250	0	2197	16	0
2	2	2075	0	1994	8	0
3	3	1834	0	1816	18	0
4	4	519	0	497	5	0
5	4	15	0	27	0	0
6	1	18	0	31	5	0
7	1	213	0	0	2	0
7	2	136	0	0	1	0
7	3	144	0	0	1	0
7	4	52	0	0	0	0
All	All	7256	0	6562	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:1:158:MET:SD	1:1:177:THR:HG23	2.23	0.79
1:1:159:TYR:HB2	6:1:2000:PLM:HE2	1.72	0.70
1:1:237:PHE:CE2	6:1:2000:PLM:H21	2.32	0.64
1:1:177:THR:HG22	1:1:180:ASN:HB2	1.79	0.63
4:4:14:GLU:HB3	4:4:20:TYR:CD2	2.33	0.63
1:1:107:VAL:HG13	1:1:239:ILE:HD13	1.80	0.63
1:1:144:GLU:HG2	7:1:2107:HOH:O	1.99	0.61
3:3:72:LEU:HB2	3:3:208:MET:CE	2.30	0.61
1:1:177:THR:HG21	1:1:182:SER:OG	2.01	0.59
3:3:97:HIS:HA	3:3:102:GLU:HG2	1.85	0.59
3:3:72:LEU:HB2	3:3:208:MET:HE2	1.85	0.58
1:1:235:ASN:HA	1:1:237:PHE:CZ	2.39	0.58
7:2:1049:HOH:O	3:3:124:MET:HG2	2.06	0.55
1:1:191:PRO:HG2	3:3:13:TYR:HB2	1.88	0.55
1:1:273:PRO:HB3	2:2:189:ASN:HB2	1.91	0.52
2:2:213:VAL:HG22	3:3:37:PRO:HG2	1.90	0.52
1:1:237:PHE:HE2	6:1:2000:PLM:H21	1.76	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:3:85:LEU:HD21	3:3:95:LEU:HD11	1.94	0.50
1:1:132:MET:SD	6:1:2000:PLM:H62	2.52	0.49
1:1:282:PRO:HG2	2:2:141:MET:HE1	1.94	0.49
3:3:87:LEU:HD13	3:3:190:ILE:HD11	1.94	0.49
2:2:37:ARG:HG3	3:3:37:PRO:HB3	1.94	0.49
3:3:167:VAL:O	3:3:169:PRO:HD3	2.12	0.49
1:1:237:PHE:CE2	6:1:2000:PLM:H52	2.48	0.48
3:3:97:HIS:ND1	3:3:102:GLU:OE2	2.43	0.48
4:4:22:GLY:O	4:4:24:THR:N	2.47	0.48
3:3:234:LYS:HD3	3:3:234:LYS:HA	1.66	0.47
2:2:30:ASN:HD21	4:4:59:ASP:HB2	1.80	0.46
3:3:102:GLU:OE2	3:3:230:HIS:ND1	2.46	0.46
1:1:96:ALA:HA	1:1:249:ASN:O	2.16	0.46
3:3:146:LYS:HB2	7:3:1072:HOH:O	2.15	0.45
1:1:149:HIS:HB3	7:1:2174:HOH:O	2.18	0.44
2:2:187:LEU:O	2:2:187:LEU:HG	2.17	0.43
4:4:6:SER:HB2	4:4:27:TYR:CE1	2.54	0.43
3:3:145:ARG:O	3:3:149:MET:HB3	2.18	0.43
3:3:120:PHE:HA	3:3:210:ILE:HG22	2.01	0.43
2:2:143:THR:HG23	2:2:173:ARG:HA	2.01	0.41
3:3:64:THR:O	3:3:67:MET:HG2	2.20	0.41
3:3:114:LEU:HA	3:3:114:LEU:HD23	1.85	0.41
2:2:70:SER:OG	2:2:246:GLU:HG2	2.21	0.40
4:4:20:TYR:CD2	4:4:21:GLY:N	2.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	1	284/302 (94%)	275 (97%)	8 (3%)	1 (0%)	39 42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	2	265/272 (97%)	252 (95%)	11 (4%)	2 (1%)	24	22
3	3	233/237 (98%)	224 (96%)	9 (4%)	0	100	100
4	4	66/68 (97%)	54 (82%)	6 (9%)	6 (9%)	1	0
All	All	848/879 (96%)	805 (95%)	34 (4%)	9 (1%)	17	14

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	4	15	ASN
4	4	23	SER
4	4	24	THR
4	4	19	ALA
4	4	21	GLY
2	2	48	ASN
1	1	145	THR
2	2	114	ALA
4	4	60	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	249/260 (96%)	245 (98%)	4 (2%)	70	82
2	2	227/232 (98%)	221 (97%)	6 (3%)	54	66
3	3	210/211 (100%)	205 (98%)	5 (2%)	57	69
4	4	57/57 (100%)	54 (95%)	3 (5%)	28	32
All	All	743/760 (98%)	725 (98%)	18 (2%)	57	69

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	1	147	ASN
1	1	177	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	1	224	LEU
1	1	286	TYR
2	2	66	LEU
2	2	74	GLU
2	2	139	THR
2	2	236	PRO
2	2	246	GLU
2	2	264	ARG
3	3	85	LEU
3	3	89	PRO
3	3	149	MET
3	3	208	MET
3	3	218	ASN
4	4	18	ARG
4	4	61	LEU
4	4	66	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	1	69	HIS
2	2	52	GLN
2	2	272	GLN
3	3	6	ASN
3	3	218	ASN
4	4	31	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PLM	1	2000	-	14,17,17	0.60	0	14,17,17	0.78	0
5	MYR	4	1	4	14,14,15	0.66	1 (7%)	12,13,15	1.21	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PLM	1	2000	-	-	0/13/15/15	0/0/0/0
5	MYR	4	1	4	-	0/11/12/13	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	4	1	MYR	C3-C2	2.11	1.60	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	4	1	MYR	C4-C3-C2	-2.93	102.43	113.86
5	4	1	MYR	C5-C4-C3	-2.10	103.69	114.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	1	2000	PLM	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.