



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2016 – 04:50 AM EDT

PDB ID : 5HZG
Title : The crystal structure of the strigolactone-induced AtD14-D3-ASK1 complex
Authors : Yao, R.F.; Ming, Z.H.; Yan, L.M.; Rao, Z.H.; Lou, Z.Y.; Xie, D.X.
Deposited on : 2016-02-02
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	<div> <div>0%</div> <div>66% 25% . .</div> </div>
1	E	267	<div> <div>3%</div> <div>70% 22% 5% .</div> </div>
2	B	740	<div> <div>2%</div> <div>50% 24% 8% . 17%</div> </div>
2	F	740	<div> <div>3%</div> <div>49% 24% 9% . 17%</div> </div>
3	C	169	<div> <div>27%</div> <div>56% 17% . . 23%</div> </div>
3	G	169	<div> <div>19%</div> <div>59% 18% . 22%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	6OM	A	301	-	X	X	X
4	6OM	E	301	-	X	X	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Strigolactone esterase D14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			2001	1285	341	369	6			
1	E	257	Total	C	N	O	S	0	0	0
			2001	1285	341	369	6			

- Molecule 2 is a protein called F-box/LRR-repeat MAX2 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	615	Total	C	N	O	S	0	0	0
			4791	3055	842	862	32			
2	F	614	Total	C	N	O	S	0	0	0
			4782	3048	840	862	32			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	GLY	-	expression tag	UNP Q5VMP0
B	-18	ALA	-	expression tag	UNP Q5VMP0
B	-17	MET	-	expression tag	UNP Q5VMP0
B	-16	GLY	-	expression tag	UNP Q5VMP0
B	-15	SER	-	expression tag	UNP Q5VMP0
B	-14	GLY	-	expression tag	UNP Q5VMP0
B	-13	ILE	-	expression tag	UNP Q5VMP0
B	-12	GLN	-	expression tag	UNP Q5VMP0
B	-11	ARG	-	expression tag	UNP Q5VMP0
B	-10	PRO	-	expression tag	UNP Q5VMP0
B	-9	THR	-	expression tag	UNP Q5VMP0
B	-8	SER	-	expression tag	UNP Q5VMP0
B	-7	THR	-	expression tag	UNP Q5VMP0
B	-6	SER	-	expression tag	UNP Q5VMP0
B	-5	SER	-	expression tag	UNP Q5VMP0
B	-4	LEU	-	expression tag	UNP Q5VMP0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	VAL	-	expression tag	UNP Q5VMP0
B	-2	ALA	-	expression tag	UNP Q5VMP0
B	-1	ALA	-	expression tag	UNP Q5VMP0
B	0	ALA	-	expression tag	UNP Q5VMP0
F	-19	GLY	-	expression tag	UNP Q5VMP0
F	-18	ALA	-	expression tag	UNP Q5VMP0
F	-17	MET	-	expression tag	UNP Q5VMP0
F	-16	GLY	-	expression tag	UNP Q5VMP0
F	-15	SER	-	expression tag	UNP Q5VMP0
F	-14	GLY	-	expression tag	UNP Q5VMP0
F	-13	ILE	-	expression tag	UNP Q5VMP0
F	-12	GLN	-	expression tag	UNP Q5VMP0
F	-11	ARG	-	expression tag	UNP Q5VMP0
F	-10	PRO	-	expression tag	UNP Q5VMP0
F	-9	THR	-	expression tag	UNP Q5VMP0
F	-8	SER	-	expression tag	UNP Q5VMP0
F	-7	THR	-	expression tag	UNP Q5VMP0
F	-6	SER	-	expression tag	UNP Q5VMP0
F	-5	SER	-	expression tag	UNP Q5VMP0
F	-4	LEU	-	expression tag	UNP Q5VMP0
F	-3	VAL	-	expression tag	UNP Q5VMP0
F	-2	ALA	-	expression tag	UNP Q5VMP0
F	-1	ALA	-	expression tag	UNP Q5VMP0
F	0	ALA	-	expression tag	UNP Q5VMP0

- Molecule 3 is a protein called SKP1-like protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	130	Total	C	N	O	S	0	0	0
			1024	645	168	207	4			
3	G	132	Total	C	N	O	S	0	0	0
			1039	655	170	210	4			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	-8	MET	-	expression tag	UNP Q39255
C	-7	ASP	-	expression tag	UNP Q39255
C	-6	TYR	-	expression tag	UNP Q39255
C	-5	LYS	-	expression tag	UNP Q39255
C	-4	ASP	-	expression tag	UNP Q39255
C	-3	ASP	-	expression tag	UNP Q39255

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	ASP	-	expression tag	UNP Q39255
C	-1	ASP	-	expression tag	UNP Q39255
C	0	LYS	-	expression tag	UNP Q39255
G	-8	MET	-	expression tag	UNP Q39255
G	-7	ASP	-	expression tag	UNP Q39255
G	-6	TYR	-	expression tag	UNP Q39255
G	-5	LYS	-	expression tag	UNP Q39255
G	-4	ASP	-	expression tag	UNP Q39255
G	-3	ASP	-	expression tag	UNP Q39255
G	-2	ASP	-	expression tag	UNP Q39255
G	-1	ASP	-	expression tag	UNP Q39255
G	0	LYS	-	expression tag	UNP Q39255

- Molecule 4 is (2Z)-2-methylbut-2-ene-1,4-diol (three-letter code: 6OM) (formula: C₅H₁₀O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 5 2	0	0
4	E	1	Total C O 7 5 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total O 3 3	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	5	Total 5	O 5	0	0
5	E	2	Total 2	O 2	0	0
5	F	10	Total 10	O 10	0	0
5	G	4	Total 4	O 4	0	0


i


- Molecule 1: Strigolactone esterase D14






• Molecule 2: F-box/LRR-repeat MAX2 homolog






• Molecule 3: SKP1-like protein 1A



• Molecule 3: SKP1-like protein 1A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.40Å 172.98Å 186.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.07 – 3.30 49.07 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.07-3.30) 99.5 (49.07-3.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.246 , 0.316 0.256 , 0.317	Depositor DCC
R_{free} test set	2669 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	81.2	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	15676	wwPDB-VP
Average B, all atoms (Å ²)	89.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 6OM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.56	0/2046	1.05	8/2785 (0.3%)
1	E	0.56	0/2046	1.17	22/2785 (0.8%)
2	B	0.64	2/4904 (0.0%)	1.27	53/6669 (0.8%)
2	F	0.63	3/4892 (0.1%)	1.31	72/6650 (1.1%)
3	C	0.52	0/1036	0.99	8/1397 (0.6%)
3	G	0.53	0/1052	0.85	6/1421 (0.4%)
All	All	0.60	5/15976 (0.0%)	1.20	169/21707 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	E	0	1
2	B	0	9
2	F	0	15
3	G	0	1
All	All	0	28

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	435	PRO	N-CD	13.99	1.67	1.47
2	B	186	PRO	N-CD	6.95	1.57	1.47
2	F	435	PRO	N-CD	6.95	1.57	1.47
2	F	568	PRO	N-CD	6.09	1.56	1.47
2	F	186	PRO	N-CD	5.12	1.55	1.47

The worst 5 of 169 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	597	ALA	CB-CA-C	-24.47	73.39	110.10
1	A	170	ALA	CB-CA-C	22.46	143.79	110.10
2	B	692	PRO	N-CA-CB	-18.29	81.35	103.30
2	F	656	LEU	CB-CA-C	-17.40	77.14	110.20
2	F	42	ALA	N-CA-C	16.11	154.50	111.00

There are no chirality outliers.

5 of 28 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	155	TRP	Peptide
1	A	96	HIS	Peptide
2	B	186	PRO	Peptide
2	B	241	THR	Peptide
2	B	400	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2001	0	1972	45	0
1	E	2001	0	1972	52	0
2	B	4791	0	4814	226	0
2	F	4782	0	4808	218	0
3	C	1024	0	1011	75	0
3	G	1039	0	1030	33	0
4	A	7	0	0	4	0
4	E	7	0	0	5	0
5	A	3	0	0	0	0
5	B	5	0	0	2	0
5	E	2	0	0	0	0
5	F	10	0	0	0	0
5	G	4	0	0	0	0
All	All	15676	0	15607	595	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 595 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:92:PRO:HB2	2:F:95:HIS:CD2	1.26	1.61
2:B:389:MET:CE	2:B:392:CYS:HB2	1.32	1.59
1:E:170:ALA:CB	1:E:172:VAL:HB	1.36	1.52
2:B:389:MET:HE1	2:B:392:CYS:CA	1.36	1.52
2:B:389:MET:CE	2:B:392:CYS:CB	1.88	1.52

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	253/267 (95%)	224 (88%)	28 (11%)	1 (0%)	39	76
1	E	253/267 (95%)	221 (87%)	32 (13%)	0	100	100
2	B	607/740 (82%)	517 (85%)	89 (15%)	1 (0%)	52	85
2	F	604/740 (82%)	516 (85%)	86 (14%)	2 (0%)	46	81
3	C	123/169 (73%)	101 (82%)	20 (16%)	2 (2%)	12	48
3	G	126/169 (75%)	108 (86%)	18 (14%)	0	100	100
All	All	1966/2352 (84%)	1687 (86%)	273 (14%)	6 (0%)	46	81

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	169	PRO
3	C	98	LEU
2	F	598	LEU
3	C	99	PHE
2	F	65	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/226 (95%)	183 (86%)	31 (14%)	4	18
1	E	214/226 (95%)	185 (86%)	29 (14%)	5	21
2	B	519/615 (84%)	420 (81%)	99 (19%)	2	8
2	F	519/615 (84%)	425 (82%)	94 (18%)	2	9
3	C	110/146 (75%)	103 (94%)	7 (6%)	22	60
3	G	112/146 (77%)	101 (90%)	11 (10%)	10	37
All	All	1688/1974 (86%)	1417 (84%)	271 (16%)	3	14

5 of 271 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	695	ASP
1	E	162	LEU
2	F	693	GLU
2	B	699	THR
1	E	6	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	533	GLN
1	E	196	ASN
3	G	46	ASN
2	B	683	GLN
2	F	95	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	6OM	A	301	1	5,6,6	3.67	5 (100%)	6,6,6	3.40	2 (33%)
4	6OM	E	301	-	5,6,6	3.66	5 (100%)	6,6,6	3.40	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	6OM	A	301	1	-	1/5/5/5	0/0/0/0
4	6OM	E	301	-	-	1/5/5/5	0/0/0/0

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	6OM	C3-C2	-3.94	1.45	1.50
4	E	301	6OM	C3-C2	-3.91	1.45	1.50
4	E	301	6OM	C4-C1	-3.43	1.41	1.50
4	A	301	6OM	C4-C1	-3.42	1.41	1.50
4	E	301	6OM	O2-C5	-2.82	1.31	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	301	6OM	C3-C2-C1	-7.62	115.22	126.61
4	E	301	6OM	C3-C2-C1	-7.60	115.25	126.61
4	E	301	6OM	O2-C5-C1	-2.60	106.12	111.76
4	A	301	6OM	O2-C5-C1	-2.58	106.15	111.76

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	301	6OM	O1-C3-C2-C1
4	A	301	6OM	O1-C3-C2-C1

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	301	6OM	4	0
4	E	301	6OM	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/267 (96%)	-0.12	2 (0%) 87 84	41, 65, 119, 148	0
1	E	257/267 (96%)	0.15	8 (3%) 52 46	53, 86, 155, 192	0
2	B	615/740 (83%)	0.06	12 (1%) 68 62	40, 78, 124, 159	0
2	F	614/740 (82%)	0.13	20 (3%) 50 43	41, 74, 126, 171	0
3	C	130/169 (76%)	1.64	46 (35%) 0 1	77, 152, 200, 232	0
3	G	132/169 (78%)	1.12	32 (24%) 1 1	77, 140, 182, 210	0
All	All	2005/2352 (85%)	0.24	120 (5%) 25 20	40, 80, 158, 232	0

The worst 5 of 120 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	73	VAL	10.5
3	G	73	VAL	7.9
3	C	90	PHE	7.2
3	C	76	ALA	6.5
3	C	77	ALA	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	6OM	A	301	7/7	0.75	0.36	3.44	50,51,55,55	0
4	6OM	E	301	7/7	0.80	0.42	2.23	50,51,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.