



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 19, 2016 – 04:10 PM EST

PDB ID : 5HZI  
Title : Crystal structure of photoinhibitable Intersectin1 containing C450M mutant LOV2 domain  
Authors : Tarnawski, M.; Dagliyan, O.; Chu, P.H.; Shirvanyants, D.; Dokholyan, N.V.; Hahn, K.M.; Schlichting, I.  
Deposited on : 2016-02-02  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	<div> <div>6%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>5%</div> </div> </div>
1	B	502	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>5%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7887 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Intersectin-1,NPH1-1,Intersectin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	476	Total	C	N	O	S	0	0	0
			3891	2483	675	713	20			
1	B	476	Total	C	N	O	S	0	0	0
			3891	2483	675	713	20			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1355	MET	CYS	engineered mutation	UNP O49003
A	1724	LEU	-	expression tag	UNP Q15811
A	1725	GLU	-	expression tag	UNP Q15811
A	1726	HIS	-	expression tag	UNP Q15811
A	1727	HIS	-	expression tag	UNP Q15811
A	1728	HIS	-	expression tag	UNP Q15811
A	1729	HIS	-	expression tag	UNP Q15811
A	1730	HIS	-	expression tag	UNP Q15811
A	1731	HIS	-	expression tag	UNP Q15811
B	1355	MET	CYS	engineered mutation	UNP O49003
B	1724	LEU	-	expression tag	UNP Q15811
B	1725	GLU	-	expression tag	UNP Q15811
B	1726	HIS	-	expression tag	UNP Q15811
B	1727	HIS	-	expression tag	UNP Q15811
B	1728	HIS	-	expression tag	UNP Q15811
B	1729	HIS	-	expression tag	UNP Q15811
B	1730	HIS	-	expression tag	UNP Q15811
B	1731	HIS	-	expression tag	UNP Q15811

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 17	N 4	O 9	P 1	0	0
2	B	1	Total 31	C 17	N 4	O 9	P 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	23	Total O 23 23	0	0
3	B	20	Total O 20 20	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Intersectin-1,NPH1-1,Intersectin-1



- Molecule 1: Intersectin-1,NPH1-1,Intersectin-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.06 Å   111.53 Å   88.33 Å 90.00°   90.63°   90.00°	Depositor
Resolution (Å)	47.15 – 2.60 47.15 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.15-2.60) 99.4 (47.15-2.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.38 (at 2.61 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.238 , 0.285 0.237 , 0.284	Depositor DCC
$R_{free}$ test set	1839 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	56.4	Xtriage
Anisotropy	0.676	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 30.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.128 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7887	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.21	0/3964	0.36	0/5340
1	B	0.21	0/3964	0.36	0/5340
All	All	0.21	0/7928	0.36	0/10680

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3891	0	3978	31	0
1	B	3891	0	3978	37	0
2	A	31	0	19	4	0
2	B	31	0	19	3	0
3	A	23	0	0	1	0
3	B	20	0	0	1	0
All	All	7887	0	7994	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1303:LYS:NZ	1:B:1453:MET:O	1.92	1.02
1:A:1303:LYS:NZ	1:A:1453:MET:O	1.94	1.00
1:B:1347:ARG:HH12	1:B:1510:LYS:HE3	1.49	0.78
1:A:1347:ARG:HH12	1:A:1510:LYS:HE3	1.50	0.77
1:B:1372:ARG:NH2	2:B:1801:FMN:O2P	2.23	0.70
1:A:1359:GLN:NE2	2:A:1801:FMN:O4'	2.25	0.69
1:B:1540:PRO:HG2	1:B:1543:HIS:HB2	1.76	0.68
1:B:1359:GLN:NE2	2:B:1801:FMN:O4'	2.28	0.65
1:A:1540:PRO:HG2	1:A:1543:HIS:HB2	1.77	0.64
1:B:1618:ASN:OD1	1:B:1695:ARG:NH2	2.31	0.64
1:A:1618:ASN:OD1	1:A:1695:ARG:NH2	2.30	0.64
1:A:1378:GLN:O	1:A:1405:ARG:NH2	2.34	0.61
1:A:1560:CYS:O	1:A:1564:ASN:ND2	2.33	0.59
1:B:1398:LEU:HB3	1:B:1419:LEU:HB3	1.84	0.59
1:A:1398:LEU:HB3	1:A:1419:LEU:HB3	1.84	0.59
1:B:1362:GLU:OE2	1:B:1394:LYS:NZ	2.32	0.59
1:B:1449:LYS:HA	1:B:1452:LYS:HE2	1.85	0.59
1:A:1449:LYS:HA	1:A:1452:LYS:HE2	1.84	0.59
1:B:1398:LEU:HD23	1:B:1419:LEU:HD23	1.85	0.58
1:A:1398:LEU:HD23	1:A:1419:LEU:HD23	1.86	0.58
1:B:1560:CYS:O	1:B:1564:ASN:ND2	2.33	0.57
1:A:1372:ARG:NH2	2:A:1801:FMN:O2P	2.39	0.55
1:A:1347:ARG:NH1	1:A:1510:LYS:HE3	2.21	0.55
1:B:1404:MET:HG2	1:B:1445:ASP:OD1	2.07	0.54
1:B:1670:VAL:HG22	1:B:1685:ILE:HG22	1.90	0.54
1:A:1238:ARG:NH2	1:A:1456:LYS:O	2.33	0.54
1:A:1628:ASP:OD2	1:A:1629:PHE:HD1	1.91	0.53
1:B:1347:ARG:NH1	1:B:1510:LYS:HE3	2.22	0.53
1:A:1497:LYS:NZ	1:A:1497:LYS:HB3	2.24	0.53
1:A:1670:VAL:HG22	1:A:1685:ILE:HG22	1.90	0.52
1:B:1669:LEU:HB2	1:B:1686:SER:HB3	1.92	0.52
1:B:1628:ASP:OD2	1:B:1629:PHE:HD1	1.93	0.52
1:B:1604:ARG:NH2	3:B:1902:HOH:O	2.43	0.51
1:A:1669:LEU:O	1:A:1686:SER:N	2.38	0.51
1:B:1497:LYS:NZ	1:B:1497:LYS:HB3	2.24	0.51
1:A:1682:ILE:HG13	1:A:1695:ARG:HG3	1.92	0.50
1:A:1695:ARG:NH2	3:A:1902:HOH:O	2.44	0.50
1:B:1619:LYS:HZ3	1:B:1659:TYR:HE1	1.60	0.49
1:A:1583:VAL:HG22	1:A:1656:TYR:HB2	1.94	0.49
1:A:1355:MET:HG2	2:A:1801:FMN:N5	2.27	0.48
1:A:1381:VAL:HG12	1:A:1401:LEU:HB3	1.95	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1362:GLU:OE2	1:A:1394:LYS:NZ	2.41	0.47
1:B:1381:VAL:HG12	1:B:1401:LEU:HB3	1.97	0.47
1:B:1513:PRO:O	1:B:1516:SER:OG	2.33	0.46
1:B:1669:LEU:O	1:B:1686:SER:N	2.36	0.46
1:B:1355:MET:HG2	2:B:1801:FMN:N5	2.30	0.46
1:A:1669:LEU:HB2	1:A:1686:SER:HB3	1.98	0.46
1:A:1264:GLN:HA	1:A:1282:PHE:CE2	2.52	0.45
1:B:1583:VAL:HG22	1:B:1656:TYR:HB2	1.98	0.45
1:B:1619:LYS:NZ	1:B:1659:TYR:HE1	2.15	0.44
1:A:1513:PRO:O	1:A:1516:SER:OG	2.35	0.44
1:A:1619:LYS:NZ	1:A:1659:TYR:HE1	2.16	0.44
1:B:1457:MET:HG2	1:B:1545:ASP:HB2	1.98	0.44
1:B:1672:LEU:HD11	1:B:1706:VAL:HG11	1.99	0.44
1:A:1258:LEU:O	1:A:1262:ILE:HG12	2.18	0.43
1:B:1682:ILE:HG13	1:B:1695:ARG:HG3	2.00	0.43
1:A:1404:MET:HG2	1:A:1445:ASP:OD1	2.18	0.43
1:B:1402:GLN:HA	1:B:1403:PRO:HD3	1.85	0.42
1:B:1258:LEU:O	1:B:1262:ILE:HG12	2.20	0.42
1:B:1533:LYS:HE2	1:B:1537:GLU:OE2	2.20	0.42
1:B:1264:GLN:HA	1:B:1282:PHE:CE2	2.55	0.42
1:B:1256:LEU:HB3	1:B:1289:ILE:HG12	2.02	0.42
1:A:1613:TYR:O	1:A:1695:ARG:N	2.51	0.41
1:B:1378:GLN:O	1:B:1405:ARG:NH2	2.54	0.41
1:B:1241:TYR:CD2	1:B:1534:ASN:HB3	2.56	0.41
1:B:1512:MET:HG2	1:B:1516:SER:OG	2.20	0.41
1:B:1604:ARG:NH1	1:B:1628:ASP:OD2	2.54	0.40
2:A:1801:FMN:H1'1	2:A:1801:FMN:H9	1.85	0.40
1:A:1448:ALA:O	1:A:1452:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	470/502 (94%)	450 (96%)	19 (4%)	1 (0%)	52	77
1	B	470/502 (94%)	449 (96%)	20 (4%)	1 (0%)	52	77
All	All	940/1004 (94%)	899 (96%)	39 (4%)	2 (0%)	52	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1283	VAL
1	B	1283	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	435/458 (95%)	435 (100%)	0	100	100
1	B	435/458 (95%)	435 (100%)	0	100	100
All	All	870/916 (95%)	870 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1359	GLN
1	B	1359	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMN	A	1801	-	32,33,33	1.18	3 (9%)	34,50,50	1.62	5 (14%)
2	FMN	B	1801	-	32,33,33	1.18	4 (12%)	34,50,50	1.63	5 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	1801	-	-	0/18/18/18	0/3/3/3
2	FMN	B	1801	-	-	0/18/18/18	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1801	FMN	C5A-N5	2.00	1.38	1.35
2	A	1801	FMN	C1'-N10	2.05	1.50	1.48
2	B	1801	FMN	C1'-N10	2.07	1.50	1.48
2	A	1801	FMN	C4-N3	2.96	1.38	1.33
2	B	1801	FMN	C4-N3	2.99	1.38	1.33
2	B	1801	FMN	C4A-N5	3.31	1.38	1.33
2	A	1801	FMN	C4A-N5	3.35	1.38	1.33

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1801	FMN	N3-C2-N1	-4.44	120.21	127.69
2	A	1801	FMN	N3-C2-N1	-4.38	120.32	127.69
2	A	1801	FMN	C4A-C4-N3	-2.84	119.81	123.52
2	B	1801	FMN	C4A-C4-N3	-2.79	119.88	123.52
2	A	1801	FMN	C4A-N5-C5A	2.68	119.88	116.72
2	B	1801	FMN	C4A-N5-C5A	2.73	119.94	116.72
2	B	1801	FMN	C5A-C9A-N10	3.24	120.00	117.58
2	A	1801	FMN	C5A-C9A-N10	3.31	120.06	117.58
2	A	1801	FMN	C4-N3-C2	5.89	120.07	115.16
2	B	1801	FMN	C4-N3-C2	5.92	120.09	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1801	FMN	4	0
2	B	1801	FMN	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	476/502 (94%)	0.22	28 (5%) 26 19	35, 68, 150, 238	0
1	B	476/502 (94%)	0.12	17 (3%) 46 38	33, 69, 140, 234	0
All	All	952/1004 (94%)	0.17	45 (4%) 35 28	33, 69, 144, 238	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1507	PRO	6.0
1	A	1616	LYS	5.9
1	B	1498	GLU	4.8
1	B	1721	LYS	4.7
1	A	1496	PHE	4.4
1	B	1507	PRO	4.2
1	A	1590	GLU	3.8
1	A	1485	LEU	3.7
1	A	1498	GLU	3.7
1	A	1587	GLY	3.6
1	A	1273	LEU	3.4
1	A	1588	LEU	3.4
1	B	1496	PHE	3.3
1	B	1684	HIS	3.2
1	A	1721	LYS	3.2
1	B	1671	LYS	3.2
1	A	1276	LYS	3.2
1	A	1486	ILE	3.1
1	A	1591	GLN	3.1
1	A	1494	PRO	2.9
1	A	1592	LEU	2.9
1	A	1687	HIS	2.8
1	B	1503	LEU	2.7
1	A	1489	LYS	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	1671	LYS	2.7
1	A	1431	ARG	2.7
1	B	1486	ILE	2.5
1	B	1692	TYR	2.4
1	A	1716	TYR	2.4
1	A	1651	LYS	2.3
1	B	1685	ILE	2.3
1	B	1281	ILE	2.3
1	B	1672	LEU	2.3
1	B	1688	ILE	2.3
1	A	1684	HIS	2.2
1	B	1691	VAL	2.2
1	A	1281	ILE	2.2
1	A	1647	VAL	2.2
1	A	1670	VAL	2.2
1	B	1654	LEU	2.2
1	A	1492	GLU	2.1
1	A	1272	LEU	2.1
1	B	1687	HIS	2.1
1	A	1669	LEU	2.1
1	B	1669	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	FMN	A	1801	31/31	0.92	0.17	-0.05	29,40,64,74	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FMN	B	1801	31/31	0.94	0.14	-0.59	41,56,71,77	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.