



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2016 – 10:52 PM EDT

PDB ID : 5I42
Title : Structure of HIV-1 Reverse Transcriptase in complex with a DNA aptamer, AZTTP, and CA(2+) ion
Authors : Das, K.; Arnold, E.
Deposited on : 2016-02-11
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	<div> <div>8%</div> <div> <div></div> <div>80%</div> <div>20%</div> </div> </div>
1	C	555	<div> <div>9%</div> <div> <div></div> <div>78%</div> <div>21%</div> <div>.</div> </div> </div>
2	B	444	<div> <div>2%</div> <div> <div></div> <div>74%</div> <div>18%</div> <div>7%</div> </div> </div>
2	D	444	<div> <div>4%</div> <div> <div></div> <div>74%</div> <div>18%</div> <div>7%</div> </div> </div>
3	E	38	<div> <div></div> <div> <div>50%</div> <div>39%</div> <div>8%</div> <div>.</div> </div> </div>
3	F	38	<div> <div></div> <div> <div>47%</div> <div>45%</div> <div>8%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 17320 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	554	Total	C	N	O	S	0	0	0
			4509	2919	750	833	7			
1	C	554	Total	C	N	O	S	0	0	0
			4509	2919	750	833	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	280	SER	CYS	engineered mutation	UNP P03366
C	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 2 is a protein called HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	412	Total	C	N	O	S	0	0	0
			3398	2212	562	618	6			
2	D	412	Total	C	N	O	S	0	0	0
			3400	2212	563	619	6			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	MET	-	initiating methionine	UNP P03366
B	-14	ALA	-	expression tag	UNP P03366
B	-13	HIS	-	expression tag	UNP P03366
B	-12	HIS	-	expression tag	UNP P03366
B	-11	HIS	-	expression tag	UNP P03366
B	-10	HIS	-	expression tag	UNP P03366
B	-9	HIS	-	expression tag	UNP P03366
B	-8	HIS	-	expression tag	UNP P03366
B	-7	ALA	-	expression tag	UNP P03366
B	-6	LEU	-	expression tag	UNP P03366

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLU	-	expression tag	UNP P03366
B	-4	VAL	-	expression tag	UNP P03366
B	-3	LEU	-	expression tag	UNP P03366
B	-2	PHE	-	expression tag	UNP P03366
B	-1	GLN	-	expression tag	UNP P03366
B	0	GLY	-	expression tag	UNP P03366
B	280	SER	CYS	engineered mutation	UNP P03366
D	-15	MET	-	initiating methionine	UNP P03366
D	-14	ALA	-	expression tag	UNP P03366
D	-13	HIS	-	expression tag	UNP P03366
D	-12	HIS	-	expression tag	UNP P03366
D	-11	HIS	-	expression tag	UNP P03366
D	-10	HIS	-	expression tag	UNP P03366
D	-9	HIS	-	expression tag	UNP P03366
D	-8	HIS	-	expression tag	UNP P03366
D	-7	ALA	-	expression tag	UNP P03366
D	-6	LEU	-	expression tag	UNP P03366
D	-5	GLU	-	expression tag	UNP P03366
D	-4	VAL	-	expression tag	UNP P03366
D	-3	LEU	-	expression tag	UNP P03366
D	-2	PHE	-	expression tag	UNP P03366
D	-1	GLN	-	expression tag	UNP P03366
D	0	GLY	-	expression tag	UNP P03366
D	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 3 is a DNA chain called DNA (38-MER).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	35	Total	C	N	O	P	0	0	0
			720	340	130	215	35			
3	F	35	Total	C	N	O	P	0	0	0
			720	340	130	215	35			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		

- Molecule 5 is 3'-AZIDO-3'-DEOXYTHYMIDINE-5'-TRIPHOSPHATE (three-letter code:

AZT) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
5	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT



• Molecule 1: HIV-1 REVERSE TRANSCRIPTASE P66 SUBUNIT



• Molecule 2: HIV-1 REVERSE TRANSCRIPTASE P51 SUBUNIT



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.15Å 132.19Å 139.66Å 90.00° 97.15° 90.00°	Depositor
Resolution (Å)	48.45 – 3.30 48.45 – 3.30	Depositor EDS
% Data completeness (in resolution range)	94.4 (48.45-3.30) 94.5 (48.45-3.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.87 (at 3.33Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, R_{free}	0.221 , 0.247 0.219 , 0.245	Depositor DCC
R_{free} test set	1888 reflections (4.08%)	DCC
Wilson B-factor (Å ²)	105.3	Xtriage
Anisotropy	0.356	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17320	wwPDB-VP
Average B, all atoms (Å ²)	117.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, CA, AZT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/4627	0.44	0/6286
1	C	0.25	0/4627	0.45	0/6286
2	B	0.26	0/3495	0.44	0/4748
2	D	0.27	0/3497	0.45	0/4751
3	E	0.75	2/759 (0.3%)	1.06	3/1170 (0.3%)
3	F	0.81	1/759 (0.1%)	1.05	2/1170 (0.2%)
All	All	0.33	3/17764 (0.0%)	0.54	5/24411 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	16	DC	C4'-O4'	15.77	1.60	1.45
3	E	16	DC	C4'-O4'	7.06	1.52	1.45
3	E	21	DC	C4'-O4'	5.63	1.50	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	F	16	DC	O4'-C4'-C3'	-12.41	98.56	106.00
3	F	16	DC	C5'-C4'-O4'	7.64	123.81	109.30
3	E	12	DG	O4'-C4'-C3'	-7.09	101.67	104.50
3	E	12	DG	C5'-C4'-O4'	6.54	121.72	109.30
3	E	21	DC	C5'-C4'-O4'	5.26	119.29	109.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4509	0	4570	78	0
1	C	4509	0	4570	84	0
2	B	3398	0	3431	60	0
2	D	3400	0	3433	56	0
3	E	720	0	397	13	0
3	F	720	0	397	16	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	31	0	12	4	0
5	C	31	0	12	3	0
All	All	17320	0	16822	280	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 280 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:ASN:HB2	1:A:289:LEU:CD2	2.07	0.85
1:C:34:LEU:HD21	1:C:132:ILE:HD12	1.59	0.83
1:C:448:ARG:HH21	3:F:19:DT:H5''	1.47	0.78
1:A:448:ARG:HH21	3:E:19:DT:H5''	1.48	0.78
1:A:110:ASP:OD2	1:A:111:VAL:N	2.16	0.78

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	552/555 (100%)	532 (96%)	20 (4%)	0	100	100
1	C	552/555 (100%)	532 (96%)	19 (3%)	1 (0%)	52	85
2	B	408/444 (92%)	395 (97%)	13 (3%)	0	100	100
2	D	408/444 (92%)	394 (97%)	14 (3%)	0	100	100
All	All	1920/1998 (96%)	1853 (96%)	66 (3%)	1 (0%)	56	89

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	55	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	495/495 (100%)	489 (99%)	6 (1%)	78	90
1	C	495/495 (100%)	489 (99%)	6 (1%)	78	90
2	B	373/403 (93%)	372 (100%)	1 (0%)	94	98
2	D	374/403 (93%)	371 (99%)	3 (1%)	86	93
All	All	1737/1796 (97%)	1721 (99%)	16 (1%)	84	92

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	42	GLU
1	C	85	GLN
1	C	391	LEU
2	B	11	LYS
2	D	11	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	255	ASN
2	B	208	HIS
1	C	373	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OMC	E	3	3	15,22,23	3.11	6 (40%)	20,31,34	1.44	1 (5%)
3	OMC	E	5	3	15,22,23	3.10	6 (40%)	20,31,34	1.66	1 (5%)
3	OMC	F	3	3	15,22,23	3.14	6 (40%)	20,31,34	1.54	2 (10%)
3	OMC	F	5	3	15,22,23	3.11	6 (40%)	20,31,34	1.48	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMC	E	3	3	-	0/5/27/28	0/2/2/2
3	OMC	E	5	3	-	0/5/27/28	0/2/2/2
3	OMC	F	3	3	-	0/5/27/28	0/2/2/2
3	OMC	F	5	3	-	0/5/27/28	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	3	OMC	C4-N4	2.33	1.41	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	3	OMC	C4-N4	2.35	1.41	1.35
3	F	5	OMC	C4-N4	2.36	1.41	1.35
3	E	5	OMC	C4-N4	2.39	1.41	1.35
3	E	3	OMC	C5-C4	2.54	1.47	1.41

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	3	OMC	N4-C4-N3	2.01	120.01	116.50
3	E	3	OMC	C6-C5-C4	5.29	119.51	117.44
3	F	5	OMC	C6-C5-C4	5.66	119.65	117.44
3	F	3	OMC	C6-C5-C4	5.82	119.72	117.44
3	E	5	OMC	C6-C5-C4	6.55	120.00	117.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	3	OMC	1	0
3	F	5	OMC	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	AZT	A	602	4	22,32,32	2.23	4 (18%)	27,49,49	2.17	6 (22%)
5	AZT	C	602	4	22,32,32	2.26	4 (18%)	27,49,49	2.27	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	AZT	A	602	4	-	0/21/37/37	0/2/2/2
5	AZT	C	602	4	-	0/21/37/37	0/2/2/2

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	602	AZT	C3'-N3'	-2.38	1.42	1.48
5	A	602	AZT	C3'-N3'	-2.18	1.42	1.48
5	A	602	AZT	C2-N3	2.38	1.43	1.38
5	C	602	AZT	C2-N3	2.43	1.43	1.38
5	A	602	AZT	C4-N3	6.38	1.44	1.33

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	602	AZT	C5-C4-N3	-7.46	119.09	125.35
5	A	602	AZT	C5-C4-N3	-6.98	119.49	125.35
5	C	602	AZT	O2A-PA-O1A	-2.26	100.78	112.56
5	A	602	AZT	O2A-PA-O1A	-2.18	101.23	112.56
5	C	602	AZT	O2B-PB-O1B	-2.12	101.52	112.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	602	AZT	4	0
5	C	602	AZT	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	554/555 (99%)	0.34	42 (7%)	17 14	56, 111, 172, 201	0
1	C	554/555 (99%)	0.40	49 (8%)	12 10	57, 127, 181, 210	0
2	B	412/444 (92%)	0.03	10 (2%)	62 55	59, 103, 142, 171	0
2	D	412/444 (92%)	0.17	19 (4%)	36 30	66, 118, 161, 179	0
3	E	33/38 (86%)	-0.54	0	100 100	93, 131, 145, 167	0
3	F	33/38 (86%)	-0.55	0	100 100	110, 143, 170, 178	0
All	All	1998/2074 (96%)	0.23	120 (6%)	25 20	56, 115, 172, 210	0

The worst 5 of 120 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	26	LEU	8.5
1	C	133	PRO	7.6
1	C	29	GLU	6.8
1	A	135	ILE	6.0
1	A	136	ASN	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	OMC	E	3	21/22	0.93	0.19	-	102,118,122,127	0
3	OMC	E	5	21/22	0.93	0.25	-	84,98,104,109	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	OMC	F	5	21/22	0.93	0.28	-	113,123,128,130	0
3	OMC	F	3	21/22	0.96	0.24	-	117,124,130,144	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	AZT	C	602	31/31	0.94	0.18	-0.46	134,150,161,167	0
5	AZT	A	602	31/31	0.92	0.18	-0.81	128,142,157,160	0
4	CA	A	601	1/1	0.94	0.09	-1.51	150,150,150,150	0
4	CA	C	601	1/1	0.94	0.11	-1.71	170,170,170,170	0

6.5 Other polymers [i](#)

There are no such residues in this entry.