



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:40 AM GMT

PDB ID : 2I6Q  
Title : Complement component C2a  
Authors : Milder, F.J.; Raaijmakers, H.C.A.; Vandeputte, D.A.A.; Schouten, A.;  
Huizinga, E.G.; Romijn, R.A.; Hemrika, W.; Roos, A.; Daha, M.R.; Gros,  
P.  
Deposited on : 2006-08-29  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	517	<div> <div>5%</div> <div>77%</div> <div>19%</div> <div>..</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement C2a fragment.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	503	Total	C	N	O	S	0	1	0
			3994	2513	706	748	27			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	216	HIS	-	EXPRESSION TAG	UNP P06681
A	217	HIS	-	EXPRESSION TAG	UNP P06681
A	218	HIS	-	EXPRESSION TAG	UNP P06681
A	219	HIS	-	EXPRESSION TAG	UNP P06681
A	220	HIS	-	EXPRESSION TAG	UNP P06681
A	221	HIS	-	EXPRESSION TAG	UNP P06681
A	222	GLY	-	CLONING ARTIFACT	UNP P06681
A	223	SER	-	CLONING ARTIFACT	UNP P06681

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	3	Total	C	N	O	0	0
			38	22	2	14		

- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mn	0	0
			1	1		

- Molecule 6 is MALONATE ION (three-letter code: MLI) (formula: C<sub>3</sub>H<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	3	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	133	Total	O	0	0
			133	133		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Complement C2a fragment



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.48 Å   77.19 Å   70.86 Å 90.00°   109.55°   90.00°	Depositor
Resolution (Å)	50.50 – 2.10 50.50 – 2.10	Depositor EDS
% Data completeness (in resolution range)	89.6 (50.50-2.10) 89.7 (50.50-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 2.10 Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.189 , 0.240 0.189 , 0.240	Depositor DCC
$R_{free}$ test set	1389 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.3	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 53.5	EDS
Estimated twinning fraction	0.029 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 27470 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4226	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG, MLI, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/4078	0.65	21/5511 (0.4%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	240	ASP	CB-CG-OD2	6.98	124.58	118.30
1	A	651	ASP	CB-CG-OD2	5.96	123.66	118.30
1	A	356	ASP	CB-CG-OD2	5.70	123.43	118.30
1	A	494	ASP	CB-CG-OD2	5.68	123.41	118.30
1	A	504	ASP	CB-CG-OD2	5.65	123.38	118.30
1	A	249	ASP	CB-CG-OD2	5.55	123.29	118.30
1	A	707	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	725	ASP	CB-CG-OD2	5.38	123.14	118.30
1	A	291	ASP	CB-CG-OD2	5.32	123.08	118.30
1	A	411	ASP	CB-CG-OD2	5.32	123.08	118.30
1	A	421	ASP	CB-CG-OD2	5.25	123.02	118.30
1	A	295	ASP	CB-CG-OD2	5.21	122.99	118.30
1	A	434	ASP	CB-CG-OD2	5.19	122.97	118.30
1	A	540	ASP	CB-CG-OD2	5.17	122.95	118.30
1	A	384	ASP	CB-CG-OD2	5.16	122.95	118.30
1	A	397	ASP	CB-CG-OD2	5.12	122.91	118.30
1	A	454	ASP	CB-CG-OD2	5.09	122.88	118.30
1	A	399	ASP	CB-CG-OD2	5.06	122.86	118.30
1	A	262[A]	ASP	CB-CG-OD2	5.02	122.82	118.30
1	A	262[B]	ASP	CB-CG-OD2	5.02	122.82	118.30
1	A	582	ASP	CB-CG-OD2	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3994	0	3942	57	1
2	A	38	0	34	0	0
3	A	39	0	34	0	0
4	A	14	0	13	0	0
5	A	1	0	0	0	0
6	A	7	0	2	0	0
7	A	133	0	0	1	0
All	All	4226	0	4025	57	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:428:VAL:HG12	1:A:432:MET:CE	1.80	1.12
1:A:428:VAL:HG12	1:A:432:MET:HE3	1.35	1.06
1:A:296:MET:SD	1:A:573:ARG:NH1	2.36	0.99
1:A:498:TRP:HB2	1:A:515:ILE:HD12	1.57	0.86
1:A:256:SER:HB3	1:A:425:LEU:HD22	1.56	0.85
1:A:428:VAL:CG1	1:A:432:MET:CE	2.60	0.78
1:A:428:VAL:CG1	1:A:432:MET:HE3	2.12	0.78
1:A:587:LEU:O	1:A:667:ARG:NH2	2.20	0.74
1:A:574:ARG:NH2	1:A:582:ASP:HB3	2.05	0.72
1:A:256:SER:HB3	1:A:425:LEU:CD2	2.20	0.71
1:A:574:ARG:HH22	1:A:582:ASP:HB3	1.56	0.70
1:A:508:GLN:CD	1:A:508:GLN:H	1.95	0.67
1:A:358:LYS:HE3	1:A:397:ASP:O	1.98	0.64
1:A:700:SER:O	1:A:701:LYS:HG3	1.99	0.62
1:A:467:LYS:HD2	1:A:497:LEU:HD13	1.82	0.61
1:A:256:SER:CB	1:A:425:LEU:HD22	2.32	0.58
1:A:467:LYS:HD3	1:A:491:ASP:OD2	2.04	0.58
1:A:291:ASP:CG	1:A:294:ARG:HH21	2.07	0.57
1:A:574:ARG:HH11	1:A:575:PRO:HD2	1.72	0.54
1:A:553:MET:SD	1:A:559:PRO:HD3	2.48	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:498:TRP:HB2	1:A:515:ILE:CD1	2.36	0.53
1:A:223:SER:HA	1:A:410:LYS:HE3	1.89	0.53
1:A:508:GLN:N	1:A:508:GLN:OE1	2.39	0.52
1:A:616:THR:O	1:A:620:GLU:HG2	2.10	0.50
1:A:243:GLN:HB2	1:A:315:THR:HB	1.93	0.50
1:A:448:MET:CE	1:A:669:PHE:HD1	2.25	0.49
1:A:705:PRO:O	1:A:706:ARG:HD3	2.13	0.49
1:A:574:ARG:NH1	1:A:578:SER:OG	2.47	0.48
1:A:379:ASN:HD21	1:A:381:LYS:HG2	1.78	0.47
1:A:563:PRO:O	1:A:564:CYS:HB2	2.16	0.46
1:A:376:LEU:O	1:A:377:ASN:HB2	2.16	0.46
1:A:509:TRP:HA	1:A:509:TRP:CE3	2.52	0.45
1:A:370:ASP:O	1:A:374:GLU:HG3	2.18	0.44
1:A:372:ILE:O	1:A:375:ILE:HG22	2.17	0.44
1:A:533:GLY:HA3	1:A:631:ASN:HB3	2.00	0.44
1:A:291:ASP:CB	1:A:294:ARG:HH21	2.31	0.43
1:A:466:PRO:HD2	1:A:470:GLU:O	2.18	0.43
1:A:400:TRP:CZ2	1:A:404:ASN:ND2	2.79	0.43
1:A:320:TYR:CD1	1:A:368:ALA:HB2	2.53	0.43
1:A:530:LYS:HA	1:A:534:ILE:O	2.19	0.43
1:A:519:VAL:HG12	1:A:718:TRP:HZ2	1.84	0.43
1:A:625:GLU:OE1	1:A:627:THR:HB	2.19	0.42
1:A:428:VAL:CG1	1:A:432:MET:HE2	2.47	0.42
1:A:400:TRP:CE2	1:A:404:ASN:ND2	2.87	0.42
1:A:495:HIS:HA	1:A:498:TRP:CD2	2.55	0.42
1:A:595:ALA:HB2	1:A:665:LEU:HD23	2.02	0.42
1:A:527:PHE:HB3	1:A:530:LYS:CE	2.50	0.42
1:A:437:LYS:HB2	7:A:6:HOH:O	2.20	0.42
1:A:460:TRP:HA	1:A:558:ARG:O	2.20	0.42
1:A:349:HIS:O	1:A:386:LEU:HA	2.19	0.42
1:A:595:ALA:HB2	1:A:665:LEU:CD2	2.51	0.41
1:A:702:VAL:HG13	1:A:703:PRO:HA	2.02	0.41
1:A:700:SER:C	1:A:701:LYS:HG3	2.40	0.41
1:A:638:VAL:HG23	1:A:639:VAL:HG23	2.02	0.41
1:A:574:ARG:NE	1:A:586:GLU:OE2	2.54	0.41
1:A:634:ASP:OD1	1:A:636:ARG:HG2	2.21	0.41
1:A:419:LEU:HD21	1:A:428:VAL:HG21	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:701:LYS:NZ	1:A:721:GLN:OE1[1_455]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	500/517 (97%)	480 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	445/455 (98%)	431 (97%)	14 (3%)	47	50

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	224	LYS
1	A	248	ASN
1	A	296	MET
1	A	300	ILE
1	A	331	ASN
1	A	336	LEU
1	A	423	LYS
1	A	440	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	526	VAL
1	A	549	GLN
1	A	574	ARG
1	A	629	PHE
1	A	667	ARG
1	A	706	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	331	ASN
1	A	377	ASN
1	A	379	ASN
1	A	549	GLN
1	A	585	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	1313	1,2	14,14,15	0.53	0	15,19,21	0.63	0
2	NAG	A	1314	2	14,14,15	0.61	0	15,19,21	1.19	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FUC	A	1315	2	10,10,11	0.34	0	14,14,16	1.10	1 (7%)
3	NAG	A	1447	1,3	14,14,15	0.50	0	15,19,21	0.87	0
3	NAG	A	1448	3	14,14,15	0.54	0	15,19,21	0.73	0
3	BMA	A	1449	3	11,11,12	0.59	0	14,15,17	1.17	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1313	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1314	2	-	0/6/23/26	0/1/1/1
2	FUC	A	1315	2	-	0/0/17/20	0/1/1/1
3	NAG	A	1447	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1448	3	-	0/6/23/26	0/1/1/1
3	BMA	A	1449	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1449	BMA	C1-C2-C3	2.01	111.92	109.54
3	A	1449	BMA	C3-C4-C5	2.05	113.78	110.20
2	A	1315	FUC	O5-C5-C6	2.28	109.90	106.13
2	A	1314	NAG	C4-C3-C2	3.50	116.67	111.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	MLI	A	1004	5	0,6,6	0.00	-	0,7,7	0.00	-
4	NAG	A	1601	1	14,14,15	0.52	0	15,19,21	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MLI	A	1004	5	-	0/0/4/4	0/0/0/0
4	NAG	A	1601	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	503/517 (97%)	0.36	25 (4%) 32 41	11, 20, 35, 47	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	490	ARG	5.5
1	A	632	LEU	4.3
1	A	491	ASP	4.2
1	A	414	ARG	4.1
1	A	379	ASN	3.5
1	A	223	SER	3.5
1	A	631	ASN	3.3
1	A	699	ARG	3.3
1	A	378	ILE	3.2
1	A	380	GLN	3.1
1	A	532	GLN	3.0
1	A	686	LEU	3.0
1	A	469	GLN	3.0
1	A	533	GLY	2.9
1	A	408	SER	2.9
1	A	700	SER	2.7
1	A	383	ASN	2.7
1	A	695	LYS	2.7
1	A	630	PRO	2.7
1	A	411	ASP	2.5
1	A	633	THR	2.5
1	A	377	ASN	2.3
1	A	381	LYS	2.2
1	A	694	ARG	2.2
1	A	531	ASN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	1447	14/15	0.97	0.13	0.71	6,10,20,21	0
2	FUC	A	1315	10/11	0.85	0.16	-0.37	49,50,52,52	0
2	NAG	A	1313	14/15	0.88	0.14	-0.40	38,44,49,51	0
3	BMA	A	1449	11/12	0.77	0.22	-	44,47,48,48	0
2	NAG	A	1314	14/15	0.76	0.29	-	55,57,59,60	0
3	NAG	A	1448	14/15	0.95	0.14	-	24,27,33,40	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	MLI	A	1004	7/7	0.97	0.12	-0.17	11,13,14,15	0
4	NAG	A	1601	14/15	0.92	0.14	-0.31	31,37,41,42	0
5	MN	A	1001	1/1	0.99	0.02	-6.62	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.