



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:42 AM GMT

PDB ID : 2IAG
Title : Crystal structure of human prostacyclin synthase
Authors : Chiang, C.-W.; Yeh, H.-C.; Wang, L.-H.; Chan, N.-L.
Deposited on : 2006-09-08
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div>..</div> </div> </div>
1	B	482	<div> <div>2%</div> <div> <div></div> <div>78%</div> <div>18%</div> <div>..</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostacyclin synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	471	Total	C	N	O	S	0	0	0
			3816	2454	670	677	15			
1	B	468	Total	C	N	O	S	0	0	0
			3795	2441	668	671	15			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	501	HIS	-	EXPRESSION TAG	UNP Q16647
A	502	HIS	-	EXPRESSION TAG	UNP Q16647
A	503	HIS	-	EXPRESSION TAG	UNP Q16647
A	504	HIS	-	EXPRESSION TAG	UNP Q16647
B	501	HIS	-	EXPRESSION TAG	UNP Q16647
B	502	HIS	-	EXPRESSION TAG	UNP Q16647
B	503	HIS	-	EXPRESSION TAG	UNP Q16647
B	504	HIS	-	EXPRESSION TAG	UNP Q16647

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	212	Total O 212 212	0	0
4	B	223	Total O 223 223	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Prostacyclin synthase



• Molecule 1: Prostacyclin synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.73Å 105.60Å 73.95Å 90.00° 92.01° 90.00°	Depositor
Resolution (Å)	20.15 – 2.15 20.15 – 2.15	Depositor EDS
% Data completeness (in resolution range)	91.0 (20.15-2.15) 90.9 (20.15-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.16 (at 2.15Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.201 , 0.263 0.201 , 0.259	Depositor DCC
R_{free} test set	2601 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.8	EDS
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 51303 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8133	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/3917	0.69	3/5311 (0.1%)
1	B	0.53	0/3897	0.66	2/5282 (0.0%)
All	All	0.53	0/7814	0.68	5/10593 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	480	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	A	144	LEU	CA-CB-CG	5.54	128.05	115.30
1	A	310	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	B	480	ARG	NE-CZ-NH2	-5.16	117.72	120.30
1	B	77	LEU	CA-CB-CG	5.09	127.01	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	186	LEU	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	186	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3816	0	3800	51	0
1	B	3795	0	3775	46	0
2	B	1	0	0	0	0
3	A	43	0	30	3	0
3	B	43	0	30	1	0
4	A	212	0	0	2	0
4	B	223	0	0	5	0
All	All	8133	0	7635	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:ARG:HG3	1:A:236:ARG:HH11	1.30	0.96
1:B:100:ALA:O	1:B:104:MET:HG3	1.71	0.90
1:A:140:MET:CE	1:A:175:ALA:HB1	2.03	0.88
1:B:284:THR:HG23	1:B:285:GLN:HE21	1.38	0.87
1:A:140:MET:HE3	1:A:175:ALA:HB1	1.57	0.86
1:B:423:ASP:CB	4:B:881:HOH:O	2.25	0.85
1:A:236:ARG:CG	1:A:236:ARG:HH11	1.89	0.85
1:B:423:ASP:HB2	4:B:881:HOH:O	1.79	0.83
1:B:316:ILE:HD11	1:B:337:SER:O	1.84	0.78
1:A:77:LEU:HB3	1:A:385:LEU:HD22	1.64	0.77
1:B:125:LYS:HD2	1:B:129:LEU:HD11	1.67	0.76
1:B:284:THR:HG23	1:B:285:GLN:NE2	2.01	0.75
1:A:140:MET:HE1	1:A:175:ALA:CB	2.16	0.74
1:A:365:LEU:HD11	1:A:377:LEU:HD12	1.69	0.73
1:A:330:LEU:HD11	1:A:334:VAL:HB	1.70	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:LEU:HD13	1:A:222:LEU:HD11	1.73	0.69
1:A:211:ASP:O	1:A:480:ARG:NH2	2.25	0.67
1:B:423:ASP:HB3	4:B:881:HOH:O	1.91	0.66
1:B:371:ASP:OD1	1:B:373:ARG:HD3	1.96	0.66
1:A:187:PRO:HD2	1:A:188:ARG:H	1.60	0.65
1:A:208:ARG:HH12	1:A:487:GLN:HE22	1.44	0.64
1:B:280:GLN:O	1:B:284:THR:HB	1.98	0.64
1:A:125:LYS:HE2	1:A:129:LEU:HD11	1.81	0.63
1:A:140:MET:HE1	1:A:175:ALA:HB3	1.80	0.62
1:A:140:MET:HE1	1:A:175:ALA:HB1	1.78	0.62
1:B:211:ASP:O	1:B:480:ARG:NH2	2.23	0.62
1:A:140:MET:CE	1:A:175:ALA:CB	2.74	0.61
1:A:187:PRO:CD	1:A:188:ARG:H	2.13	0.61
1:A:236:ARG:HG3	1:A:236:ARG:NH1	2.10	0.59
1:A:362:VAL:O	1:A:379:ARG:HD3	2.02	0.58
1:A:219:ARG:NH2	1:A:479:SER:O	2.35	0.58
1:B:216:LYS:HB3	1:B:222:LEU:CD2	2.34	0.58
1:A:306:LEU:O	1:A:310:ARG:HG3	2.04	0.57
1:B:458:VAL:HG11	1:B:464:LEU:HD22	1.88	0.56
1:B:125:LYS:HD2	1:B:129:LEU:CD1	2.36	0.56
1:B:219:ARG:HH12	1:B:479:SER:HB2	1.70	0.55
1:B:145:HIS:O	1:B:149:LEU:HB2	2.07	0.54
1:A:447:ALA:HB2	3:A:600:HEM:HMC2	1.89	0.54
1:B:500:PRO:O	1:B:502:HIS:N	2.41	0.54
1:B:284:THR:HG22	1:B:285:GLN:HG2	1.90	0.54
1:A:76:VAL:HG12	1:A:78:LEU:HD13	1.89	0.53
1:A:236:ARG:CG	1:A:236:ARG:NH1	2.60	0.53
1:A:145:HIS:NE2	1:A:149:LEU:HD12	2.24	0.52
1:A:330:LEU:CD1	1:A:334:VAL:HB	2.37	0.52
1:A:201:ALA:O	1:A:205:HIS:CD2	2.63	0.51
1:A:487:GLN:NE2	4:A:777:HOH:O	2.43	0.51
1:B:97:HIS:CD2	1:B:101:ILE:HD11	2.45	0.51
1:A:186:LEU:HB2	1:A:250:ALA:HB1	1.93	0.51
1:B:68:LEU:HD13	1:B:73:TYR:CE1	2.46	0.50
1:B:284:THR:CG2	1:B:285:GLN:HE21	2.15	0.50
1:A:145:HIS:CE1	1:A:149:LEU:HD12	2.46	0.49
1:B:84:ASP:OD2	1:B:428:LYS:HD2	2.11	0.49
1:A:330:LEU:HD22	1:A:331:PRO:HD2	1.94	0.49
1:B:89:GLU:HG3	1:B:94:LEU:HD12	1.94	0.49
1:A:83:TYR:CD1	1:A:385:LEU:HD13	2.48	0.49
1:B:96:PHE:HB2	1:B:439:ASN:ND2	2.27	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:LEU:HB3	1:A:215:PRO:HD3	1.94	0.49
1:B:77:LEU:HB3	1:B:385:LEU:HD22	1.95	0.48
1:B:91:ARG:O	1:B:362:VAL:HG21	2.14	0.48
1:A:201:ALA:O	1:A:205:HIS:HD2	1.97	0.48
1:B:206:THR:HA	1:B:209:GLN:HE21	1.79	0.47
1:B:357:ILE:HB	1:B:385:LEU:HB2	1.96	0.47
1:B:245:ARG:NH2	4:B:796:HOH:O	2.47	0.47
1:B:83:TYR:CD1	1:B:385:LEU:HD13	2.51	0.46
1:B:294:TRP:CG	1:B:352:THR:HG21	2.50	0.46
1:A:187:PRO:CD	1:A:188:ARG:N	2.76	0.46
1:B:219:ARG:NH1	1:B:479:SER:HB2	2.31	0.46
1:A:227:LYS:HB2	1:A:227:LYS:HE2	1.74	0.46
1:B:154:GLU:HG3	4:B:869:HOH:O	2.16	0.45
1:A:43:ALA:HA	1:A:69:VAL:HG23	1.99	0.45
3:B:600:HEM:HHC	3:B:600:HEM:HBB2	1.98	0.45
1:A:140:MET:HE2	1:A:144:LEU:HD22	1.98	0.44
1:A:499:ARG:HA	1:A:500:PRO:HD3	1.79	0.44
1:B:284:THR:CG2	1:B:285:GLN:HG2	2.47	0.44
1:B:148:LEU:HD22	1:B:171:PHE:CG	2.53	0.44
1:B:116:SER:HA	1:B:117:PRO:HD3	1.88	0.44
1:B:239:LYS:HD2	1:B:239:LYS:HA	1.73	0.42
1:B:316:ILE:CG2	1:B:333:LYS:HB2	2.50	0.42
1:A:187:PRO:HD2	1:A:188:ARG:N	2.33	0.42
1:B:457:LEU:O	1:B:461:HIS:HB2	2.18	0.42
1:B:97:HIS:CD2	1:B:117:PRO:HG3	2.54	0.42
1:A:23:ARG:HG3	1:A:32:LEU:HD21	2.00	0.42
1:B:186:LEU:HB2	1:B:250:ALA:HB1	2.01	0.42
1:A:74:VAL:HA	1:A:382:ARG:O	2.19	0.42
1:A:89:GLU:HA	1:A:90:PRO:HD3	1.86	0.42
1:B:134:GLN:O	1:B:138:GLU:HG3	2.19	0.42
1:A:238:TRP:CE2	1:A:275:ARG:HG2	2.54	0.42
1:A:501:HIS:HB3	1:B:444:ARG:HD3	2.02	0.41
1:B:141:TYR:HB2	1:B:453:GLN:NE2	2.35	0.41
1:A:293:PHE:CE2	1:A:488:PRO:HG3	2.55	0.41
1:A:125:LYS:HE3	4:A:657:HOH:O	2.21	0.41
1:A:284:THR:HB	3:A:600:HEM:CBC	2.51	0.41
1:A:389:LEU:HD12	1:A:389:LEU:HA	1.89	0.40
1:B:398:TYR:CD2	1:B:408:ARG:HD3	2.56	0.40
1:B:313:LEU:HD22	1:B:456:PHE:HD1	1.85	0.40
1:A:468:ASN:ND2	1:A:470:ASP:H	2.19	0.40
3:A:600:HEM:HBB2	3:A:600:HEM:HMB2	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:TRP:CG	1:A:352:THR:HG21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	467/482 (97%)	458 (98%)	9 (2%)	0	100	100
1	B	464/482 (96%)	447 (96%)	13 (3%)	4 (1%)	21	13
All	All	931/964 (97%)	905 (97%)	22 (2%)	4 (0%)	39	34

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	501	HIS
1	B	188	ARG
1	B	332	GLN
1	B	187	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	408/418 (98%)	381 (93%)	27 (7%)	21	14

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	405/418 (97%)	375 (93%)	30 (7%)	17	11
All	All	813/836 (97%)	756 (93%)	57 (7%)	19	12

All (57) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	VAL
1	A	77	LEU
1	A	78	LEU
1	A	110	VAL
1	A	112	LEU
1	A	128	LEU
1	A	142	THR
1	A	144	LEU
1	A	148	LEU
1	A	169	TYR
1	A	186	LEU
1	A	188	ARG
1	A	213	LEU
1	A	217	LEU
1	A	227	LYS
1	A	236	ARG
1	A	248	ARG
1	A	260	LEU
1	A	300	LEU
1	A	304	GLU
1	A	306	LEU
1	A	317	LEU
1	A	335	LEU
1	A	348	SER
1	A	363	VAL
1	A	379	ARG
1	A	468	ASN
1	B	44	LEU
1	B	69	VAL
1	B	72	ARG
1	B	77	LEU
1	B	78	LEU
1	B	105	GLU
1	B	110	VAL
1	B	112	LEU
1	B	128	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	142	THR
1	B	148	LEU
1	B	149	LEU
1	B	169	TYR
1	B	186	LEU
1	B	188	ARG
1	B	213	LEU
1	B	230	MET
1	B	240	LEU
1	B	248	ARG
1	B	261	LEU
1	B	284	THR
1	B	296	LEU
1	B	300	LEU
1	B	306	LEU
1	B	314	GLU
1	B	317	LEU
1	B	318	TRP
1	B	389	LEU
1	B	442	LEU
1	B	459	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	GLN
1	A	160	HIS
1	A	205	HIS
1	A	449	ASN
1	A	453	GLN
1	A	468	ASN
1	A	487	GLN
1	A	490	HIS
1	B	195	GLN
1	B	209	GLN
1	B	285	GLN
1	B	332	GLN
1	B	449	ASN
1	B	468	ASN
1	B	490	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HEM	A	600	1	30,50,50	2.20	10 (33%)	24,82,82	2.40	7 (29%)
3	HEM	B	600	1	30,50,50	2.26	9 (30%)	24,82,82	2.29	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	600	1	-	0/10/54/54	0/0/8/8
3	HEM	B	600	1	-	0/10/54/54	0/0/8/8

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	600	HEM	C3B-C4B	-7.45	1.45	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	600	HEM	C3B-C4B	-6.21	1.46	1.51
3	B	600	HEM	C3D-C4D	-5.24	1.44	1.51
3	A	600	HEM	C3D-C4D	-3.93	1.46	1.51
3	B	600	HEM	C2C-C1C	-3.58	1.45	1.52
3	A	600	HEM	C2C-C1C	-2.97	1.46	1.52
3	A	600	HEM	C2D-C1D	-2.40	1.44	1.51
3	B	600	HEM	C2B-C1B	-2.27	1.44	1.51
3	B	600	HEM	C3B-CAB	2.07	1.55	1.51
3	B	600	HEM	CAA-C2A	2.21	1.55	1.52
3	A	600	HEM	C3C-CAC	2.24	1.55	1.51
3	A	600	HEM	C1C-NC	2.27	1.38	1.36
3	A	600	HEM	C3B-CAB	2.38	1.55	1.51
3	A	600	HEM	C4C-NC	2.40	1.39	1.36
3	B	600	HEM	C4C-NC	2.41	1.39	1.36
3	B	600	HEM	C3C-CAC	2.87	1.56	1.51
3	A	600	HEM	FE-NC	3.03	2.07	1.95
3	A	600	HEM	FE-ND	3.18	2.14	1.97
3	B	600	HEM	FE-NC	4.74	2.14	1.95

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	600	HEM	CMA-C3A-C4A	-2.52	124.19	128.36
3	B	600	HEM	CMA-C3A-C4A	-2.44	124.32	128.36
3	A	600	HEM	CAA-CBA-CGA	-2.18	108.74	112.75
3	B	600	HEM	CBA-CAA-C2A	-2.17	108.64	112.53
3	B	600	HEM	C2D-C3D-C4D	2.08	105.02	101.50
3	B	600	HEM	CMD-C2D-C3D	2.91	127.23	114.35
3	B	600	HEM	CMB-C2B-C3B	3.40	125.01	116.53
3	B	600	HEM	CMC-C2C-C3C	3.42	125.06	116.53
3	A	600	HEM	CMD-C2D-C3D	3.45	129.62	114.35
3	A	600	HEM	CAD-C3D-C4D	3.69	125.49	112.47
3	B	600	HEM	C3B-C4B-CHC	3.77	128.47	123.16
3	B	600	HEM	CAD-C3D-C4D	3.87	126.12	112.47
3	A	600	HEM	CMC-C2C-C3C	4.21	127.04	116.53
3	A	600	HEM	CMB-C2B-C3B	4.97	128.93	116.53
3	B	600	HEM	CAD-C3D-C2D	5.44	128.85	113.22
3	A	600	HEM	CAD-C3D-C2D	5.88	130.13	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	HEM	3	0
3	B	600	HEM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	471/482 (97%)	0.07	13 (2%) 56 66	10, 26, 45, 53	0
1	B	468/482 (97%)	-0.04	9 (1%) 70 78	13, 25, 38, 55	0
All	All	939/964 (97%)	0.01	22 (2%) 64 72	10, 25, 42, 55	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	318	TRP	5.3
1	A	328	THR	4.5
1	A	38	PRO	4.0
1	A	39	TRP	3.7
1	A	363	VAL	3.7
1	A	370	ALA	3.5
1	A	114	HIS	3.4
1	A	187	PRO	3.4
1	B	102	PHE	3.3
1	B	186	LEU	3.3
1	B	114	HIS	2.9
1	B	317	LEU	2.6
1	A	329	THR	2.6
1	A	102	PHE	2.6
1	B	333	LYS	2.5
1	A	85	ALA	2.4
1	B	99	TYR	2.2
1	B	332	GLN	2.1
1	A	92	THR	2.1
1	A	30	PRO	2.1
1	B	490	HIS	2.1
1	A	372	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	HEM	B	600	43/43	0.95	0.13	1.13	19,23,40,44	0
3	HEM	A	600	43/43	0.96	0.12	0.35	20,23,38,40	0
2	NA	B	701	1/1	0.99	0.08	-4.15	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.