



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 28, 2016 – 02:05 AM EST

PDB ID : 5IE5
Title : Crystal structure of a lactonase double mutant in complex with substrate a
Authors : Zheng, Y.Y.; Xu, Z.X.; Liu, W.D.; Chen, C.C.; Guo, R.T.
Deposited on : 2016-02-25
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028442
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	<div> <div>8%</div> <div>84%</div> <div>13%</div> <div>.</div> </div>
1	B	264	<div> <div>5%</div> <div>83%</div> <div>15%</div> <div>.</div> </div>
1	C	264	<div> <div>46%</div> <div>40%</div> <div>18%</div> <div>5%</div> <div>38%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	36J	B	301	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Zearalenone hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2023	1283	343	386	11			
1	B	264	Total	C	N	O	S	0	0	0
			2023	1283	343	386	11			
1	C	164	Total	C	N	O	S	0	0	0
			1255	806	207	235	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	SER	engineered mutation	UNP Q8NKB0
A	153	HIS	VAL	engineered mutation	UNP Q8NKB0
B	102	ALA	SER	engineered mutation	UNP Q8NKB0
B	153	HIS	VAL	engineered mutation	UNP Q8NKB0
C	102	ALA	SER	engineered mutation	UNP Q8NKB0
C	153	HIS	VAL	engineered mutation	UNP Q8NKB0

- Molecule 2 is (3S,7R,11E)-7,14,16-trihydroxy-3-methyl-3,4,5,6,7,8,9,10-octahydro-1H-2-benzoxacyclotetradecin-1-one (three-letter code: 36J) (formula: C₁₈H₂₄O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			23	18	5		
2	B	1	Total	C	O	0	0
			23	18	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	116	Total	O	0	0
			116	116		
3	B	114	Total	O	0	0
			114	114		
3	C	71	Total	O	0	0
			71	71		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Zearalenone hydrolase




• Molecule 1: Zearalenone hydrolase



• Molecule 1: Zearalenone hydrolase





4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	86.16 Å 86.16 Å 471.86 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 2.39 24.84 – 2.38	Depositor EDS
% Data completeness (in resolution range)	99.5 (25.00-2.39) 99.6 (24.84-2.38)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.62 (at 2.39 Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.214 , 0.270 0.264 , 0.305	Depositor DCC
R_{free} test set	2045 reflections (4.82%)	DCC
Wilson B-factor (Å ²)	40.8	Xtriage
Anisotropy	0.646	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5648	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 36J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/2075	0.77	2/2834 (0.1%)
1	B	0.56	0/2075	0.78	2/2834 (0.1%)
1	C	0.45	0/1279	0.63	0/1734
All	All	0.52	0/5429	0.75	4/7402 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	5
All	All	0	8

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	120	ARG	NE-CZ-NH2	-7.60	116.50	120.30
1	A	2	ARG	NE-CZ-NH1	-7.19	116.70	120.30
1	A	120	ARG	NE-CZ-NH2	-6.16	117.22	120.30
1	B	2	ARG	NE-CZ-NH1	-6.08	117.26	120.30

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	120	ARG	Sidechain
1	A	197	VAL	Mainchain
1	B	120	ARG	Sidechain
1	C	214	ALA	Peptide
1	C	223	ASP	Peptide
1	C	238	LEU	Peptide
1	C	250	ASP	Peptide
1	C	91	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2023	0	1989	30	0
1	B	2023	0	1989	17	0
1	C	1255	0	1222	29	0
2	A	23	0	22	7	0
2	B	23	0	22	5	0
3	A	116	0	0	1	1
3	B	114	0	0	0	1
3	C	71	0	0	1	0
All	All	5648	0	5244	82	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:MET:CE	1:A:245:TYR:HE2	1.75	1.00
1:C:88:LEU:O	1:C:92:ASP:O	1.88	0.92
1:C:209:ASP:OD1	1:C:234:ASN:ND2	2.13	0.82
1:A:67:ALA:HB1	1:A:71:THR:HG21	1.62	0.80
1:A:38:MET:CE	1:A:245:TYR:CE2	2.65	0.76
1:A:38:MET:HE2	1:A:245:TYR:HE2	1.50	0.76
1:B:76:THR:HG22	1:B:79:LYS:H	1.51	0.76
1:B:109:VAL:HG22	1:B:196:PRO:HG2	1.70	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:HIS:CD2	2:B:301:36J:H5	2.27	0.70
1:C:221:PHE:O	1:C:224:ASN:HB2	1.92	0.69
1:A:158:VAL:HG21	2:A:300:36J:H2	1.81	0.61
1:A:38:MET:HE3	1:A:245:TYR:CE2	2.36	0.60
1:A:38:MET:HE3	1:A:245:TYR:HE2	1.62	0.60
1:C:227:THR:O	1:C:231:ALA:HB3	2.03	0.59
2:A:300:36J:OAD	2:A:300:36J:OAB	2.16	0.57
1:A:129:THR:HG21	1:A:223:ASP:OD1	2.05	0.56
2:B:301:36J:OAP	2:B:301:36J:H15	2.06	0.56
1:A:102:ALA:HB1	2:A:300:36J:CAQ	2.36	0.56
1:A:81:ALA:HB1	1:A:111:LEU:HD13	1.88	0.55
1:A:154:MET:HE1	2:A:300:36J:H1	1.87	0.54
2:B:301:36J:OAD	2:B:301:36J:OAB	2.21	0.54
1:C:16:TYR:O	1:C:56:PHE:HA	2.08	0.54
1:C:26:VAL:O	1:C:53:VAL:C	2.46	0.54
1:C:97:THR:OG1	1:C:121:ASN:ND2	2.41	0.54
1:A:129:THR:HG23	3:A:476:HOH:O	2.09	0.53
1:C:9:THR:OG1	1:C:11:ASN:ND2	2.42	0.52
1:C:228:ALA:HB1	1:C:233:VAL:HB	1.92	0.51
1:C:216:THR:C	3:C:308:HOH:O	2.48	0.51
1:C:92:ASP:O	1:C:93:ILE:HB	2.11	0.51
1:A:59:PRO:CB	1:A:71:THR:HG23	2.40	0.51
1:A:105:ALA:HB1	1:A:124:CYS:HB2	1.93	0.51
1:A:11:ASN:H	1:A:11:ASN:ND2	2.09	0.51
1:A:197:VAL:HG22	1:A:203:LEU:HD21	1.93	0.50
1:A:38:MET:HE2	1:A:245:TYR:CE2	2.41	0.50
1:C:120:ARG:HG2	1:C:121:ASN:HD22	1.76	0.50
1:A:146:ILE:HG23	1:A:150:LEU:HD22	1.94	0.50
1:B:102:ALA:HB1	2:B:301:36J:CAQ	2.42	0.50
1:A:59:PRO:HB2	1:A:71:THR:HG23	1.92	0.49
1:C:253:ALA:O	1:C:257:VAL:HG13	2.12	0.49
1:B:105:ALA:HB1	1:B:124:CYS:HB2	1.95	0.49
1:C:96:ALA:O	1:C:119:ILE:HA	2.13	0.49
1:A:242:HIS:CD2	2:A:300:36J:H3	2.48	0.48
1:A:31:ASP:OD2	1:A:38:MET:HE1	2.14	0.48
1:A:146:ILE:CG2	1:A:150:LEU:HD22	2.43	0.48
1:B:187:TYR:CD2	1:B:191:ILE:HD13	2.49	0.48
1:A:134:HIS:CE1	1:A:135:LEU:HD13	2.50	0.47
1:C:39:PHE:O	1:C:43:VAL:N	2.44	0.47
1:C:7:ILE:O	1:C:7:ILE:HG23	2.14	0.46
1:C:97:THR:CG2	1:C:123:MET:CE	2.94	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:MET:SD	1:A:182:VAL:HG22	2.56	0.45
2:A:300:36J:H15	2:A:300:36J:OAP	2.16	0.45
1:C:42:SER:O	1:C:46:ILE:HB	2.17	0.45
1:C:91:LEU:HB2	1:C:93:ILE:HD12	1.98	0.45
1:B:76:THR:HG23	1:B:78:GLN:OE1	2.16	0.45
1:C:75:VAL:HG12	1:C:191:ILE:HD11	1.98	0.45
1:B:153:HIS:ND1	1:B:157:ASP:OD2	2.47	0.44
1:A:68:PRO:O	1:A:71:THR:HB	2.18	0.44
1:B:154:MET:O	1:B:159:SER:HB3	2.17	0.44
1:C:197:VAL:HG22	1:C:197:VAL:O	2.17	0.44
1:B:102:ALA:HB2	1:B:242:HIS:CE1	2.53	0.44
1:C:17:TYR:HA	1:C:56:PHE:HA	1.99	0.44
1:C:252:PHE:C	1:C:252:PHE:CD1	2.91	0.44
1:A:102:ALA:HB2	1:A:242:HIS:NE2	2.33	0.43
1:A:153:HIS:ND1	1:A:157:ASP:OD2	2.42	0.43
1:B:102:ALA:HB2	1:B:242:HIS:NE2	2.33	0.43
1:C:7:ILE:O	1:C:7:ILE:CG2	2.66	0.43
1:A:134:HIS:ND1	1:A:135:LEU:HD13	2.33	0.43
1:C:104:GLY:O	1:C:108:VAL:HG13	2.19	0.43
1:C:121:ASN:HA	1:C:207:PRO:HB2	2.00	0.42
1:B:144:GLU:OE2	1:B:177:HIS:CE1	2.72	0.42
1:C:252:PHE:O	1:C:256:VAL:HG23	2.19	0.42
1:B:80:LEU:O	1:B:84:VAL:HG13	2.19	0.42
1:B:75:VAL:HG21	1:B:187:TYR:CE1	2.55	0.42
1:B:158:VAL:O	1:B:158:VAL:HG22	2.19	0.41
2:B:301:36J:CAG	2:B:301:36J:OAP	2.67	0.41
2:A:300:36J:CAG	2:A:300:36J:OAP	2.68	0.41
1:C:25:ASP:O	1:C:120:ARG:NH2	2.53	0.41
1:C:108:VAL:HA	1:C:111:LEU:HB2	2.03	0.41
1:B:6:THR:HA	1:B:15:TRP:O	2.21	0.41
1:A:81:ALA:CB	1:A:111:LEU:HD13	2.51	0.40
1:A:61:MET:O	1:A:62:SER:C	2.57	0.40
1:B:3:THR:O	1:B:18:GLU:HA	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:491:HOH:O	3:B:482:HOH:O[6_555]	2.09	0.11

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/264 (99%)	251 (96%)	9 (3%)	2 (1%)	24	35
1	B	262/264 (99%)	253 (97%)	7 (3%)	2 (1%)	24	35
1	C	142/264 (54%)	131 (92%)	8 (6%)	3 (2%)	9	10
All	All	666/792 (84%)	635 (95%)	24 (4%)	7 (1%)	17	25

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	92	ASP
1	C	37	GLN
1	C	93	ILE
1	B	187	TYR
1	A	161	GLY
1	A	187	TYR
1	B	161	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/220 (100%)	202 (92%)	18 (8%)	14	21
1	B	220/220 (100%)	202 (92%)	18 (8%)	14	21
1	C	134/220 (61%)	114 (85%)	20 (15%)	4	4
All	All	574/660 (87%)	518 (90%)	56 (10%)	10	14

All (56) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	11	ASN
1	A	56	PHE
1	A	71	THR
1	A	74	GLU
1	A	111	LEU
1	A	129	THR
1	A	132	LEU
1	A	135	LEU
1	A	141	LEU
1	A	150	LEU
1	A	162	SER
1	A	191	ILE
1	A	197	VAL
1	A	200	LEU
1	A	203	LEU
1	A	235	ILE
1	A	238	LEU
1	B	1	MET
1	B	41	SER
1	B	56	PHE
1	B	74	GLU
1	B	76	THR
1	B	131	LEU
1	B	132	LEU
1	B	141	LEU
1	B	142	GLU
1	B	162	SER
1	B	172	VAL
1	B	176	LEU
1	B	189	ARG
1	B	191	ILE
1	B	200	LEU
1	B	203	LEU
1	B	235	ILE
1	B	251	VAL
1	C	11	ASN
1	C	36	CYS
1	C	37	GLN
1	C	41	SER
1	C	42	SER
1	C	57	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	86	SER
1	C	87	VAL
1	C	88	LEU
1	C	106	SER
1	C	111	LEU
1	C	113	LEU
1	C	191	ILE
1	C	203	LEU
1	C	206	LYS
1	C	216	THR
1	C	238	LEU
1	C	252	PHE
1	C	254	LYS
1	C	264	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	121	ASN
1	B	121	ASN
1	B	137	ASN
1	B	177	HIS
1	B	261	GLN
1	C	11	ASN
1	C	49	GLN
1	C	121	ASN
1	C	224	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	36J	A	300	-	24,24,24	1.54	3 (12%)	32,32,32	1.50	7 (21%)
2	36J	B	301	-	24,24,24	1.58	3 (12%)	32,32,32	1.32	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	36J	A	300	-	-	0/22/22/22	0/1/2/2
2	36J	B	301	-	-	0/22/22/22	0/1/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	36J	CAT-CAG	-5.04	1.39	1.47
2	A	300	36J	CAT-CAG	-4.96	1.39	1.47
2	B	301	36J	CAU-CAQ	-4.09	1.40	1.50
2	A	300	36J	CAU-CAQ	-4.06	1.40	1.50
2	A	300	36J	CAG-CAF	2.53	1.40	1.31
2	B	301	36J	CAG-CAF	2.58	1.40	1.31

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	36J	CAN-CAK-CAJ	-3.22	109.41	114.61
2	A	300	36J	OAD-CAS-CAU	-3.04	115.35	121.18
2	A	300	36J	CAN-CAK-CAJ	-2.94	109.86	114.61
2	B	301	36J	OAD-CAS-CAU	-2.77	115.87	121.18
2	A	300	36J	CAK-CAN-CAW	-2.35	106.76	114.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	36J	CAT-CAG-CAF	-2.34	119.94	125.35
2	B	301	36J	CAJ-CAF-CAG	-2.21	120.88	125.20
2	B	301	36J	CAA-CAV-CAM	-2.19	108.22	113.98
2	A	300	36J	CAS-CAH-CAR	-2.17	117.57	119.64
2	B	301	36J	CAV-OAP-CAQ	2.06	121.72	117.68
2	B	301	36J	OAP-CAV-CAA	2.11	112.65	107.89
2	A	300	36J	CAV-OAP-CAQ	2.92	123.42	117.68
2	A	300	36J	OAP-CAV-CAA	3.27	115.26	107.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	36J	7	0
2	B	301	36J	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	264/264 (100%)	0.33	22 (8%)	14 14	16, 25, 47, 62	0
1	B	264/264 (100%)	0.36	14 (5%)	30 30	18, 26, 44, 75	0
1	C	164/264 (62%)	3.50	121 (73%)	0 0	30, 45, 59, 63	0
All	All	692/792 (87%)	1.10	157 (22%)	1 1	16, 28, 55, 75	0

All (157) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	36	CYS	9.2
1	C	46	ILE	8.7
1	C	185	ARG	8.7
1	C	241	MET	8.4
1	C	24	PRO	7.9
1	C	184	ALA	7.8
1	C	18	GLU	7.8
1	C	87	VAL	7.7
1	C	222	PHE	7.6
1	C	47	ALA	7.4
1	C	72	TYR	7.3
1	C	189	ARG	7.0
1	C	28	LEU	6.9
1	C	191	ILE	6.9
1	C	16	TYR	6.9
1	C	8	SER	6.8
1	C	110	ALA	6.5
1	C	12	GLY	6.4
1	C	14	THR	6.3
1	C	193	PRO	6.2
1	C	39	PHE	6.2
1	C	38	MET	6.1
1	C	215	ALA	6.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	17	TYR	5.8
1	C	188	PRO	5.7
1	C	202	ALA	5.6
1	C	111	LEU	5.5
1	C	13	ILE	5.4
1	C	40	ASP	5.4
1	C	80	LEU	5.4
1	C	19	GLN	5.3
1	C	21	GLY	5.2
1	C	223	ASP	5.1
1	C	240	GLY	5.1
1	C	108	VAL	5.0
1	C	190	THR	5.0
1	C	90	ALA	4.9
1	C	73	THR	4.9
1	C	259	THR	4.9
1	A	48	ALA	4.9
1	C	114	GLY	4.8
1	C	102	ALA	4.8
1	C	192	PRO	4.7
1	C	22	THR	4.6
1	C	186	GLY	4.5
1	C	197	VAL	4.5
1	C	116	PRO	4.4
1	C	49	GLN	4.4
1	C	187	TYR	4.4
1	C	115	TYR	4.4
1	C	242	HIS	4.4
1	C	42	SER	4.3
1	C	252	PHE	4.3
1	C	45	GLN	4.2
1	C	44	SER	4.2
1	C	239	PRO	4.1
1	C	77	ALA	4.1
1	C	183	TRP	4.1
1	C	200	LEU	4.1
1	C	194	SER	4.1
1	C	41	SER	4.0
1	C	75	VAL	4.0
1	C	51	PHE	3.9
1	C	205	GLY	3.9
1	C	231	ALA	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	37	GLN	3.9
1	B	29	VAL	3.9
1	C	67	ALA	3.9
1	C	81	ALA	3.8
1	C	105	ALA	3.8
1	C	78	GLN	3.8
1	C	74	GLU	3.8
1	C	83	TYR	3.8
1	C	101	CYS	3.8
1	C	92	ASP	3.7
1	C	11	ASN	3.7
1	C	232	GLY	3.7
1	C	89	ASP	3.7
1	A	99	TRP	3.7
1	A	29	VAL	3.6
1	B	27	VAL	3.6
1	C	94	LYS	3.6
1	C	99	TRP	3.5
1	C	71	THR	3.5
1	C	76	THR	3.5
1	C	258	GLU	3.5
1	C	79	LYS	3.5
1	C	256	VAL	3.5
1	C	91	LEU	3.4
1	C	206	LYS	3.4
1	C	25	ASP	3.4
1	C	243	PHE	3.4
1	B	28	LEU	3.4
1	A	107	THR	3.3
1	A	140	VAL	3.3
1	C	15	TRP	3.3
1	A	103	SER	3.2
1	C	30	PRO	3.2
1	C	262	LYS	3.2
1	B	136	SER	3.2
1	C	112	LEU	3.2
1	B	127	LEU	3.2
1	B	30	PRO	3.1
1	C	221	PHE	3.1
1	C	117	ASP	3.1
1	C	56	PHE	3.1
1	A	104	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	137	ASN	3.0
1	A	30	PRO	3.0
1	A	101	CYS	3.0
1	C	9	THR	3.0
1	C	86	SER	3.0
1	B	98	VAL	3.0
1	C	48	ALA	2.9
1	A	33	LEU	2.9
1	A	108	VAL	2.8
1	C	50	GLY	2.8
1	A	102	ALA	2.8
1	A	100	GLY	2.8
1	B	101	CYS	2.8
1	C	238	LEU	2.7
1	B	39	PHE	2.7
1	A	32	GLY	2.7
1	C	113	LEU	2.7
1	C	68	PRO	2.7
1	C	207	PRO	2.6
1	C	53	VAL	2.6
1	C	88	LEU	2.6
1	C	253	ALA	2.6
1	A	105	ALA	2.6
1	C	20	GLU	2.5
1	A	124	CYS	2.5
1	C	69	PRO	2.5
1	B	100	GLY	2.5
1	C	109	VAL	2.5
1	B	99	TRP	2.5
1	C	104	GLY	2.5
1	C	103	SER	2.4
1	C	201	GLU	2.4
1	C	199	ASP	2.4
1	B	33	LEU	2.4
1	C	43	VAL	2.3
1	C	57	ASP	2.3
1	A	166	GLN	2.3
1	A	58	MET	2.3
1	A	106	SER	2.3
1	C	100	GLY	2.3
1	B	166	GLN	2.3
1	C	251	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	123	MET	2.2
1	A	80	LEU	2.1
1	C	250	ASP	2.1
1	C	82	SER	2.1
1	A	31	ASP	2.1
1	A	34	GLY	2.0
1	C	214	ALA	2.0
1	C	107	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	36J	B	301	23/23	0.91	0.29	2.09	39,55,91,94	0
2	36J	A	300	23/23	0.86	0.31	1.67	41,64,100,103	0

6.5 Other polymers [i](#)

There are no such residues in this entry.