



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:46 AM GMT

PDB ID : 2IGN
Title : Crystal structure of recombinant pyranose 2-oxidase H167A mutant
Authors : Divne, C.
Deposited on : 2006-09-22
Resolution : 1.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1226 (1.66-1.66)
Clashscore	102246	1323 (1.66-1.66)
Ramachandran outliers	100387	1295 (1.66-1.66)
Sidechain outliers	100360	1295 (1.66-1.66)
RSRZ outliers	91569	1227 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	623	<div> <div>4%</div> <div>78%</div> <div>13%</div> <div>7%</div> </div>
1	B	623	<div> <div>3%</div> <div>77%</div> <div>14%</div> <div>7%</div> </div>
1	C	623	<div> <div>5%</div> <div>79%</div> <div>12%</div> <div>7%</div> </div>
1	D	623	<div> <div>4%</div> <div>79%</div> <div>13%</div> <div>7%</div> </div>
1	E	623	<div> <div>5%</div> <div>79%</div> <div>13%</div> <div>7%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	623	
1	G	623	
1	H	623	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MES	A	8002	-	-	-	X
3	MES	B	8003	-	-	X	X
3	MES	C	8008	-	-	X	X
3	MES	D	8005	-	-	X	X
3	MES	E	8007	-	-	X	X
3	MES	G	8006	-	-	X	X
3	MES	H	8004	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 41758 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyranose oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	577	Total	C	N	O	S	0	1	0
			4551	2874	777	875	25			
1	B	577	Total	C	N	O	S	0	2	0
			4560	2879	779	877	25			
1	D	577	Total	C	N	O	S	0	2	0
			4560	2879	779	877	25			
1	C	577	Total	C	N	O	S	0	2	0
			4560	2879	779	877	25			
1	E	577	Total	C	N	O	S	0	2	0
			4560	2879	779	877	25			
1	F	577	Total	C	N	O	S	0	1	0
			4551	2874	777	875	25			
1	H	577	Total	C	N	O	S	0	3	0
			4568	2884	780	878	26			
1	G	577	Total	C	N	O	S	0	3	0
			4570	2885	782	878	25			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	167	ALA	HIS	ENGINEERED	UNP Q7ZA32
B	167	ALA	HIS	ENGINEERED	UNP Q7ZA32
C	167	ALA	HIS	ENGINEERED	UNP Q7ZA32
D	167	ALA	HIS	ENGINEERED	UNP Q7ZA32
E	167	ALA	HIS	ENGINEERED	UNP Q7ZA32
F	167	ALA	HIS	ENGINEERED	UNP Q7ZA32
G	167	ALA	HIS	ENGINEERED	UNP Q7ZA32
H	167	ALA	HIS	ENGINEERED	UNP Q7ZA32

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	H	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	G	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	G	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	705	Total	O	0	0
			705	705		
4	B	670	Total	O	0	0
			670	670		
4	C	528	Total	O	0	0
			528	528		
4	D	577	Total	O	0	0
			577	577		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	565	Total 565	O 565	0	0
4	F	498	Total 498	O 498	0	0
4	G	604	Total 604	O 604	0	0
4	H	611	Total 611	O 611	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Pyranose oxidase




• Molecule 1: Pyranose oxidase





• Molecule 1: Pyranose oxidase







• Molecule 1: Pyranose oxidase



• Molecule 1: Pyranose oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	168.07Å 103.00Å 168.56Å 90.00° 106.43° 90.00°	Depositor
Resolution (Å)	39.00 – 1.65 38.98 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.00-1.65) 99.9 (38.98-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.172 , 0.205 0.179 , 0.209	Depositor DCC
R_{free} test set	6563 reflections (1.00%)	DCC
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.6	EDS
Estimated twinning fraction	0.014 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 659652 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	41758	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 19.82 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.9631e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	5/4666 (0.1%)	0.92	11/6345 (0.2%)
1	B	0.90	2/4675 (0.0%)	0.92	13/6357 (0.2%)
1	C	0.84	0/4675	0.86	7/6357 (0.1%)
1	D	0.82	2/4675 (0.0%)	0.87	11/6357 (0.2%)
1	E	0.81	1/4675 (0.0%)	0.84	5/6357 (0.1%)
1	F	0.81	2/4666 (0.0%)	0.84	5/6345 (0.1%)
1	G	0.89	5/4686 (0.1%)	0.90	11/6372 (0.2%)
1	H	0.89	2/4683 (0.0%)	0.92	12/6367 (0.2%)
All	All	0.86	19/37401 (0.1%)	0.88	75/50857 (0.1%)

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	310	GLU	CG-CD	7.49	1.63	1.51
1	D	310	GLU	CB-CG	7.16	1.65	1.52
1	B	482	GLU	CG-CD	6.94	1.62	1.51
1	G	81	ASP	CA-C	6.58	1.70	1.52
1	B	421	GLU	CB-CG	5.97	1.63	1.52

The worst 5 of 75 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	139	ARG	NE-CZ-NH2	-19.15	110.72	120.30
1	G	139	ARG	NE-CZ-NH2	-18.35	111.12	120.30
1	E	139	ARG	NE-CZ-NH2	-17.48	111.56	120.30
1	G	139	ARG	NE-CZ-NH1	17.45	129.03	120.30
1	F	139	ARG	NE-CZ-NH2	-17.08	111.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4551	0	4401	126	0
1	B	4560	0	4408	152	0
1	C	4560	0	4408	107	0
1	D	4560	0	4408	93	0
1	E	4560	0	4408	98	0
1	F	4551	0	4401	74	0
1	G	4570	0	4414	170	0
1	H	4568	0	4416	163	1
2	A	53	0	31	0	0
2	B	53	0	30	2	0
2	C	53	0	28	1	0
2	D	53	0	30	1	0
2	E	53	0	29	0	0
2	F	53	0	30	1	0
2	G	53	0	30	1	0
2	H	53	0	29	0	0
3	A	12	0	12	4	0
3	B	12	0	12	9	0
3	C	12	0	12	10	0
3	D	12	0	12	14	0
3	E	12	0	12	11	0
3	F	12	0	12	4	0
3	G	12	0	12	9	0
3	H	12	0	12	17	0
4	A	705	0	0	100	2
4	B	670	0	0	111	1
4	C	528	0	0	73	0
4	D	577	0	0	56	0
4	E	565	0	0	64	0
4	F	498	0	0	48	0
4	G	604	0	0	110	3
4	H	611	0	0	122	1
All	All	41758	0	35597	993	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 993 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:528:LYS:HE3	4:B:8620:HOH:O	1.23	1.38
1:B:452:ASP:HB2	4:B:8464:HOH:O	1.26	1.34
1:G:104[A]:VAL:HG23	4:G:8521:HOH:O	1.28	1.33
1:B:142:SER:HB3	4:B:8632:HOH:O	1.28	1.33
1:A:204:SER:HB2	4:A:8479:HOH:O	1.16	1.31

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:8534:HOH:O	4:G:8089:HOH:O[2_646]	1.69	0.51
4:A:8707:HOH:O	4:G:8089:HOH:O[2_646]	1.73	0.47
4:H:8261:HOH:O	4:G:8557:HOH:O[2_655]	1.81	0.39
1:H:235:ASP:OD2	4:B:8535:HOH:O[2_666]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	576/623 (92%)	563 (98%)	13 (2%)	0	100	100
1	B	577/623 (93%)	564 (98%)	13 (2%)	0	100	100
1	C	577/623 (93%)	558 (97%)	19 (3%)	0	100	100
1	D	577/623 (93%)	564 (98%)	13 (2%)	0	100	100
1	E	577/623 (93%)	561 (97%)	15 (3%)	1 (0%)	52	30
1	F	576/623 (92%)	563 (98%)	13 (2%)	0	100	100
1	G	578/623 (93%)	562 (97%)	16 (3%)	0	100	100
1	H	578/623 (93%)	562 (97%)	16 (3%)	0	100	100
All	All	4616/4984 (93%)	4497 (97%)	118 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	343	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	505/541 (93%)	490 (97%)	15 (3%)	48	18
1	B	506/541 (94%)	492 (97%)	14 (3%)	51	21
1	C	506/541 (94%)	494 (98%)	12 (2%)	57	28
1	D	506/541 (94%)	494 (98%)	12 (2%)	57	28
1	E	506/541 (94%)	496 (98%)	10 (2%)	63	38
1	F	505/541 (93%)	491 (97%)	14 (3%)	51	21
1	G	507/541 (94%)	497 (98%)	10 (2%)	63	38
1	H	507/541 (94%)	493 (97%)	14 (3%)	51	21
All	All	4048/4328 (94%)	3947 (98%)	101 (2%)	55	26

5 of 101 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	389	LEU
1	E	206	PHE
1	G	385	THR
1	C	418	GLN
1	C	496	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	105	ASN
1	F	105	ASN
1	G	108	GLN
1	E	418	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	108	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	7001	-	48,58,58	1.55	9 (18%)	54,89,89	3.40	11 (20%)
3	MES	A	8002	-	11,12,12	0.61	0	14,16,16	5.54	6 (42%)
2	FAD	B	7002	-	48,58,58	1.11	1 (2%)	54,89,89	3.38	17 (31%)
3	MES	B	8003	-	11,12,12	0.57	0	14,16,16	6.38	6 (42%)
2	FAD	C	7004	-	48,58,58	1.44	9 (18%)	54,89,89	2.94	16 (29%)
3	MES	C	8008	-	11,12,12	0.58	0	14,16,16	4.98	7 (50%)
2	FAD	D	7003	-	48,58,58	1.32	6 (12%)	54,89,89	2.63	18 (33%)
3	MES	D	8005	-	11,12,12	0.81	0	14,16,16	8.94	6 (42%)
2	FAD	E	7005	-	48,58,58	1.41	7 (14%)	54,89,89	3.09	16 (29%)
3	MES	E	8007	-	11,12,12	0.80	0	14,16,16	6.76	10 (71%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	F	7006	-	48,58,58	1.34	9 (18%)	54,89,89	3.26	17 (31%)
3	MES	F	8001	-	11,12,12	0.70	0	14,16,16	3.98	7 (50%)
2	FAD	G	7008	-	48,58,58	1.34	9 (18%)	54,89,89	3.19	15 (27%)
3	MES	G	8006	-	11,12,12	0.65	0	14,16,16	7.59	7 (50%)
2	FAD	H	7007	-	48,58,58	1.29	5 (10%)	54,89,89	3.12	16 (29%)
3	MES	H	8004	-	11,12,12	0.50	0	14,16,16	4.84	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	7001	-	-	0/30/50/50	0/6/6/6
3	MES	A	8002	-	-	0/6/14/14	0/1/1/1
2	FAD	B	7002	-	-	0/30/50/50	0/6/6/6
3	MES	B	8003	-	-	0/6/14/14	0/1/1/1
2	FAD	C	7004	-	-	0/30/50/50	0/6/6/6
3	MES	C	8008	-	-	0/6/14/14	0/1/1/1
2	FAD	D	7003	-	-	0/30/50/50	0/6/6/6
3	MES	D	8005	-	-	0/6/14/14	0/1/1/1
2	FAD	E	7005	-	-	0/30/50/50	0/6/6/6
3	MES	E	8007	-	-	0/6/14/14	0/1/1/1
2	FAD	F	7006	-	-	0/30/50/50	0/6/6/6
3	MES	F	8001	-	-	0/6/14/14	0/1/1/1
2	FAD	G	7008	-	-	0/30/50/50	0/6/6/6
3	MES	G	8006	-	-	0/6/14/14	0/1/1/1
2	FAD	H	7007	-	-	0/30/50/50	0/6/6/6
3	MES	H	8004	-	-	0/6/14/14	0/1/1/1

The worst 5 of 55 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	7003	FAD	O4B-C4B	-3.62	1.36	1.45
2	A	7001	FAD	C10-N10	-3.43	1.35	1.39
2	C	7004	FAD	O4B-C4B	-3.30	1.37	1.45
2	E	7005	FAD	C10-N10	-3.29	1.35	1.39
2	D	7003	FAD	C1'-N10	-3.19	1.45	1.48

The worst 5 of 182 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	8005	MES	O2S-S-C8	-31.17	80.31	106.91
3	G	8006	MES	O2S-S-C8	-26.05	84.68	106.91
3	B	8003	MES	O1S-S-C8	-18.70	90.95	106.91
3	E	8007	MES	O1S-S-C8	-18.18	91.40	106.91
3	A	8002	MES	O1S-S-C8	-16.92	92.47	106.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

13 monomers are involved in 84 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	8002	MES	4	0
2	B	7002	FAD	2	0
3	B	8003	MES	9	0
2	C	7004	FAD	1	0
3	C	8008	MES	10	0
2	D	7003	FAD	1	0
3	D	8005	MES	14	0
3	E	8007	MES	11	0
2	F	7006	FAD	1	0
3	F	8001	MES	4	0
2	G	7008	FAD	1	0
3	G	8006	MES	9	0
3	H	8004	MES	17	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	577/623 (92%)	-0.09	25 (4%)	39	38	11, 16, 37, 60	0
1	B	577/623 (92%)	-0.10	21 (3%)	46	47	11, 17, 36, 54	0
1	C	577/623 (92%)	0.16	30 (5%)	31	28	13, 23, 43, 64	0
1	D	577/623 (92%)	0.01	25 (4%)	39	38	13, 21, 40, 59	0
1	E	577/623 (92%)	0.04	32 (5%)	29	26	14, 21, 41, 58	0
1	F	577/623 (92%)	0.07	30 (5%)	31	28	13, 23, 41, 61	0
1	G	577/623 (92%)	-0.05	25 (4%)	39	38	13, 19, 38, 57	0
1	H	577/623 (92%)	-0.17	20 (3%)	48	48	12, 18, 35, 55	0
All	All	4616/4984 (92%)	-0.02	208 (4%)	37	35	11, 20, 40, 64	0

The worst 5 of 208 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	619	THR	13.1
1	C	619	THR	12.9
1	H	619	THR	12.0
1	A	619	THR	11.6
1	E	619	THR	11.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MES	E	8007	12/12	0.91	0.19	2.99	29,33,38,39	0
3	MES	G	8006	12/12	0.91	0.19	2.73	24,30,33,33	0
3	MES	A	8002	12/12	0.97	0.14	2.65	22,26,28,28	0
3	MES	D	8005	12/12	0.93	0.17	2.47	30,32,34,36	0
3	MES	C	8008	12/12	0.93	0.19	2.43	27,32,38,39	0
3	MES	B	8003	12/12	0.92	0.18	2.09	28,32,34,35	0
3	MES	H	8004	12/12	0.93	0.17	1.65	27,33,34,35	0
3	MES	F	8001	12/12	0.96	0.10	0.79	20,22,24,25	0
2	FAD	F	7006	53/53	0.98	0.10	0.31	14,17,20,22	0
2	FAD	G	7008	53/53	0.98	0.09	0.17	11,14,15,17	0
2	FAD	D	7003	53/53	0.98	0.09	0.07	14,17,19,22	0
2	FAD	A	7001	53/53	0.99	0.09	0.01	10,12,14,16	0
2	FAD	H	7007	53/53	0.99	0.09	-0.11	10,14,17,19	0
2	FAD	B	7002	53/53	0.99	0.09	-0.21	10,13,16,20	0
2	FAD	E	7005	53/53	0.98	0.09	-0.22	15,17,20,21	0
2	FAD	C	7004	53/53	0.98	0.09	-0.27	15,18,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.