



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:10 PM GMT

PDB ID : 4IGH
Title : High resolution crystal structure of human dihydroorotate dehydrogenase bound with 4-quinoline carboxylic acid analog
Authors : Deng, X.; Das, P.; Fontoura, B.M.A.; Phillips, M.A.; De Brabander, J.K.
Deposited on : 2012-12-17
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1475 (1.34-1.26)
Clashscore	102246	1031 (1.32-1.28)
Ramachandran outliers	100387	1504 (1.34-1.26)
Sidechain outliers	100360	1503 (1.34-1.26)
RSRZ outliers	91569	1476 (1.34-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FMN	A	501	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	506	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6321 atoms, of which 3039 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dihydroorotate dehydrogenase (quinone), mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	356	5830	1804	2950	530	542	4	853	22	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	397	LEU	-	EXPRESSION TAG	UNP Q02127
A	398	GLU	-	EXPRESSION TAG	UNP Q02127
A	399	HIS	-	EXPRESSION TAG	UNP Q02127
A	400	HIS	-	EXPRESSION TAG	UNP Q02127
A	401	HIS	-	EXPRESSION TAG	UNP Q02127
A	402	HIS	-	EXPRESSION TAG	UNP Q02127
A	403	HIS	-	EXPRESSION TAG	UNP Q02127
A	404	HIS	-	EXPRESSION TAG	UNP Q02127

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	
			50	17	19	4	9	1	
								9	0

- Molecule 3 is OROTIC ACID (three-letter code: ORO) (formula: $C_5H_4N_2O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O		
			14	5	3	2	4		
								0	0

- Molecule 4 is 6-FLUORO-2-[2-METHYL-4-PHENOXY-5-(PROPAN-2-YL)PHENYL]QUINOLINE-4-CARBOXYLIC ACID (three-letter code: 1EA) (formula: $C_{26}H_{22}FNO_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	H	N	O	
			50	26	1	19	1	3	
								6	0

- Molecule 5 is 3-[DECYL(DIMETHYL)AMMONIO]PROPANE-1-SULFONATE (three-letter code: ZWI) (formula: $C_{15}H_{33}NO_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	S	
			53	15	33	1	3	1	
								9	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	1	0
			13	3	7	3		
6	A	1	Total	C	H	O	2	0
			14	3	8	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	297	Total	O	0	0
			297	297		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase (quinone), mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.79 Å 90.79 Å 122.48 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	78.62 – 1.30 28.53 – 1.22	Depositor EDS
% Data completeness (in resolution range)	99.5 (78.62-1.30) 99.4 (28.53-1.22)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 1.22 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.138 , 0.148 0.214 , 0.222	Depositor DCC
R_{free} test set	4312 reflections (3.11%)	DCC
Wilson B-factor (Å ²)	12.2	Xtriage
Anisotropy	0.236	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.51 , 75.0	EDS
Estimated twinning fraction	0.016 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	2 of 171726 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6321	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1EA, ZWI, ORO, FMN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.49	34/2944 (1.2%)	1.28	34/3974 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64[A]	SER	N-CA	11.57	1.69	1.46
1	A	64[B]	SER	N-CA	11.57	1.69	1.46
1	A	62	PHE	N-CA	6.99	1.60	1.46
1	A	160	ARG	CB-CG	-6.92	1.33	1.52
1	A	300	GLU	CD-OE2	-6.89	1.18	1.25
1	A	295	GLY	C-O	-6.71	1.12	1.23
1	A	388[A]	ASP	N-CA	6.63	1.59	1.46
1	A	388[B]	ASP	N-CA	6.63	1.59	1.46
1	A	89[A]	ARG	CD-NE	-6.57	1.35	1.46
1	A	89[B]	ARG	CD-NE	-6.57	1.35	1.46
1	A	213[A]	VAL	N-CA	-6.47	1.33	1.46
1	A	213[B]	VAL	N-CA	-6.47	1.33	1.46
1	A	53	GLU	CD-OE2	-6.29	1.18	1.25
1	A	195	TYR	CE1-CZ	-5.82	1.30	1.38
1	A	61[A]	ARG	C-N	5.68	1.47	1.34
1	A	61[B]	ARG	C-N	5.68	1.47	1.34
1	A	342	ALA	C-O	-5.66	1.12	1.23
1	A	216	PRO	N-CD	5.65	1.55	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	TYR	CG-CD1	-5.62	1.31	1.39
1	A	266	GLU	CD-OE1	-5.54	1.19	1.25
1	A	212[A]	ASN	C-N	5.43	1.46	1.34
1	A	212[B]	ASN	C-N	5.43	1.46	1.34
1	A	300	GLU	CD-OE1	-5.30	1.19	1.25
1	A	211[A]	VAL	C-O	-5.26	1.13	1.23
1	A	211[B]	VAL	C-O	-5.26	1.13	1.23
1	A	339	GLY	C-O	-5.26	1.15	1.23
1	A	188[A]	SER	CB-OG	-5.24	1.35	1.42
1	A	188[B]	SER	CB-OG	-5.24	1.35	1.42
1	A	64[A]	SER	CA-C	5.20	1.66	1.52
1	A	64[B]	SER	CA-C	5.20	1.66	1.52
1	A	332	GLY	C-O	-5.19	1.15	1.23
1	A	168	GLN	C-O	-5.11	1.13	1.23
1	A	229	GLU	CD-OE2	5.09	1.31	1.25
1	A	243	GLY	C-O	-5.03	1.15	1.23

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	232	ARG	NE-CZ-NH1	15.02	127.81	120.30
1	A	232	ARG	NE-CZ-NH2	-10.39	115.11	120.30
1	A	211[A]	VAL	O-C-N	10.01	138.71	122.70
1	A	211[B]	VAL	O-C-N	10.01	138.71	122.70
1	A	200	ARG	NE-CZ-NH1	9.86	125.23	120.30
1	A	89[A]	ARG	NE-CZ-NH1	-9.13	115.73	120.30
1	A	89[B]	ARG	NE-CZ-NH1	-9.13	115.73	120.30
1	A	81[A]	VAL	CG1-CB-CG2	-9.04	96.43	110.90
1	A	81[B]	VAL	CG1-CB-CG2	-9.04	96.43	110.90
1	A	231	ARG	NE-CZ-NH1	8.43	124.52	120.30
1	A	82	ARG	NE-CZ-NH1	7.50	124.05	120.30
1	A	211[A]	VAL	C-N-CA	-7.42	103.16	121.70
1	A	211[B]	VAL	C-N-CA	-7.42	103.16	121.70
1	A	158	GLU	OE1-CD-OE2	-7.18	114.69	123.30
1	A	160	ARG	NE-CZ-NH2	6.61	123.61	120.30
1	A	211[A]	VAL	CA-C-N	-6.61	102.67	117.20
1	A	211[B]	VAL	CA-C-N	-6.61	102.67	117.20
1	A	36	ARG	NE-CZ-NH2	6.51	123.55	120.30
1	A	82	ARG	NE-CZ-NH2	-6.47	117.06	120.30
1	A	221	LEU	CA-CB-CG	5.99	129.08	115.30
1	A	387	THR	O-C-N	5.97	132.26	122.70
1	A	81[A]	VAL	CA-CB-CG2	5.94	119.81	110.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	81[B]	VAL	CA-CB-CG2	5.94	119.81	110.90
1	A	147	TYR	CB-CG-CD2	5.88	124.53	121.00
1	A	89[A]	ARG	NE-CZ-NH2	5.69	123.15	120.30
1	A	89[B]	ARG	NE-CZ-NH2	5.69	123.15	120.30
1	A	249	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	190	ASP	CB-CG-OD2	-5.60	113.26	118.30
1	A	347[A]	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	A	347[B]	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	A	327	ARG	NE-CZ-NH1	5.27	122.93	120.30
1	A	395	ARG	N-CA-C	5.15	124.90	111.00
1	A	245	ARG	NE-CZ-NH1	-5.03	117.78	120.30
1	A	393	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	212[A]	ASN	Mainchain
1	A	213[A]	VAL	Mainchain,Peptide
1	A	213[B]	VAL	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2880	2950	2929	52	0
2	A	31	19	19	0	0
3	A	11	3	3	1	0
4	A	31	19	20	2	0
5	A	20	33	33	0	0
6	A	12	15	16	4	0
7	A	297	0	0	7	0
All	All	3282	3039	3020	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149[A]:PHE:CZ	1:A:215:SER:HB2	1.91	1.05
1:A:149[A]:PHE:HZ	1:A:215:SER:HB2	1.17	1.05
1:A:149[A]:PHE:CZ	1:A:215:SER:CB	2.50	0.94
1:A:46:LEU:HD13	4:A:503:1EA:H251	1.53	0.91
1:A:149[A]:PHE:HZ	1:A:215:SER:CB	1.90	0.84
1:A:213[A]:VAL:HG22	1:A:214:SER:HB3	1.59	0.83
1:A:213[A]:VAL:CG2	1:A:214:SER:HB3	2.09	0.83
1:A:150:ASN:HD21	6:A:506:GOL:H12	1.47	0.79
1:A:149[A]:PHE:CE1	1:A:215:SER:CB	2.66	0.78
1:A:46:LEU:HD13	4:A:503:1EA:C25	2.15	0.76
1:A:87:LYS:NZ	1:A:89[A]:ARG:HG2	2.02	0.74
1:A:145:ASN:OD1	3:A:502:ORO:H5	1.91	0.70
1:A:149[A]:PHE:CE1	1:A:215:SER:OG	2.47	0.67
1:A:184[B]:LYS:HE2	1:A:188[B]:SER:HB3	1.76	0.67
1:A:61[B]:ARG:HD2	7:A:714:HOH:O	1.95	0.67
1:A:57:ARG:CZ	6:A:506:GOL:O3	2.43	0.66
1:A:213[A]:VAL:O	1:A:221:LEU:HG	1.96	0.65
1:A:34:ASP:N	7:A:870:HOH:O	2.29	0.65
1:A:213[B]:VAL:O	1:A:213[B]:VAL:HG22	2.01	0.60
1:A:150:ASN:ND2	6:A:506:GOL:H12	2.16	0.60
1:A:149[A]:PHE:HE1	1:A:215:SER:OG	1.85	0.58
1:A:56:HIS:HD2	1:A:147:TYR:OH	1.88	0.56
1:A:236:LYS:HE3	7:A:769:HOH:O	2.05	0.55
1:A:154:LEU:HD21	7:A:871:HOH:O	2.07	0.54
1:A:213[A]:VAL:HG23	1:A:214:SER:HB3	1.89	0.52
1:A:61[B]:ARG:NE	1:A:61[B]:ARG:HA	2.25	0.52
1:A:57:ARG:NE	6:A:506:GOL:O3	2.44	0.51
1:A:362:TRP:HE3	7:A:728:HOH:O	1.94	0.50
1:A:213[A]:VAL:HG23	1:A:214:SER:CA	2.41	0.50
1:A:123:PRO:HA	1:A:154:LEU:HG	1.95	0.49
1:A:56:HIS:HE1	1:A:98:PHE:O	1.96	0.47
1:A:213[B]:VAL:HG22	1:A:221:LEU:O	2.16	0.47
1:A:149[A]:PHE:HE1	1:A:215:SER:CB	2.23	0.46
1:A:213[A]:VAL:HG22	1:A:256:ILE:HA	1.97	0.46
1:A:321:TYR:OH	1:A:396:ARG:HD3	2.17	0.45
1:A:396:ARG:HD2	1:A:396:ARG:C	2.38	0.44
1:A:149[A]:PHE:CZ	1:A:215:SER:HB3	2.50	0.44
1:A:213[B]:VAL:O	1:A:213[B]:VAL:HG13	2.18	0.44
1:A:56:HIS:CD2	1:A:147:TYR:OH	2.70	0.43
1:A:213[A]:VAL:HG23	1:A:214:SER:CB	2.49	0.42
1:A:372:ARG:HG2	7:A:874:HOH:O	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149[A]:PHE:CE1	1:A:215:SER:HB3	2.54	0.42
1:A:241:ARG:HD2	1:A:241:ARG:C	2.41	0.41
1:A:364:PRO:N	1:A:365:PRO:CD	2.84	0.41
1:A:78:MET:SD	1:A:379:LYS:HE3	2.60	0.41
1:A:221:LEU:O	1:A:221:LEU:HD23	2.21	0.41
1:A:213[B]:VAL:CG2	1:A:221:LEU:O	2.69	0.41
1:A:167:LYS:NZ	7:A:837:HOH:O	2.52	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/372 (100%)	365 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/302 (103%)	305 (98%)	5 (2%)	70	30

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ARG
1	A	221	LEU
1	A	356	TYR
1	A	395	ARG
1	A	396	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FMN	A	501	-	31,33,33	1.51	7 (22%)	32,50,50	2.21	5 (15%)
3	ORO	A	502	-	4,11,11	1.52	1 (25%)	3,15,15	2.06	1 (33%)
4	1EA	A	503	-	31,34,34	2.44	10 (32%)	44,49,49	2.06	11 (25%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ZWI	A	504	-	19,19,19	1.88	1 (5%)	23,24,24	1.52	5 (21%)
6	GOL	A	505	-	5,5,5	1.23	0	5,5,5	1.84	2 (40%)
6	GOL	A	506	-	5,5,5	0.54	0	5,5,5	1.40	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	501	-	-	0/18/18/18	0/3/3/3
3	ORO	A	502	-	-	0/0/4/4	0/1/1/1
4	1EA	A	503	-	-	0/12/16/16	0/4/4/4
5	ZWI	A	504	-	-	0/19/19/19	0/0/0/0
6	GOL	A	505	-	-	0/4/4/4	0/0/0/0
6	GOL	A	506	-	-	0/4/4/4	0/0/0/0

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	1EA	C25-C24	-8.94	1.29	1.52
5	A	504	ZWI	C1-S1	-7.15	1.65	1.78
4	A	503	1EA	F1-C6	-3.66	1.27	1.36
4	A	503	1EA	O3-C16	-3.17	1.32	1.39
2	A	501	FMN	C2-N1	-2.91	1.32	1.38
2	A	501	FMN	C6-C5A	-2.65	1.37	1.41
3	A	502	ORO	C6-N1	-2.42	1.32	1.36
4	A	503	1EA	C26-C24	-2.40	1.46	1.52
4	A	503	1EA	C18-C17	-2.37	1.34	1.38
2	A	501	FMN	C9-C9A	-2.29	1.35	1.40
2	A	501	FMN	C10-N10	-2.28	1.36	1.39
2	A	501	FMN	C4-N3	-2.09	1.33	1.36
4	A	503	1EA	C9-N1	-2.03	1.34	1.37
4	A	503	1EA	C3-C4	2.14	1.47	1.42
4	A	503	1EA	C13-C14	2.69	1.45	1.39
2	A	501	FMN	C4-C4A	2.86	1.46	1.40
2	A	501	FMN	C4A-C10	3.63	1.47	1.40
4	A	503	1EA	C10-C11	3.68	1.46	1.40
4	A	503	1EA	C4-C9	4.13	1.48	1.42

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FMN	C4A-C10-N10	-4.89	116.97	120.52
4	A	503	1EA	C3-C4-C9	-3.79	115.29	117.47
4	A	503	1EA	C2-C1-C10	-3.38	116.98	122.15
5	A	504	ZWI	O1-S1-O3	-3.21	105.94	112.51
6	A	505	GOL	O2-C2-C1	-2.88	94.65	108.47
6	A	505	GOL	C3-C2-C1	-2.84	99.10	111.06
4	A	503	1EA	C15-C14-C24	-2.64	113.59	118.85
4	A	503	1EA	C8-C9-C4	-2.41	116.53	119.09
5	A	504	ZWI	C15-N1-C3	-2.20	103.64	109.42
4	A	503	1EA	C12-C13-C14	-2.12	118.40	121.83
5	A	504	ZWI	O2-S1-C1	2.15	111.73	105.91
4	A	503	1EA	C6-C5-C4	2.22	120.79	118.75
2	A	501	FMN	C6-C5A-N5	2.25	121.72	118.92
6	A	506	GOL	O3-C3-C2	2.42	122.27	109.97
4	A	503	1EA	C1-N1-C9	2.51	119.90	118.04
5	A	504	ZWI	C5-C6-C7	2.66	128.38	114.54
2	A	501	FMN	C4A-N5-C5A	2.78	120.00	116.72
4	A	503	1EA	C13-O3-C16	2.98	125.76	117.83
4	A	503	1EA	C26-C24-C25	3.39	118.49	110.32
5	A	504	ZWI	C4-C5-C6	3.54	126.43	110.71
2	A	501	FMN	C5A-C9A-N10	3.55	120.23	117.58
3	A	502	ORO	C4-C5-C6	3.56	120.50	116.87
4	A	503	1EA	C13-C14-C24	3.74	126.30	120.79
4	A	503	1EA	C25-C24-C14	8.30	126.08	111.77
2	A	501	FMN	C2-N1-C10	8.87	120.95	113.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	502	ORO	1	0
4	A	503	1EA	2	0
6	A	506	GOL	4	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/372 (95%)	1.21	76 (21%) 1 1	4, 8, 26, 84	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	37	PHE	12.5
1	A	189	VAL	9.6
1	A	45	THR	9.3
1	A	221	LEU	9.2
1	A	46	LEU	8.4
1	A	42	LEU	8.3
1	A	219	ALA	8.2
1	A	218	THR	8.1
1	A	84	LEU	7.9
1	A	38	TYR	7.6
1	A	69	PRO	7.3
1	A	213[A]	VAL	7.1
1	A	36	ARG	6.7
1	A	39	ALA	6.4
1	A	49	LEU	6.1
1	A	68	LEU	6.1
1	A	41	HIS	6.0
1	A	220	GLY	5.9
1	A	34	ASP	5.7
1	A	67	LEU	5.6
1	A	226	GLY	5.5
1	A	395	ARG	5.3
1	A	74	GLN	5.1
1	A	44	PRO	5.0
1	A	40	GLU	5.0
1	A	48	GLY	5.0
1	A	43	MET	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	396	ARG	4.6
1	A	70	ARG	4.5
1	A	228	ALA	4.4
1	A	65	LEU	4.1
1	A	35	GLU	3.9
1	A	187	THR	3.8
1	A	78	MET	3.6
1	A	384	GLY	3.4
1	A	180	VAL	3.4
1	A	382	GLY	3.3
1	A	66	GLY	3.3
1	A	47	GLN	3.2
1	A	229	GLU	3.2
1	A	215	SER	3.1
1	A	217	ASN	3.1
1	A	170	LYS	3.1
1	A	118	ILE	3.0
1	A	173[A]	GLU	3.0
1	A	372	ARG	3.0
1	A	76	SER	2.9
1	A	62	PHE	2.8
1	A	116	VAL	2.8
1	A	85	GLY	2.7
1	A	216	PRO	2.7
1	A	212[A]	ASN	2.7
1	A	77	ASP	2.6
1	A	353	VAL	2.6
1	A	209	LEU	2.6
1	A	388[A]	ASP	2.6
1	A	182	LEU	2.5
1	A	199	VAL	2.5
1	A	190	ASP	2.5
1	A	210	VAL	2.5
1	A	202	LEU	2.5
1	A	188[A]	SER	2.5
1	A	82	ARG	2.4
1	A	86	HIS	2.4
1	A	75	ASP	2.3
1	A	246	ARG	2.3
1	A	309	LEU	2.3
1	A	327	ARG	2.2
1	A	282	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	94	ILE	2.2
1	A	333	VAL	2.2
1	A	336	VAL	2.1
1	A	232	ARG	2.1
1	A	385	GLY	2.1
1	A	281	ILE	2.1
1	A	355	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FMN	A	501	31/31	0.96	0.18	2.01	7,8,18,18	9
6	GOL	A	506	6/6	0.82	0.15	1.31	13,25,60,64	2
5	ZWI	A	504	20/20	0.76	0.16	0.84	14,34,62,103	9
3	ORO	A	502	11/11	0.95	0.14	0.24	7,9,12,12	0
6	GOL	A	505	6/6	0.97	0.10	0.08	11,26,35,55	1
4	1EA	A	503	31/31	0.94	0.12	-0.27	10,18,32,33	6

6.5 Other polymers [i](#)

There are no such residues in this entry.