



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:58 PM GMT

PDB ID : 6INS
Title : X-RAY ANALYSIS OF THE SINGLE CHAIN B29-A1 PEPTIDE-LINKED
INSULIN MOLECULE. A COMPLETELY INACTIVE ANALOGUE
Authors : Derewenda, U.; Derewenda, Z.; Dodson, E.J.; Dodson, G.G.; Bing, X.;
Markussen, J.
Deposited on : 1992-11-25
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	50	
1	F	50	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INSULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	50	Total	C	N	O	S	0	0	0
			394	250	64	74	6			
1	F	50	Total	C	N	O	S	8	0	0
			394	252	63	73	6			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	F	1	Total	Zn	0	0
			1	1		
2	E	1	Total	Zn	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	97	Total	O	0	0
			97	97		
3	F	82	Total	O	0	0
			82	82		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: INSULIN

Chain E: 



• Molecule 1: INSULIN

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	79.22Å 79.22Å 37.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.00 9.90 – 2.02	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 86.4 (9.90-2.02)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.73 (at 2.02Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.179 , (Not available) 0.175 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.133	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.12 , 60.2	EDS
Estimated twinning fraction	0.043 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 5237 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	969	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	1.39	1/403 (0.2%)	2.81	38/543 (7.0%)
1	F	1.79	5/403 (1.2%)	2.34	18/543 (3.3%)
All	All	1.60	6/806 (0.7%)	2.58	56/1086 (5.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
1	F	0	1
All	All	0	2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	8(C)	THR	CB-OG1	19.27	1.81	1.43
1	F	8(C)	THR	CB-CG2	-12.38	1.11	1.52
1	F	21(D)	GLU	CB-CG	-10.81	1.31	1.52
1	F	3(C)	VAL	CB-CG1	9.04	1.71	1.52
1	F	3(C)	VAL	CB-CG2	-6.23	1.39	1.52
1	E	4(A)	GLU	CD-OE2	-5.53	1.19	1.25

All (56) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	16(B)	TYR	CB-CG-CD1	18.50	132.10	121.00
1	F	14(C)	TYR	CB-CG-CD1	-13.35	112.99	121.00
1	E	19(A)	TYR	CB-CG-CD1	-12.02	113.79	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	24(D)	PHE	CB-CG-CD2	-11.95	112.43	120.80
1	F	8(C)	THR	CA-CB-OG1	-11.09	85.72	109.00
1	F	14(C)	TYR	CB-CG-CD2	10.66	127.40	121.00
1	E	3(B)	ASN	CA-CB-CG	10.55	136.60	113.40
1	E	18(A)	ASN	CA-CB-CG	10.22	135.89	113.40
1	E	19(A)	TYR	CB-CG-CD2	10.15	127.09	121.00
1	F	3(C)	VAL	CA-CB-CG1	-8.37	98.35	110.90
1	F	8(C)	THR	CA-CB-CG2	8.28	124.00	112.40
1	E	16(B)	TYR	CB-CG-CD2	-8.27	116.04	121.00
1	E	13(B)	GLU	CB-CG-CD	7.61	134.76	114.20
1	F	18(D)	VAL	CA-CB-CG1	7.61	122.31	110.90
1	E	7(B)	CYS	CA-CB-SG	-7.09	101.23	114.00
1	E	12(B)	VAL	N-CA-CB	-7.00	96.11	111.50
1	E	18(A)	ASN	O-C-N	6.99	133.88	122.70
1	F	5(C)	GLN	CB-CG-CD	6.98	129.74	111.60
1	E	4(A)	GLU	CA-CB-CG	6.82	128.41	113.40
1	E	26(B)	TYR	CB-CG-CD2	-6.71	116.97	121.00
1	F	21(D)	GLU	CB-CG-CD	-6.70	96.11	114.20
1	E	17(A)	GLU	N-CA-CB	6.62	122.51	110.60
1	E	3(A)	VAL	N-CA-CB	-6.45	97.30	111.50
1	F	17(C)	GLU	CG-CD-OE1	6.42	131.14	118.30
1	E	12(B)	VAL	CB-CA-C	6.27	123.31	111.40
1	E	26(B)	TYR	CB-CG-CD1	6.20	124.72	121.00
1	E	16(B)	TYR	CG-CD1-CE1	6.18	126.25	121.30
1	F	24(D)	PHE	CB-CG-CD1	6.15	125.10	120.80
1	F	22(D)	ARG	O-C-N	6.13	133.63	123.20
1	E	4(A)	GLU	CG-CD-OE1	-6.03	106.24	118.30
1	F	17(D)	LEU	CA-CB-CG	5.99	129.09	115.30
1	F	6(C)	CYS	CA-CB-SG	5.84	124.51	114.00
1	E	19(A)	TYR	O-C-N	-5.83	113.38	122.70
1	E	10(B)	HIS	N-CA-CB	5.79	121.03	110.60
1	E	24(B)	PHE	CB-CG-CD2	-5.74	116.78	120.80
1	F	15(D)	LEU	CA-CB-CG	5.73	128.49	115.30
1	E	13(A)	LEU	CB-CA-C	5.73	121.08	110.20
1	E	20(A)	CYS	CA-CB-SG	5.72	124.30	114.00
1	E	2(A)	ILE	C-N-CA	5.71	135.97	121.70
1	E	12(B)	VAL	CA-CB-CG1	5.69	119.44	110.90
1	E	17(A)	GLU	CA-CB-CG	5.69	125.92	113.40
1	E	16(B)	TYR	CD1-CG-CD2	-5.68	111.65	117.90
1	E	7(B)	CYS	O-C-N	-5.64	113.62	123.20
1	E	11(B)	LEU	N-CA-CB	-5.59	99.21	110.40
1	E	15(B)	LEU	O-C-N	-5.46	113.96	122.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	1(B)	PHE	CB-CG-CD2	-5.37	117.04	120.80
1	E	20(A)	CYS	O-C-N	5.35	131.26	122.70
1	E	19(A)	TYR	CD1-CE1-CZ	-5.24	115.08	119.80
1	E	18(A)	ASN	N-CA-CB	5.24	120.03	110.60
1	E	5(A)	GLN	O-C-N	5.21	131.04	122.70
1	E	1(B)	PHE	CB-CG-CD1	5.14	124.39	120.80
1	E	8(A)	THR	N-CA-CB	-5.13	100.55	110.30
1	F	4(C)	GLU	CG-CD-OE2	5.13	128.56	118.30
1	F	2(D)	VAL	CA-C-O	5.09	130.79	120.10
1	F	2(D)	VAL	CA-C-N	-5.03	106.14	117.20
1	E	21(B)	GLU	CG-CD-OE1	5.01	128.32	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	22(B)	ARG	Sidechain
1	F	22(D)	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	394	0	363	21	1
1	F	394	0	366	19	2
2	E	1	0	0	0	0
2	F	1	0	0	1	0
3	E	97	0	0	9	13
3	F	82	0	0	4	10
All	All	969	0	729	38	15

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:30:ZN:ZN	3:F:81:HOH:O	1.18	0.93
1:E:18(A):ASN:C	3:E:74:HOH:O	2.06	0.93
1:E:18(A):ASN:O	3:E:74:HOH:O	1.86	0.91
1:E:18(A):ASN:O	3:E:73:HOH:O	1.94	0.85
1:F:11(D):LEU:HD21	1:F:2(C):ILE:HG12	1.59	0.85
1:E:28(B):PRO:HB2	3:E:83:HOH:O	1.92	0.69
1:E:3(B):ASN:ND2	3:E:36:HOH:O	2.26	0.69
1:F:18(C):ASN:ND2	3:F:111:HOH:O	2.29	0.66
1:E:14(B):ALA:O	1:E:18(B):VAL:HG23	1.97	0.65
1:F:2(C):ILE:HD12	1:F:19(C):TYR:CD2	2.32	0.64
1:F:12(C):SER:O	1:F:14(C):TYR:N	2.31	0.64
1:F:12(C):SER:O	1:F:13(C):LEU:C	2.35	0.64
1:F:3(C):VAL:O	1:F:5(C):GLN:N	2.34	0.61
1:E:27(B):THR:HG22	1:E:29(B):LYS:HD2	1.83	0.61
1:F:9(C):SER:HB3	3:F:36:HOH:O	2.00	0.61
1:E:3(B):ASN:OD1	3:E:85:HOH:O	2.16	0.60
1:E:29(B):LYS:HE2	3:F:65:HOH:O	2.03	0.59
1:E:22(B):ARG:O	1:E:21(A):ASN:HB2	2.03	0.59
1:E:3(B):ASN:ND2	3:E:35:HOH:O	2.25	0.57
1:E:29(B):LYS:HA	3:E:77:HOH:O	2.04	0.57
1:F:12(C):SER:O	1:F:15(C):GLN:N	2.39	0.56
1:F:3(C):VAL:C	1:F:5(C):GLN:N	2.61	0.53
1:F:3(C):VAL:C	1:F:5(C):GLN:H	2.12	0.53
1:F:28(D):PRO:O	1:F:29(D):LYS:HG3	2.08	0.53
1:F:5(C):GLN:O	1:F:9(C):SER:HB2	2.09	0.52
1:E:26(B):TYR:CD1	1:E:2(A):ILE:HD12	2.48	0.48
1:E:19(A):TYR:HA	3:E:74:HOH:O	2.13	0.48
1:F:2(C):ILE:HD12	1:F:19(C):TYR:CE2	2.49	0.47
1:E:27(B):THR:O	1:E:29(B):LYS:N	2.47	0.47
1:E:15(A):GLN:O	1:E:18(A):ASN:ND2	2.49	0.46
1:F:5(C):GLN:HA	1:F:9(C):SER:HB2	1.98	0.46
1:F:12(C):SER:C	1:F:14(C):TYR:N	2.69	0.45
1:E:25(B):PHE:CE2	1:F:25(D):PHE:HB3	2.52	0.44
1:E:12(B):VAL:HG13	1:F:12(D):VAL:HG23	1.99	0.43
1:E:27(B):THR:HA	1:E:28(B):PRO:HD2	1.81	0.42
1:F:11(D):LEU:HD21	1:F:2(C):ILE:CG1	2.41	0.42
1:E:28(B):PRO:HG3	1:F:21(D):GLU:O	2.20	0.41
1:E:7(B):CYS:O	1:E:11(B):LEU:HG	2.21	0.41

All (15) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:1(D):PHE:N	3:E:40:HOH:O[3_555]	1.12	1.08
3:E:73:HOH:O	3:F:55:HOH:O[6_455]	1.17	1.03
3:F:43:HOH:O	3:F:58:HOH:O[9_555]	1.73	0.47
3:E:65:HOH:O	3:F:77:HOH:O[1_556]	1.77	0.43
3:E:48:HOH:O	3:E:118:HOH:O[6_455]	1.79	0.41
3:E:71:HOH:O	3:F:74:HOH:O[6_455]	1.81	0.39
1:E:15(A):GLN:OE1	3:F:101:HOH:O[6_455]	1.84	0.36
3:E:104:HOH:O	3:E:105:HOH:O[2_555]	1.87	0.33
3:E:127:HOH:O	3:F:68:HOH:O[2_555]	1.90	0.30
3:E:126:HOH:O	3:F:40:HOH:O[1_556]	2.00	0.20
3:E:57:HOH:O	3:F:98:HOH:O[6_455]	2.00	0.20
3:E:49:HOH:O	3:E:66:HOH:O[1_556]	2.02	0.18
3:E:65:HOH:O	3:F:63:HOH:O[1_556]	2.03	0.17
1:F:4(C):GLU:OE2	3:E:95:HOH:O[1_554]	2.09	0.11
3:E:96:HOH:O	3:F:40:HOH:O[1_556]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	48/50 (96%)	41 (85%)	4 (8%)	3 (6%)	2	0
1	F	48/50 (96%)	37 (77%)	5 (10%)	6 (12%)	0	0
All	All	96/100 (96%)	78 (81%)	9 (9%)	9 (9%)	1	0

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	2(B)	VAL
1	E	28(B)	PRO
1	F	29(D)	LYS
1	F	4(C)	GLU
1	F	13(C)	LEU
1	E	3(B)	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	28(D)	PRO
1	F	20(D)	GLY
1	F	3(C)	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	44/45 (98%)	36 (82%)	8 (18%)	2	1
1	F	43/45 (96%)	37 (86%)	6 (14%)	4	2
All	All	87/90 (97%)	73 (84%)	14 (16%)	3	1

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	1(B)	PHE
1	E	12(B)	VAL
1	E	2(A)	ILE
1	E	3(A)	VAL
1	E	5(A)	GLN
1	E	8(A)	THR
1	E	9(A)	SER
1	E	13(A)	LEU
1	F	4(D)	GLN
1	F	6(D)	LEU
1	F	15(D)	LEU
1	F	17(D)	LEU
1	F	2(C)	ILE
1	F	5(C)	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	3(B)	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	18(A)	ASN
1	F	15(C)	GLN
1	F	18(C)	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	50/50 (100%)	-0.60	0 100 100	7, 23, 35, 45	0
1	F	50/50 (100%)	-0.62	0 100 100	12, 21, 40, 46	3 (6%)
All	All	100/100 (100%)	-0.61	0 100 100	7, 22, 40, 46	3 (3%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ZN	F	30	1/1	1.00	0.03	-	21,21,21,21	1
2	ZN	E	30	1/1	1.00	0.02	-	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.