



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:32 AM GMT

PDB ID : 3ITG
Title : Structure the proline utilization A proline dehydrogenase domain (PutA86-630) inactivated with N-propargylglycine
Authors : Tanner, J.J.
Deposited on : 2009-08-28
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	602	<div> <div></div> <div>58%8%34%</div> </div>
1	B	602	<div> <div></div> <div>58%8%33%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional protein putA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	399	Total	C	N	O	S	0	1	0
			3060	1944	526	579	11			
1	B	401	Total	C	N	O	S	0	1	0
			3073	1954	529	579	11			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	670	SER	-	EXPRESSION TAG	UNP P09546
A	671	SER	-	EXPRESSION TAG	UNP P09546
A	672	SER	-	EXPRESSION TAG	UNP P09546
A	673	VAL	-	EXPRESSION TAG	UNP P09546
A	674	ASP	-	EXPRESSION TAG	UNP P09546
A	675	LYS	-	EXPRESSION TAG	UNP P09546
A	676	LEU	-	EXPRESSION TAG	UNP P09546
A	677	ALA	-	EXPRESSION TAG	UNP P09546
A	678	ALA	-	EXPRESSION TAG	UNP P09546
A	679	ALA	-	EXPRESSION TAG	UNP P09546
A	680	LEU	-	EXPRESSION TAG	UNP P09546
A	681	GLU	-	EXPRESSION TAG	UNP P09546
A	682	HIS	-	EXPRESSION TAG	UNP P09546
A	683	HIS	-	EXPRESSION TAG	UNP P09546
A	684	HIS	-	EXPRESSION TAG	UNP P09546
A	685	HIS	-	EXPRESSION TAG	UNP P09546
A	686	HIS	-	EXPRESSION TAG	UNP P09546
A	687	HIS	-	EXPRESSION TAG	UNP P09546
B	670	SER	-	EXPRESSION TAG	UNP P09546
B	671	SER	-	EXPRESSION TAG	UNP P09546
B	672	SER	-	EXPRESSION TAG	UNP P09546
B	673	VAL	-	EXPRESSION TAG	UNP P09546
B	674	ASP	-	EXPRESSION TAG	UNP P09546
B	675	LYS	-	EXPRESSION TAG	UNP P09546
B	676	LEU	-	EXPRESSION TAG	UNP P09546

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	677	ALA	-	EXPRESSION TAG	UNP P09546
B	678	ALA	-	EXPRESSION TAG	UNP P09546
B	679	ALA	-	EXPRESSION TAG	UNP P09546
B	680	LEU	-	EXPRESSION TAG	UNP P09546
B	681	GLU	-	EXPRESSION TAG	UNP P09546
B	682	HIS	-	EXPRESSION TAG	UNP P09546
B	683	HIS	-	EXPRESSION TAG	UNP P09546
B	684	HIS	-	EXPRESSION TAG	UNP P09546
B	685	HIS	-	EXPRESSION TAG	UNP P09546
B	686	HIS	-	EXPRESSION TAG	UNP P09546
B	687	HIS	-	EXPRESSION TAG	UNP P09546

- Molecule 2 is DIHYDROFLAVINE-ADENINE DINUCLEOTIDE (three-letter code: FDA) (formula: $C_{27}H_{35}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	124	Total	O	0	0
			124	124		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	117	Total	O	0	0
			117	117		

ASN	SER	ALA	GLY	LEU	ASP	LEU	ALA	ASN	GLU	ARG	HIS	ARG	LEU	ALA	SER	SER	SER	SER	ALA	LEU	GLN	LYS	TRP	ALA	ALA	LEU	PRO	PRO	LEU	LEU	GLU	GLN	GLU	PRO	PRO	VAL	ALA	ALA	ALA	GLY	GLY	GLU	GLU	SER	MET	ASN	ILE	VAL	VAL	PRO	PRO	GLU	GLU	ASP	LYS	LEU	ALA
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

ALA
ALA
LEU
GLU
HIS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	127.48Å 133.36Å 133.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.75 – 2.15 31.41 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.4 (30.75-2.15) 96.9 (31.41-2.15)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.16Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.193 , 0.234 0.193 , 0.234	Depositor DCC
R_{free} test set	3048 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.236	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.2	EDS
Estimated twinning fraction	0.467 for -h,-l,-k 0.015 for l,-k,h 0.016 for -k,-h,l 0.012 for k,-l,-h 0.012 for -l,h,-k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 60416 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6480	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FDA, LYX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/3109	0.47	0/4231
1	B	0.33	0/3122	0.48	0/4246
All	All	0.33	0/6231	0.48	0/8477

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3060	0	2965	25	0
1	B	3073	0	2985	34	0
2	A	53	0	32	3	0
2	B	53	0	32	3	0
3	A	124	0	0	0	0
3	B	117	0	0	0	0
All	All	6480	0	6014	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:540:TYR:HB2	2:B:2002:FDA:HM72	1.69	0.74
1:A:540:TYR:HB2	2:A:2001:FDA:HM72	1.69	0.73
1:B:485:ALA:HB1	2:B:2002:FDA:HM83	1.69	0.71
1:A:135:LEU:HD23	1:A:570:ILE:HD13	1.74	0.68
1:A:485:ALA:HB1	2:A:2001:FDA:HM83	1.79	0.65
1:A:518:GLU:O	1:A:522:GLU:HG2	1.99	0.62
1:B:135:LEU:HD23	1:B:570:ILE:HD13	1.84	0.60
1:A:263:GLU:OE2	1:A:546:HIS:HD2	1.86	0.58
1:B:492:LEU:HD11	1:B:524:VAL:HG21	1.84	0.58
1:A:307:ALA:O	1:A:311:ILE:HG13	2.06	0.56
1:B:485:ALA:HB1	2:B:2002:FDA:C8M	2.34	0.56
1:B:518:GLU:O	1:B:522:GLU:HG2	2.07	0.55
1:A:516:MET:HE3	1:A:567:VAL:HG11	1.91	0.52
1:B:547:GLU:C	1:B:549:LEU:H	2.14	0.51
1:B:92:ARG:NH1	1:B:382:ASP:OD2	2.42	0.51
1:A:492:LEU:HD11	1:A:524:VAL:HG21	1.94	0.50
1:A:92:ARG:NH2	1:A:382:ASP:OD2	2.42	0.50
1:B:572:ASP:OD1	1:B:574:SER:HB3	2.13	0.49
1:B:288:GLY:HA2	1:B:548:THR:HG21	1.94	0.49
1:B:120:VAL:HG13	1:B:497:GLN:HB3	1.95	0.48
1:A:350:TYR:HB3	1:A:351:PRO:HD3	1.96	0.48
1:A:575:LEU:HD12	1:A:576:PRO:HD2	1.96	0.48
1:B:433:VAL:HG12	1:B:485:ALA:HB3	1.95	0.47
1:B:326:ILE:HG23	1:B:367:ILE:HG13	1.96	0.47
1:B:350:TYR:HB3	1:B:351:PRO:HD3	1.96	0.46
1:A:547:GLU:C	1:A:549:LEU:H	2.19	0.46
1:A:545:THR:HB	1:A:547:GLU:OE1	2.16	0.46
1:B:287:LEU:HD12	1:B:287:LEU:HA	1.77	0.46
1:A:410:CYS:HB3	1:A:411:PRO:HD3	1.97	0.45
1:A:548:THR:HG22	1:A:548:THR:O	2.15	0.45
1:A:485:ALA:HB1	2:A:2001:FDA:C8M	2.46	0.45
1:B:263:GLU:OE2	1:B:546:HIS:HD2	2.00	0.45
1:B:475:ALA:C	1:B:477:PRO:HD3	2.37	0.45
1:B:326:ILE:HG13	1:B:327:SER:N	2.33	0.44
1:A:361:ARG:HG3	1:A:396:TRP:CG	2.52	0.44
1:A:402:VAL:HG22	1:A:431:ARG:HD2	1.98	0.44
1:B:400:GLY:HA2	1:B:429:MET:O	2.18	0.44
1:A:275:LEU:HD23	1:A:278:LYS:HD3	2.00	0.43
1:B:569:ARG:NH2	1:B:580:LEU:HD12	2.33	0.43
1:B:547:GLU:C	1:B:549:LEU:N	2.71	0.43
1:B:567:VAL:O	1:B:570:ILE:HG22	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:475:ALA:C	1:A:477:PRO:HD3	2.39	0.43
1:B:288:GLY:HA2	1:B:548:THR:CG2	2.49	0.43
1:B:120:VAL:HG13	1:B:497:GLN:CB	2.50	0.42
1:B:396:TRP:CZ2	1:B:398:GLY:HA3	2.54	0.42
1:B:547:GLU:O	1:B:549:LEU:N	2.52	0.42
1:B:275:LEU:HD23	1:B:278:LYS:HD3	2.01	0.42
1:A:100:ARG:CZ	1:A:450:LEU:HD13	2.50	0.42
1:A:565:SER:O	1:A:569:ARG:HG3	2.20	0.42
1:B:342:TYR:CD1	1:B:379:ILE:HD13	2.55	0.41
1:A:261:THR:OG1	1:A:541:ALA:HB1	2.19	0.41
1:B:492:LEU:C	1:B:492:LEU:HD23	2.40	0.41
1:B:342:TYR:HD1	1:B:379:ILE:HD13	1.86	0.41
1:A:326:ILE:HG13	1:A:327:SER:N	2.36	0.41
1:A:126:LYS:O	1:A:130:GLN:HG2	2.21	0.41
1:B:402:VAL:HG22	1:B:431:ARG:HB3	2.03	0.41
1:B:307:ALA:O	1:B:311:ILE:HG13	2.21	0.41
1:B:535:ARG:HA	1:B:536:PRO:HD3	1.97	0.41
1:B:592:ALA:HB2	1:B:598:THR:HA	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/602 (65%)	385 (98%)	7 (2%)	1 (0%)	46	42
1	B	395/602 (66%)	386 (98%)	8 (2%)	1 (0%)	46	42
All	All	788/1204 (65%)	771 (98%)	15 (2%)	2 (0%)	46	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	548	THR
1	A	548	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/489 (62%)	296 (98%)	7 (2%)	58	62
1	B	303/489 (62%)	295 (97%)	8 (3%)	54	55
All	All	606/978 (62%)	591 (98%)	15 (2%)	55	58

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	TYR
1	A	327	SER
1	A	349	LEU
1	A	406	TYR
1	A	440	SER
1	A	578	ASP
1	A	593	GLN
1	B	99	TYR
1	B	287	LEU
1	B	327	SER
1	B	349	LEU
1	B	361	ARG
1	B	406	TYR
1	B	440	SER
1	B	593	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	271	ASN
1	A	305	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	316	ASN
1	A	362	GLN
1	A	497	GLN
1	A	534	ASN
1	A	546	HIS
1	A	594	GLN
1	B	271	ASN
1	B	305	GLN
1	B	316	ASN
1	B	368	ASN
1	B	497	GLN
1	B	501	GLN
1	B	534	ASN
1	B	546	HIS
1	B	594	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LYX	A	329	1,2	10,11,64	2.04	1 (10%)	8,11,93	1.17	1 (12%)
1	LYX	B	329	1,2	10,11,64	2.01	1 (10%)	8,11,93	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LYX	A	329	1,2	-	0/8/10/83	0/0/0/3
1	LYX	B	329	1,2	-	0/8/10/83	0/0/0/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	329	LYX	C23-NZ	-5.96	1.27	1.46
1	B	329	LYX	C23-NZ	-5.87	1.28	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	329	LYX	CE-NZ-C23	2.09	120.75	113.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FDA	A	2001	1	48,58,58	0.99	1 (2%)	54,89,89	2.34	10 (18%)
2	FDA	B	2002	1	48,58,58	1.00	2 (4%)	54,89,89	2.27	11 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FDA	A	2001	1	-	0/30/50/50	0/6/6/6
2	FDA	B	2002	1	-	0/30/50/50	0/6/6/6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2002	FDA	C4X-N5	2.13	1.36	1.33
2	A	2001	FDA	C4-C4X	3.90	1.49	1.41
2	B	2002	FDA	C4-C4X	4.10	1.49	1.41

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	FDA	N3A-C2A-N1A	-10.60	120.78	128.89
2	B	2002	FDA	N3A-C2A-N1A	-10.39	120.94	128.89
2	A	2001	FDA	C4X-C10-N10	-5.20	117.45	120.52
2	B	2002	FDA	C4-C4X-C10	-4.79	116.87	119.94
2	A	2001	FDA	C4-C4X-C10	-4.53	117.04	119.94
2	B	2002	FDA	C4X-C10-N10	-4.37	117.94	120.52
2	A	2001	FDA	C4X-C4-N3	-4.11	117.97	123.59
2	B	2002	FDA	C4X-C4-N3	-4.04	118.06	123.59
2	B	2002	FDA	C9A-C5X-N5	-2.61	118.50	122.36
2	A	2001	FDA	C9A-C5X-N5	-2.48	118.68	122.36
2	A	2001	FDA	C4X-N5-C5X	-2.18	114.26	116.76
2	B	2002	FDA	C4A-C5A-N7A	-2.08	107.57	109.48
2	B	2002	FDA	C5X-C9A-N10	2.11	119.22	117.62
2	A	2001	FDA	C6-C5X-N5	2.14	121.72	118.96
2	B	2002	FDA	C1'-N10-C9A	2.16	121.28	118.86
2	B	2002	FDA	C6-C5X-N5	2.34	121.97	118.96
2	A	2001	FDA	C5X-C9A-N10	2.72	119.68	117.62
2	A	2001	FDA	C4-C4X-N5	3.50	122.96	118.72
2	B	2002	FDA	C4-C4X-N5	3.72	123.24	118.72
2	B	2002	FDA	C4-N3-C2	6.82	121.14	115.25
2	A	2001	FDA	C4-N3-C2	7.18	121.45	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2001	FDA	3	0
2	B	2002	FDA	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	398/602 (66%)	-0.06	4 (1%) 84 88	16, 30, 68, 102	0
1	B	400/602 (66%)	-0.07	4 (1%) 84 88	15, 30, 70, 106	0
All	All	798/1204 (66%)	-0.07	8 (1%) 84 88	15, 30, 69, 106	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	549	LEU	5.2
1	B	549	LEU	4.2
1	B	553	LEU	2.8
1	A	575	LEU	2.7
1	B	567	VAL	2.6
1	B	256	GLY	2.4
1	A	550	LEU	2.2
1	A	553	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	LYX	A	329	12/62	0.97	0.13	-	20,24,34,35	0
1	LYX	B	329	12/62	0.98	0.12	-	18,25,35,38	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FDA	B	2002	53/53	0.98	0.12	0.11	17,25,34,37	0
2	FDA	A	2001	53/53	0.98	0.11	-0.09	17,25,34,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.