



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:33 AM GMT

PDB ID : 3IWJ
Title : Crystal structure of aminoaldehyde dehydrogenase 2 from Pisum sativum (PsAMADH2)
Authors : Kopecny, D.; Morera, S.; Briozzo, P.
Deposited on : 2009-09-02
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	 6% 82% 17% ..
1	B	503	 6% 82% 16% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	507	-	-	-	X
4	GOL	B	506	-	-	-	X
4	GOL	B	508	-	-	-	X
4	GOL	B	509	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative aminoaldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	500	Total	C	N	O	S	0	0	0
			3811	2445	636	713	17			
1	B	500	Total	C	N	O	S	0	1	0
			3817	2450	637	713	17			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Na 1 1	0	0
3	A	1	Total Na 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	140	Total O 140 140	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	150	Total	O	0	0
			150	150		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative aminoaldehyde dehydrogenase



- Molecule 1: Putative aminoaldehyde dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.25Å 86.73Å 179.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.15 24.32 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (25.00-2.15) 99.3 (24.32-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 2.15Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.234 , 0.289 0.238 , 0.295	Depositor DCC
R_{free} test set	2827 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.56$, $\langle L^2 \rangle = 0.40$	Xtriage
Outliers	1 of 56542 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8016	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.13 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.7922e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAD, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3899	0.56	0/5311
1	B	0.40	0/3908	0.55	0/5322
All	All	0.40	0/7807	0.55	0/10633

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3811	0	3852	53	0
1	B	3817	0	3865	47	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	18	0	24	0	0
4	B	24	0	32	2	0
5	A	140	0	0	1	0
5	B	150	0	0	0	0
All	All	8016	0	7797	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:33:GLN:HE21	1:A:372:GLY:H	1.04	0.96
1:B:33:GLN:HE21	1:B:372:GLY:H	1.18	0.86
1:B:8:ARG:O	1:B:24:ARG:NH2	2.10	0.84
1:A:157:LEU:HG	1:A:171:LYS:HB3	1.77	0.67
1:A:425:VAL:HG22	1:A:440:PHE:CE2	2.29	0.65
1:A:163:TYR:HB2	1:A:167:MET:HG2	1.79	0.65
1:A:425:VAL:HG22	1:A:440:PHE:HE2	1.63	0.64
1:B:9:GLN:HA	1:B:9:GLN:HE21	1.62	0.64
1:B:459:TRP:CZ2	4:B:508:GOL:H2	2.33	0.63
1:B:384:THR:HG22	1:B:387:MET:HG3	1.79	0.63
1:A:8:ARG:O	1:A:24:ARG:NH2	2.23	0.62
1:A:292:GLN:HE22	1:A:337:VAL:H	1.45	0.62
1:A:105:ASP:OD2	1:A:332:ARG:NH2	2.33	0.60
1:A:33:GLN:HE21	1:A:372:GLY:N	1.88	0.60
1:A:157:LEU:HD22	1:A:182:ALA:HB1	1.84	0.59
1:A:144:LYS:HE2	5:A:578:HOH:O	2.03	0.59
1:B:9:GLN:HA	1:B:9:GLN:NE2	2.18	0.58
1:B:425:VAL:HG13	1:B:440:PHE:CE2	2.37	0.58
1:A:384:THR:HG22	1:A:387:MET:HG3	1.86	0.58
1:A:502:LYS:O	1:A:503:LEU:HB2	2.02	0.58
1:B:15:ASP:OD1	1:B:202:LYS:HE2	2.04	0.58
1:A:13:ASN:H	1:A:212:ASN:ND2	2.02	0.57
1:A:285:GLY:HA2	1:A:451:GLN:HG2	1.86	0.57
1:B:323:ILE:HD12	1:B:333:LEU:HD23	1.85	0.57
1:B:292:GLN:HE22	1:B:337:VAL:H	1.53	0.56
1:A:288:TRP:CE2	1:A:289:THR:HG22	2.40	0.56
1:B:432:ARG:O	1:B:436:VAL:HG23	2.06	0.55
1:A:284:PHE:O	1:A:288:TRP:HB3	2.07	0.55
1:A:425:VAL:CG2	1:A:440:PHE:HE2	2.21	0.54
1:A:293:ILE:HB	1:A:296:ALA:HB2	1.90	0.54
1:B:293:ILE:HB	1:B:296:ALA:HB2	1.89	0.53
1:B:311:LEU:HA	1:B:314:ILE:HG22	1.91	0.53
1:B:13:ASN:H	1:B:212:ASN:ND2	2.06	0.53
1:A:171:LYS:O	1:A:174:PRO:HD2	2.08	0.52
1:A:15:ASP:OD1	1:A:202:LYS:HE2	2.08	0.52
1:A:311:LEU:HA	1:A:314:ILE:HG22	1.92	0.52
1:A:152:ILE:HD13	1:A:233:LYS:HD2	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:TYR:HB2	1:B:167:MET:HG2	1.92	0.52
1:B:450:SER:C	1:B:452:PRO:HD3	2.31	0.52
1:A:35:ILE:HD12	1:A:35:ILE:N	2.26	0.51
1:A:13:ASN:H	1:A:212:ASN:HD22	1.58	0.50
1:A:271:GLU:HB3	1:A:303:HIS:ND1	2.27	0.50
1:B:157:LEU:HG	1:B:171:LYS:HB3	1.93	0.50
1:A:39:ILE:HB	1:A:215:THR:HG21	1.94	0.50
1:B:425:VAL:HG13	1:B:440:PHE:CZ	2.47	0.49
1:B:283:ILE:HG23	1:B:287:PHE:HD1	1.77	0.49
1:B:383:VAL:HG21	1:B:400:CYS:HB2	1.95	0.49
1:A:273:VAL:HG21	1:A:426:ILE:HG22	1.95	0.49
1:A:273:VAL:HG21	1:A:426:ILE:CG2	2.43	0.49
1:B:33:GLN:HE21	1:B:372:GLY:N	1.97	0.48
1:A:342:TYR:HB2	1:A:373:PHE:HB3	1.96	0.48
1:B:105:ASP:OD2	1:B:332:ARG:NH2	2.46	0.48
1:A:323:ILE:HD12	1:A:333:LEU:HD23	1.96	0.48
1:B:71:GLY:HA3	1:B:125:ALA:O	2.14	0.47
1:A:71:GLY:HA3	1:A:125:ALA:O	2.14	0.47
1:A:286:CYS:O	1:A:291:GLY:HA2	2.14	0.47
1:A:9:GLN:HA	1:A:9:GLN:HE21	1.79	0.47
1:B:311:LEU:O	1:B:315:VAL:HG23	2.14	0.47
1:B:157:LEU:HD22	1:B:182:ALA:HB1	1.97	0.47
1:A:9:GLN:HA	1:A:9:GLN:NE2	2.30	0.46
1:A:450:SER:C	1:A:452:PRO:HD3	2.36	0.46
1:B:247:MET:HG3	1:B:257:VAL:HG11	1.98	0.45
1:A:163:TYR:OH	1:A:289:THR:HG21	2.16	0.45
1:A:228:HIS:HB3	1:A:231:VAL:HG23	1.97	0.45
1:A:450:SER:O	1:A:451:GLN:HB2	2.17	0.45
1:A:40:PRO:O	1:A:215:THR:HG22	2.15	0.45
1:B:91:GLU:CD	1:B:91:GLU:H	2.19	0.45
1:A:323:ILE:HD11	1:A:374:PHE:HB3	1.98	0.45
1:A:51:VAL:HG11	1:A:228:HIS:CE1	2.51	0.45
1:A:383:VAL:HG21	1:A:400:CYS:HB2	1.99	0.44
1:A:288:TRP:CD2	1:A:289:THR:HG22	2.52	0.44
1:B:270:PHE:HB2	1:B:427:SER:HB2	1.99	0.44
1:B:201:CYS:SG	1:B:211:LEU:HD23	2.58	0.44
1:B:285:GLY:HA2	1:B:451:GLN:HG2	2.00	0.44
1:B:342:TYR:HB2	1:B:373:PHE:HB3	1.98	0.44
1:B:323:ILE:HD11	1:B:374:PHE:HB3	2.00	0.44
1:A:332:ARG:NH1	1:B:496:TRP:O	2.47	0.43
1:A:39:ILE:HD12	1:A:215:THR:CG2	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:TRP:CD2	1:A:471:LEU:HD21	2.54	0.43
1:B:250:ALA:HB1	1:B:255:LYS:HB3	1.99	0.42
1:B:423:ALA:O	1:B:445:VAL:HA	2.19	0.42
1:B:194:CYS:O	1:B:197:LEU:HB3	2.20	0.42
1:B:300:LEU:HD11	1:B:310:PHE:CE2	2.54	0.42
1:B:450:SER:O	1:B:451:GLN:HB2	2.20	0.42
1:B:124:LEU:HB3	1:B:476:LEU:HD22	2.02	0.42
1:B:391:ARG:HB2	1:B:391:ARG:HE	1.75	0.42
1:B:163:TYR:CE2	4:B:508:GOL:H11	2.54	0.41
1:A:443:GLY:HA3	1:A:460:GLY:O	2.20	0.41
1:B:418:VAL:O	1:B:463:LYS:HB3	2.21	0.41
1:A:237:THR:HA	1:A:260:GLU:O	2.21	0.41
1:A:161:TRP:CD1	1:A:189:LEU:HD12	2.55	0.41
1:A:133:LYS:HB3	1:A:133:LYS:HE2	1.82	0.41
1:A:124:LEU:HB3	1:A:476:LEU:HD22	2.03	0.41
1:B:425:VAL:HG11	1:B:436:VAL:CG1	2.50	0.41
1:B:159:THR:HB	1:B:160:PRO:HD2	2.02	0.41
1:A:418:VAL:O	1:A:464:ARG:HB2	2.21	0.40
1:B:121:TYR:CE2	1:B:174:PRO:HB3	2.57	0.40
1:B:284:PHE:O	1:B:288:TRP:HB3	2.21	0.40
1:B:268:VAL:HB	1:B:425:VAL:HG12	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	498/503 (99%)	478 (96%)	18 (4%)	2 (0%)	39	34
1	B	499/503 (99%)	479 (96%)	19 (4%)	1 (0%)	52	51
All	All	997/1006 (99%)	957 (96%)	37 (4%)	3 (0%)	46	42

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	451	GLN
1	B	451	GLN
1	A	22	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	405/408 (99%)	385 (95%)	20 (5%)	31	26
1	B	406/408 (100%)	388 (96%)	18 (4%)	35	31
All	All	811/816 (99%)	773 (95%)	38 (5%)	32	28

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ASN
1	A	91	GLU
1	A	96	GLU
1	A	97	SER
1	A	138	LEU
1	A	144	LYS
1	A	157	LEU
1	A	185	LYS
1	A	217	LEU
1	A	225	LEU
1	A	254	VAL
1	A	289	THR
1	A	300	LEU
1	A	323	ILE
1	A	353	LYS
1	A	360	LEU
1	A	384	THR
1	A	395	PHE
1	A	430	LEU
1	A	492	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	29	ASN
1	B	55	LYS
1	B	91	GLU
1	B	96	GLU
1	B	138	LEU
1	B	157	LEU
1	B	185	LYS
1	B	217	LEU
1	B	225	LEU
1	B	275	LEU
1	B	300	LEU
1	B	323	ILE
1	B	344	LYS
1	B	360	LEU
1	B	384	THR
1	B	385	THR
1	B	395	PHE
1	B	492	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	29	ASN
1	A	33	GLN
1	A	212	ASN
1	A	292	GLN
1	A	478	ASN
1	B	9	GLN
1	B	29	ASN
1	B	33	GLN
1	B	212	ASN
1	B	292	GLN
1	B	312	ASN
1	B	478	ASN
1	B	484	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	A	504	-	23,29,48	1.05	1 (4%)	27,45,73	1.61	3 (11%)
4	GOL	A	506	-	5,5,5	0.36	0	5,5,5	0.60	0
4	GOL	A	507	-	5,5,5	0.38	0	5,5,5	0.26	0
4	GOL	A	508	-	5,5,5	0.35	0	5,5,5	0.33	0
2	NAD	B	504	-	23,29,48	0.99	1 (4%)	27,45,73	1.78	4 (14%)
4	GOL	B	506	-	5,5,5	0.35	0	5,5,5	0.22	0
4	GOL	B	507	-	5,5,5	0.39	0	5,5,5	0.17	0
4	GOL	B	508	-	5,5,5	0.38	0	5,5,5	0.38	0
4	GOL	B	509	-	5,5,5	0.35	0	5,5,5	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	504	-	-	0/12/32/62	0/3/3/5
4	GOL	A	506	-	-	0/4/4/4	0/0/0/0
4	GOL	A	507	-	-	0/4/4/4	0/0/0/0
4	GOL	A	508	-	-	0/4/4/4	0/0/0/0
2	NAD	B	504	-	-	0/12/32/62	0/3/3/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	506	-	-	0/4/4/4	0/0/0/0
4	GOL	B	507	-	-	0/4/4/4	0/0/0/0
4	GOL	B	508	-	-	0/4/4/4	0/0/0/0
4	GOL	B	509	-	-	0/4/4/4	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	504	NAD	C5A-C4A	3.23	1.47	1.40
2	A	504	NAD	C5A-C4A	3.62	1.48	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	504	NAD	N3A-C2A-N1A	-7.09	123.46	128.89
2	A	504	NAD	N3A-C2A-N1A	-6.37	124.02	128.89
2	B	504	NAD	C4A-C5A-N7A	-2.95	106.77	109.48
2	A	504	NAD	C4A-C5A-N7A	-2.90	106.81	109.48
2	B	504	NAD	C2B-C1B-N9A	-2.32	110.75	114.29
2	A	504	NAD	O4B-C1B-N9A	2.36	113.03	108.10
2	B	504	NAD	O4B-C1B-N9A	2.88	114.13	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	508	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	500/503 (99%)	0.38	30 (6%) 25 34	26, 38, 50, 55	0
1	B	500/503 (99%)	0.39	28 (5%) 28 38	25, 38, 51, 59	0
All	All	1000/1006 (99%)	0.39	58 (5%) 26 36	25, 38, 51, 59	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	367	GLU	4.7
1	A	367	GLU	4.1
1	B	267	LEU	3.9
1	B	399	LEU	3.8
1	B	424	ALA	3.6
1	A	503	LEU	3.4
1	B	211	LEU	3.2
1	A	370	LYS	3.2
1	B	141	ASP	3.1
1	A	267	LEU	3.0
1	B	286	CYS	3.0
1	B	273	VAL	3.0
1	A	141	ASP	2.9
1	B	353	LYS	2.9
1	A	329	GLU	2.9
1	A	211	LEU	2.8
1	A	399	LEU	2.8
1	A	391	ARG	2.7
1	A	286	CYS	2.7
1	A	424	ALA	2.7
1	B	368	HIS	2.5
1	B	213	ILE	2.5
1	A	61	ASN	2.4
1	A	285	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	498	GLN	2.4
1	B	425	VAL	2.4
1	A	4	PRO	2.4
1	B	333	LEU	2.4
1	B	502	LYS	2.4
1	B	4	PRO	2.3
1	A	308	THR	2.3
1	B	334	GLY	2.3
1	B	397	PRO	2.3
1	B	398	VAL	2.3
1	B	287	PHE	2.2
1	B	386	ASN	2.2
1	A	184	LEU	2.2
1	B	175	ALA	2.2
1	B	370	LYS	2.2
1	A	498	GLN	2.2
1	B	503	LEU	2.2
1	B	87	GLU	2.2
1	A	175	ALA	2.2
1	A	323	ILE	2.2
1	B	360	LEU	2.1
1	A	273	VAL	2.1
1	A	155	VAL	2.1
1	A	500	PRO	2.1
1	A	353	LYS	2.1
1	A	301	ILE	2.1
1	A	491	GLU	2.1
1	A	312	ASN	2.1
1	A	297	THR	2.1
1	A	282	ALA	2.0
1	A	450	SER	2.0
1	B	312	ASN	2.0
1	A	172	VAL	2.0
1	B	172	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GOL	B	508	6/6	0.77	0.24	4.58	60,60,61,61	0
4	GOL	B	506	6/6	0.86	0.19	2.76	38,41,42,42	0
4	GOL	A	507	6/6	0.79	0.20	2.22	59,60,60,60	0
4	GOL	B	509	6/6	0.72	0.23	2.06	83,84,84,84	0
4	GOL	A	506	6/6	0.84	0.18	1.25	35,36,37,37	0
2	NAD	B	504	27/44	0.89	0.12	-0.07	26,33,51,52	0
3	NA	A	505	1/1	0.87	0.14	-0.16	35,35,35,35	0
2	NAD	A	504	27/44	0.91	0.10	-0.49	27,32,52,53	0
3	NA	B	505	1/1	0.85	0.11	-2.51	43,43,43,43	0
4	GOL	B	507	6/6	0.72	0.18	-	66,67,67,67	0
4	GOL	A	508	6/6	0.80	0.17	-	68,69,69,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.