



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:37 PM GMT

PDB ID : 4IYN  
Title : Structure of mitochondrial Hsp90 (TRAP1) with ADP-ALF4-  
Authors : Partridge, J.R.; Lavery, L.A.; Agard, D.A.  
Deposited on : 2013-01-28  
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	719	<div> <div>7%</div> <div> <div></div> <div>73%</div> <div>10%</div> <div>17%</div> </div> </div>
1	B	719	<div> <div>3%</div> <div> <div></div> <div>75%</div> <div>9%</div> <div>15%</div> </div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TNF receptor-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	596	Total	C	N	O	S	0	2	0
			4789	3032	831	904	22			
1	B	609	Total	C	N	O	S	1	6	0
			4954	3140	858	933	23			

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Co	0	0
			2	2		
2	A	2	Total	Co	0	0
			2	2		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is TETRAFLUOROALUMINATE ION (three-letter code: ALF) (formula:  $\text{AlF}_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Al	F	0	0
			5	1	4		
4	B	1	Total	Al	F	0	0
			5	1	4		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula:  $\text{Mg}$ ).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	125	Total	O	0	0
			125	125		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	152	Total 152	O 152	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TNF receptor-associated protein 1



- Molecule 1: TNF receptor-associated protein 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.90 Å 96.60 Å 125.90 Å 90.00° 134.26° 90.00°	Depositor
Resolution (Å)	29.81 – 2.31 49.27 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.81-2.31) 94.3 (49.27-2.31)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.20 (at 2.32 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.191 , 0.231 0.188 , 0.228	Depositor DCC
$R_{free}$ test set	1889 reflections (2.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	48.8	Xtriage
Anisotropy	0.493	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 45.6	EDS
Estimated twinning fraction	0.000 for h+2*l,k,-h-l 0.007 for h,-k,-h-l 0.014 for -h-2*l,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 66761 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10090	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ALF, MG, CO, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/4881	0.57	0/6578
1	B	0.43	0/5056	0.57	0/6811
All	All	0.43	0/9937	0.57	0/13389

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4789	0	4804	55	0
1	B	4954	0	4958	49	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	27	0	12	1	0
3	B	27	0	12	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	125	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	152	0	0	8	0
All	All	10090	0	9786	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:456:PHE:CD1	1:B:463:MET:HE1	2.07	0.89
1:B:456:PHE:HD1	1:B:463:MET:HE1	1.42	0.81
1:B:463:MET:SD	1:B:482:LEU:CB	2.73	0.77
1:A:490:LEU:HD11	1:A:496:THR:HG21	1.68	0.76
1:B:463:MET:SD	1:B:482:LEU:HB2	2.26	0.75
1:A:611:ASN:HB3	1:A:656:THR:HG23	1.71	0.72
1:B:637:LEU:O	1:B:638:ARG:NE	2.24	0.71
1:A:598:LEU:HB3	1:A:602:ARG:HH21	1.57	0.70
1:B:463:MET:SD	1:B:482:LEU:HB3	2.32	0.69
1:B:412:PRO:HG2	1:B:421:GLN:HB2	1.75	0.68
1:B:456:PHE:CD1	1:B:463:MET:CE	2.77	0.67
1:A:625:ILE:HD13	1:A:657:LEU:HB3	1.85	0.59
1:A:606:VAL:HG23	1:A:607:GLN:HG2	1.86	0.58
1:A:116:VAL:CG1	1:A:218:VAL:HB	2.35	0.57
1:B:456:PHE:HD1	1:B:463:MET:CE	2.13	0.57
1:A:490:LEU:CD1	1:A:496:THR:CG2	2.83	0.56
1:A:490:LEU:CD1	1:A:496:THR:HG21	2.36	0.56
1:A:676:ASN:HB3	1:A:679:LEU:HB3	1.88	0.55
1:B:366:SER:HB3	1:B:369:ASP:HB2	1.88	0.55
1:B:179:ASN:OD1	1:B:182:ASP:N	2.39	0.55
1:A:612:ILE:HG23	1:A:657:LEU:HD11	1.89	0.55
1:B:456:PHE:CE1	1:B:463:MET:HE1	2.41	0.54
1:B:456:PHE:CE1	1:B:463:MET:CE	2.90	0.54
1:A:126:VAL:HG13	1:A:129:ARG:CZ	2.37	0.54
1:A:123:GLU:O	1:A:126:VAL:HG23	2.08	0.54
1:B:435[A]:GLN:HG3	6:B:908:HOH:O	2.06	0.54
1:B:232:VAL:HB	1:B:246:TRP:HB3	1.91	0.53
1:A:490:LEU:HD11	1:A:496:THR:CG2	2.38	0.53
1:A:612:ILE:HG23	1:A:657:LEU:CD1	2.39	0.53
1:A:330:TYR:OH	1:A:361:PRO:HB3	2.09	0.53
1:A:518:PRO:HD2	1:A:630:MET:HE3	1.91	0.52
1:B:368:PHE:CE2	1:B:372:ARG:HD2	2.45	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:LYS:O	1:A:290:THR:HG23	2.09	0.52
1:B:591:SER:OG	1:B:592:GLU:OE1	2.27	0.51
1:B:476:LYS:NZ	6:B:1003:HOH:O	2.43	0.51
1:A:341:ARG:NH2	1:A:362:GLU:O	2.45	0.50
1:B:347[A]:ARG:HG3	1:B:355:ARG:HD3	1.93	0.50
1:A:590:SER:OG	1:A:591:SER:N	2.43	0.50
1:A:316:MET:HB2	1:A:321:ILE:HD11	1.93	0.49
1:A:605:LEU:HD22	1:A:608:ARG:HD2	1.94	0.49
1:B:211:SER:N	6:B:1038:HOH:O	2.42	0.49
1:B:553:HIS:HD2	6:B:1003:HOH:O	1.96	0.49
1:A:549:LEU:HA	1:A:552:LEU:HB2	1.95	0.49
1:A:138:ALA:O	1:A:176:VAL:HG23	2.13	0.48
1:B:341:ARG:HD3	6:B:1042:HOH:O	2.13	0.48
1:B:529:PHE:CE2	1:B:533:LYS:HE3	2.48	0.48
1:B:380:LEU:HD22	1:B:420:LEU:HD11	1.94	0.48
1:A:512:ILE:HB	1:A:562:LEU:HG	1.95	0.48
1:A:657:LEU:HD13	1:A:659:ILE:HD11	1.96	0.48
1:A:440:PHE:O	1:A:444:GLN:HG2	2.14	0.48
1:A:114:ASP:OD1	1:A:118:ARG:HD2	2.13	0.48
1:B:598:LEU:HD11	1:B:614:VAL:HG23	1.96	0.47
1:B:573[A]:HIS:HE1	1:B:658:GLU:OE2	1.97	0.47
1:B:520:ARG:HE	1:B:524:GLU:CD	2.15	0.47
1:A:490:LEU:CD1	1:A:496:THR:HG23	2.45	0.47
1:A:598:LEU:HD23	1:A:601:MET:HE1	1.97	0.47
1:B:408:SER:HB3	1:B:411:ILE:HG12	1.96	0.47
1:B:608:ARG:NH1	6:B:1050:HOH:O	2.21	0.47
1:A:598:LEU:O	1:A:602:ARG:HB2	2.16	0.46
1:B:598:LEU:O	1:B:602:ARG:HG3	2.15	0.46
1:A:126:VAL:CG1	1:A:129:ARG:NH2	2.78	0.46
1:A:466:GLY:O	1:A:470:THR:HG23	2.14	0.46
1:A:139:LEU:HD22	1:A:156:MET:HB3	1.96	0.46
1:A:657:LEU:CD1	1:A:659:ILE:HD11	2.46	0.46
1:A:588:ARG:HB2	1:A:667:LYS:HE2	1.97	0.46
1:B:377:SER:HB2	1:B:403[B]:ARG:NH1	2.31	0.46
1:B:315:MET:HG2	6:B:1009:HOH:O	2.16	0.45
1:A:116:VAL:HG13	1:A:218:VAL:HB	1.97	0.45
1:B:575:LYS:HB2	1:B:577:GLU:HG2	1.98	0.45
1:A:657:LEU:C	1:A:657:LEU:HD12	2.37	0.45
1:A:126:VAL:CG1	1:A:129:ARG:CZ	2.95	0.45
1:B:116:VAL:HG11	1:B:221:TYR:HB2	1.98	0.45
1:B:317:GLU:OE2	1:B:347[A]:ARG:NH1	2.47	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:ALA:N	1:B:209:SER:HA	2.33	0.44
1:B:431:ASP:O	1:B:435[A]:GLN:HG2	2.17	0.44
1:A:517:ALA:O	1:A:543:PHE:HA	2.18	0.44
1:A:313:LEU:HG	1:A:321:ILE:HD13	2.00	0.43
1:A:278:GLU:O	1:A:281:SER:HB3	2.17	0.43
1:A:272:LEU:O	1:A:277[B]:LYS:HE3	2.18	0.43
1:A:518:PRO:HB2	1:A:522:LEU:HD12	2.01	0.43
1:A:703:PRO:HA	1:B:522:LEU:HD21	2.00	0.43
1:A:304:GLY:HA2	1:B:87:HIS:O	2.18	0.43
1:B:230:VAL:HG22	1:B:270:LEU:HD22	2.01	0.43
1:A:158:ILE:HB	1:A:300:ILE:HD13	2.01	0.43
1:A:186:ASN:O	6:A:901:HOH:O	2.21	0.43
1:B:248:SER:HB2	1:B:254:PHE:HB3	2.01	0.42
1:B:456:PHE:CE1	1:B:463:MET:HE3	2.53	0.42
1:A:521:HIS:HE1	1:A:625:ILE:O	2.02	0.42
1:B:235:GLN:NE2	1:B:241:ALA:O	2.43	0.42
1:A:522:LEU:HB3	1:A:630:MET:HE2	2.01	0.41
1:B:237:ALA:HB2	1:B:264:GLN:NE2	2.35	0.41
1:A:232:VAL:HB	1:A:246:TRP:HB3	2.01	0.41
1:A:230:VAL:HG22	1:A:270:LEU:HD22	2.02	0.41
1:B:603:ASN:O	1:B:606:VAL:HG23	2.21	0.41
1:A:118:ARG:HH21	1:A:273:LYS:NZ	2.19	0.41
1:B:495[B]:GLN:HG3	6:B:930:HOH:O	2.20	0.41
1:B:278:GLU:O	1:B:281:SER:HB3	2.21	0.40
1:A:217:GLY:HA2	3:A:802:ADP:O3A	2.22	0.40
1:A:532:MET:H	1:A:532:MET:HG2	1.63	0.40
1:A:152:ASP:O	1:A:154:ALA:N	2.55	0.40
1:B:495[B]:GLN:H	1:B:495[B]:GLN:CD	2.25	0.40
1:B:665:LEU:HD12	1:B:665:LEU:HA	1.92	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	590/719 (82%)	558 (95%)	32 (5%)	0	100	100
1	B	603/719 (84%)	585 (97%)	17 (3%)	1 (0%)	52	64
All	All	1193/1438 (83%)	1143 (96%)	49 (4%)	1 (0%)	56	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	637	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	524/634 (83%)	523 (100%)	1 (0%)	95	98
1	B	546/634 (86%)	543 (100%)	3 (0%)	92	97
All	All	1070/1268 (84%)	1066 (100%)	4 (0%)	93	97

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	549	LEU
1	B	126	VAL
1	B	234	SER
1	B	378	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	335	GLN
1	A	521	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ADP	A	802	5	22,29,29	0.99	1 (4%)	27,45,45	1.82	4 (14%)
4	ALF	A	803	-	0,4,4	0.00	-	0,6,6	0.00	-
3	ADP	B	803	5	22,29,29	1.06	2 (9%)	27,45,45	2.08	7 (25%)
4	ALF	B	804	-	0,4,4	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	802	5	-	0/12/32/32	0/3/3/3
4	ALF	A	803	-	-	0/0/0/0	0/0/0/0
3	ADP	B	803	5	-	0/12/32/32	0/3/3/3
4	ALF	B	804	-	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	803	ADP	O4'-C1'	2.07	1.43	1.41
3	A	802	ADP	C5-C4	3.25	1.47	1.40
3	B	803	ADP	C5-C4	3.35	1.48	1.40

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	ADP	N3-C2-N1	-6.95	123.57	128.89
3	B	803	ADP	N3-C2-N1	-6.80	123.69	128.89
3	B	803	ADP	C2'-C1'-N9	-4.49	107.43	114.29
3	B	803	ADP	C4-C5-N7	-3.55	106.21	109.48
3	A	802	ADP	C4-C5-N7	-3.29	106.46	109.48
3	A	802	ADP	PA-O3A-PB	-2.07	125.72	132.67
3	B	803	ADP	PA-O3A-PB	-2.03	125.85	132.67
3	A	802	ADP	O2A-PA-O1A	2.14	124.13	112.53
3	B	803	ADP	O3B-PB-O1B	2.37	118.22	110.58
3	B	803	ADP	C2-N1-C6	2.39	123.04	118.77
3	B	803	ADP	O3B-PB-O2B	2.77	117.94	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	ADP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	596/719 (82%)	0.56	49 (8%)	14 20	33, 65, 121, 154	0
1	B	609/719 (84%)	0.22	22 (3%)	46 55	34, 59, 105, 131	0
All	All	1205/1438 (83%)	0.39	71 (5%)	26 34	33, 61, 116, 154	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	374	MET	7.4
1	B	652	ILE	6.7
1	A	508	GLY	5.8
1	A	607	GLN	5.5
1	A	534	GLN	5.3
1	A	659	ILE	5.3
1	A	589	LEU	4.9
1	A	99	SER	4.9
1	B	510	ARG	4.9
1	A	670	HIS	4.0
1	B	653	LEU	4.0
1	B	463	MET	3.7
1	B	209	SER	3.7
1	A	472	GLU	3.7
1	A	617	ARG	3.5
1	A	498	LEU	3.4
1	A	672	LEU	3.3
1	A	152	ASP	3.3
1	A	606	VAL	3.3
1	B	389	THR	3.3
1	A	509	THR	3.2
1	A	597	LEU	3.2
1	A	600	TRP	3.2
1	A	601	MET	3.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	153	THR	3.1
1	A	555	ARG	3.1
1	A	96	VAL	3.0
1	B	232	VAL	2.9
1	B	237	ALA	2.9
1	B	636	PHE	2.9
1	A	590	SER	2.9
1	A	668	LYS	2.8
1	B	238	GLU	2.8
1	A	520	ARG	2.8
1	A	145	ARG	2.7
1	A	100	PHE	2.7
1	A	510	ARG	2.7
1	B	146	MET	2.7
1	A	637	LEU	2.7
1	A	507	ALA	2.7
1	A	609	VAL	2.7
1	A	679	LEU	2.7
1	A	240	ASP	2.7
1	B	262	VAL	2.7
1	B	607	GLN	2.6
1	A	716	LEU	2.6
1	B	427	ARG	2.6
1	A	665	LEU	2.5
1	A	677	PRO	2.5
1	B	576	GLU	2.5
1	A	612	ILE	2.5
1	A	669	LEU	2.4
1	A	239	ALA	2.4
1	B	393	ASP	2.4
1	B	240	ASP	2.3
1	A	537	MET	2.3
1	A	598	LEU	2.3
1	A	717	GLU	2.3
1	A	97	GLN	2.3
1	A	116	VAL	2.3
1	A	366	SER	2.3
1	B	100	PHE	2.3
1	A	535	LYS	2.2
1	B	208	ALA	2.2
1	B	373	GLU	2.2
1	A	567	THR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	335	GLN	2.1
1	A	513	TYR	2.1
1	A	594	ALA	2.0
1	B	699	GLU	2.0
1	A	680	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	ALF	A	803	5/5	0.94	0.18	0.66	29,31,33,33	0
5	MG	A	804	1/1	0.86	0.16	-0.09	37,37,37,37	0
3	ADP	A	802	27/27	0.98	0.14	-0.31	27,39,43,45	0
4	ALF	B	804	5/5	0.98	0.12	-0.67	46,50,50,51	5
3	ADP	B	803	27/27	0.98	0.10	-1.13	41,52,57,60	0
5	MG	B	805	1/1	0.96	0.11	-1.13	52,52,52,52	0
2	CO	A	805	1/1	0.15	0.19	-	109,109,109,109	1
2	CO	A	801	1/1	0.93	0.15	-	109,109,109,109	1
2	CO	B	802	1/1	0.92	0.13	-	109,109,109,109	1
2	CO	B	801	1/1	0.88	0.23	-	96,96,96,96	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.