



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 31, 2016 – 05:45 AM EDT

PDB ID : 5J6F  
Title : Crystal structure of DAH7PS-CM complex from *Geobacillus* sp. with prephenate  
Authors : Nazmi, A.R.; Othman, M.; Lang, E.J.M.; Bai, Y.; Allison, T.M.; Panjkar, S.; Arcus, V.L.; Parker, E.J.  
Deposited on : 2016-04-04  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	 82% 13% . .
1	B	368	 82% 10% . 8%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase, chorismate mutase-isozyme 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2741	1730	485	516	10			
1	B	340	Total	C	N	O	S	0	0	0
			2625	1657	463	494	11			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLY	SER	conflict	UNP L8A208
A	359	ARG	GLN	conflict	UNP L8A208
A	361	LEU	-	expression tag	UNP L8A208
A	362	GLU	-	expression tag	UNP L8A208
A	363	HIS	-	expression tag	UNP L8A208
A	364	HIS	-	expression tag	UNP L8A208
A	365	HIS	-	expression tag	UNP L8A208
A	366	HIS	-	expression tag	UNP L8A208
A	367	HIS	-	expression tag	UNP L8A208
A	368	HIS	-	expression tag	UNP L8A208
B	2	GLY	SER	conflict	UNP L8A208
B	359	ARG	GLN	conflict	UNP L8A208
B	361	LEU	-	expression tag	UNP L8A208
B	362	GLU	-	expression tag	UNP L8A208
B	363	HIS	-	expression tag	UNP L8A208
B	364	HIS	-	expression tag	UNP L8A208
B	365	HIS	-	expression tag	UNP L8A208
B	366	HIS	-	expression tag	UNP L8A208
B	367	HIS	-	expression tag	UNP L8A208
B	368	HIS	-	expression tag	UNP L8A208

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is PREPHENIC ACID (three-letter code: PRE) (formula: C<sub>10</sub>H<sub>10</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	10	6		
4	B	1	Total	C	O	0	0
			16	10	6		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase, chorismate mutase-isozyme 3



- Molecule 1: 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase, chorismate mutase-isozyme 3



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.44Å 95.44Å 167.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	82.65 – 2.75 19.61 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.0 (82.65-2.75) 99.3 (19.61-2.75)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 2.75Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, $R_{free}$	0.182 , 0.235 0.186 , 0.232	Depositor DCC
$R_{free}$ test set	1054 reflections (4.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	74.8	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 33.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.095 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5420	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PRE, MN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.99	1/2781 (0.0%)	1.10	13/3755 (0.3%)
1	B	0.95	0/2661	1.06	8/3593 (0.2%)
All	All	0.97	1/5442 (0.0%)	1.08	21/7348 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	239	GLU	CD-OE2	7.08	1.33	1.25

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	ASP	CB-CG-OD1	-7.40	111.64	118.30
1	A	63	ASP	CB-CG-OD2	7.35	124.91	118.30
1	A	5	ARG	NE-CZ-NH2	-7.10	116.75	120.30
1	A	80	LYS	CA-CB-CG	6.83	128.42	113.40
1	A	332	SER	CB-CA-C	-6.62	97.53	110.10
1	B	21	LEU	CA-CB-CG	6.57	130.42	115.30
1	A	52	ARG	NE-CZ-NH1	6.50	123.55	120.30
1	A	301	ARG	CG-CD-NE	6.34	125.11	111.80
1	B	301	ARG	NE-CZ-NH1	6.33	123.47	120.30
1	A	29	ARG	NE-CZ-NH1	6.08	123.34	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	261	ARG	NE-CZ-NH1	5.97	123.29	120.30
1	A	268	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	B	56	ASP	CB-CG-OD1	5.55	123.29	118.30
1	B	286	THR	N-CA-C	-5.54	96.03	111.00
1	B	12	ARG	NE-CZ-NH1	5.47	123.04	120.30
1	A	29	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	B	261	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	B	57	LEU	CA-CB-CG	5.18	127.21	115.30
1	A	351	ARG	NE-CZ-NH1	5.17	122.89	120.30
1	A	91	ARG	NE-CZ-NH2	-5.15	117.73	120.30
1	A	157	ARG	NE-CZ-NH1	5.03	122.81	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	156	PRO	Peptide
1	B	352	ALA	Peptide
1	B	353	PHE	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2741	0	2773	19	0
1	B	2625	0	2650	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	10	0	0	0	0
3	B	10	0	0	1	0
4	A	16	0	8	0	0
4	B	16	0	8	0	0
All	All	5420	0	5439	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:ARG:HH12	1:A:104:ASN:HB3	1.52	0.74
1:A:276:ILE:HD13	1:B:276:ILE:HD11	1.73	0.71
1:A:41:GLN:HG3	1:A:43:THR:HG23	1.73	0.71
1:A:94:LEU:HD22	1:A:246:GLU:HG3	1.74	0.69
1:B:261:ARG:NE	3:B:402:SO4:O1	2.27	0.67
1:A:41:GLN:HG3	1:A:43:THR:CG2	2.33	0.59
1:A:123:MET:O	1:A:149:LEU:HD12	2.03	0.58
1:B:123:MET:O	1:B:149:LEU:HD12	2.04	0.57
1:A:101:HIS:HB3	1:A:287:HIS:CD2	2.40	0.56
1:A:305:ILE:HB	1:A:306:PRO:HD3	1.89	0.54
1:B:305:ILE:HB	1:B:306:PRO:HD3	1.90	0.53
1:B:188:GLU:HA	1:B:206:GLN:HB3	1.93	0.50
1:A:39:GLU:OE2	1:A:90:HIS:ND1	2.43	0.49
1:A:157:ARG:NH2	1:A:331:LEU:O	2.47	0.47
1:A:95:LEU:N	1:A:246:GLU:OE2	2.48	0.47
1:A:188:GLU:HA	1:A:206:GLN:HB3	1.97	0.46
1:B:194:ASP:HA	1:B:197:ILE:HD12	1.98	0.46
1:A:122:VAL:HB	1:A:320:MET:HG3	1.99	0.45
1:A:222:GLY:HA3	1:A:251:GLN:HB2	1.99	0.45
1:A:154:TYR:CD1	1:A:154:TYR:N	2.85	0.44
1:A:5:ARG:HA	1:A:8:GLU:HG2	1.97	0.44
1:B:302:ASP:OD1	1:B:302:ASP:N	2.42	0.43
1:A:194:ASP:HA	1:A:197:ILE:HD12	1.99	0.43
1:B:5:ARG:HA	1:B:8:GLU:HG2	2.01	0.41
1:A:55:LEU:HD22	1:A:72:GLN:HG2	2.01	0.41
1:A:302:ASP:OD1	1:A:302:ASP:N	2.41	0.41
1:B:122:VAL:HB	1:B:320:MET:HG3	2.02	0.41
1:B:284:LYS:HA	1:B:284:LYS:HD3	1.92	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/368 (95%)	343 (98%)	7 (2%)	0	100	100
1	B	336/368 (91%)	332 (99%)	4 (1%)	0	100	100
All	All	686/736 (93%)	675 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/308 (94%)	277 (96%)	11 (4%)	40	73
1	B	273/308 (89%)	256 (94%)	17 (6%)	23	51
All	All	561/616 (91%)	533 (95%)	28 (5%)	30	61

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	21	LEU
1	A	63	ASP
1	A	107	VAL
1	A	113	ARG
1	A	150	ARG
1	A	191	THR
1	A	271	ARG
1	A	277	SER
1	A	302	ASP
1	A	340	ILE
1	B	12	ARG
1	B	15	GLU
1	B	45	ARG
1	B	63	ASP
1	B	107	VAL
1	B	150	ARG
1	B	157	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	158	THR
1	B	164	GLN
1	B	191	THR
1	B	239	GLU
1	B	271	ARG
1	B	277	SER
1	B	302	ASP
1	B	348	GLU
1	B	351	ARG
1	B	354	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	41	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	402	-	4,4,4	0.67	0	6,6,6	0.33	0
3	SO4	A	403	-	4,4,4	0.45	0	6,6,6	0.28	0
4	PRE	A	404	-	10,16,16	2.31	5 (50%)	5,23,23	1.14	0
3	SO4	B	402	-	4,4,4	0.62	0	6,6,6	0.52	0
3	SO4	B	403	-	4,4,4	0.47	0	6,6,6	0.23	0
4	PRE	B	404	-	10,16,16	2.22	5 (50%)	5,23,23	1.18	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	402	-	-	0/0/0/0	0/0/0/0
3	SO4	A	403	-	-	0/0/0/0	0/0/0/0
4	PRE	A	404	-	-	0/5/27/27	0/1/1/1
3	SO4	B	402	-	-	0/0/0/0	0/0/0/0
3	SO4	B	403	-	-	0/0/0/0	0/0/0/0
4	PRE	B	404	-	-	0/5/27/27	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	404	PRE	C8-C1	-3.74	1.49	1.57
4	B	404	PRE	C8-C1	-3.43	1.50	1.57
4	A	404	PRE	C1-C6	-3.16	1.42	1.52
4	B	404	PRE	C1-C6	-3.16	1.42	1.52
4	A	404	PRE	C1-C2	-3.05	1.42	1.52
4	B	404	PRE	C1-C2	-3.04	1.42	1.52
4	A	404	PRE	C3-C2	2.72	1.36	1.32
4	B	404	PRE	C3-C2	2.73	1.36	1.32
4	B	404	PRE	C5-C6	2.74	1.36	1.32
4	A	404	PRE	C5-C6	3.04	1.37	1.32

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	404	PRE	O1'-C1'-C8	2.34	123.51	120.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	352/368 (95%)	-0.52	0 100 100	54, 74, 96, 118	0
1	B	340/368 (92%)	-0.43	2 (0%) 90 88	59, 77, 107, 161	0
All	All	692/736 (94%)	-0.48	2 (0%) 94 93	54, 75, 99, 161	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	3.3
1	B	160	PRO	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	402	5/5	0.98	0.16	0.72	73,78,83,91	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	403	5/5	0.98	0.14	0.57	69,70,84,88	0
4	PRE	A	404	16/16	0.93	0.15	0.14	62,107,122,126	0
3	SO4	B	403	5/5	0.97	0.19	0.13	90,91,100,107	0
4	PRE	B	404	16/16	0.95	0.12	-0.12	63,79,84,86	0
3	SO4	B	402	5/5	0.97	0.09	-1.02	76,76,81,83	0
2	MN	A	401	1/1	0.99	0.05	-2.06	74,74,74,74	0
2	MN	B	401	1/1	0.99	0.04	-	100,100,100,100	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.