



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:55 AM GMT

PDB ID : 2JAP  
Title : CLAVULANIC ACID DEHYDROGENASE: STRUCTURAL AND BIO-CHEMICAL ANALYSIS OF THE FINAL STEP IN THE BIOSYNTHESIS OF THE BETA-LACTAMASE INHIBITOR CLAVULANIC ACID  
Authors : Mackenzie, A.K.; Kershaw, N.J.; Hernandez, H.; Robinson, C.V.; Schofield, C.J.; Andersson, I.  
Deposited on : 2006-11-29  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	<div> <div>9%</div> <div>94%</div> <div>5%</div> </div>
1	B	247	<div> <div>14%</div> <div>90%</div> <div>9%</div> </div>
1	C	247	<div> <div>11%</div> <div>91%</div> <div>9%</div> </div>
1	D	247	<div> <div>8%</div> <div>94%</div> <div>5%</div> </div>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7847 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CLAVALDEHYDE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	0	0
			1832	1136	337	353	6			
1	B	245	Total	C	N	O	S	0	0	0
			1832	1136	337	353	6			
1	C	246	Total	C	N	O	S	0	1	0
			1850	1147	342	355	6			
1	D	245	Total	C	N	O	S	0	0	0
			1832	1136	337	353	6			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is (2R,3Z,5R)-3-(2-HYDROXYETHYLIDENE)-7-OXO-4-OXA-1-AZABICYCLO[3.2.0]HEPTANE-2-CARBOXYLIC ACID (three-letter code: J01) (formula: C<sub>8</sub>H<sub>9</sub>NO<sub>5</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O		0	0
			14	8	1	5			
3	B	1	Total	C	N	O		0	0
			14	8	1	5			
3	C	1	Total	C	N	O		0	0
			14	8	1	5			
3	D	1	Total	C	N	O		0	0
			14	8	1	5			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	68	Total	O	0	0
			68	68		
4	B	56	Total	O	0	0
			56	56		
4	C	73	Total	O	0	0
			73	73		

Continued on next page...


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	56	Total	O	0	0
			56	56		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: CLAVALDEHYDE DEHYDROGENASE




#### • Molecule 1: CLAVALDEHYDE DEHYDROGENASE



#### • Molecule 1: CLAVALDEHYDE DEHYDROGENASE



#### • Molecule 1: CLAVALDEHYDE DEHYDROGENASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.08Å 123.42Å 126.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.86 – 2.10 34.87 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.2 (34.86-2.10) 99.2 (34.87-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.99 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.184 , 0.216 0.184 , 0.215	Depositor DCC
$R_{free}$ test set	2713 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.9	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.6	EDS
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	2 of 53614 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7847	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.18 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5838e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: J01, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/1852	0.59	0/2513
1	B	0.46	1/1852 (0.1%)	0.59	1/2513 (0.0%)
1	C	0.45	0/1871	0.59	0/2538
1	D	0.44	0/1852	0.58	0/2513
All	All	0.45	1/7427 (0.0%)	0.59	1/10077 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	45	ARG	NE-CZ	6.52	1.41	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	45	ARG	NE-CZ-NH1	-6.74	116.93	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	45	ARG	Sidechain



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1832	0	1871	12	0
1	B	1832	0	1871	11	0
1	C	1850	0	1891	12	0
1	D	1832	0	1871	7	0
2	A	48	0	26	5	0
2	B	48	0	26	5	0
2	C	48	0	26	5	0
2	D	48	0	26	5	0
3	A	14	0	8	4	0
3	B	14	0	8	3	0
3	C	14	0	8	3	0
3	D	14	0	8	3	0
4	A	68	0	0	2	0
4	B	56	0	0	0	0
4	C	73	0	0	1	0
4	D	56	0	0	0	0
All	All	7847	0	7640	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1248:NDP:H41N	3:A:1249:J01:H7C1	1.53	0.89
2:C:1248:NDP:H41N	3:C:1249:J01:H7C1	1.58	0.85
2:A:1248:NDP:C4N	3:A:1249:J01:H7C1	2.21	0.71
2:C:1248:NDP:C4N	3:C:1249:J01:H7C1	2.21	0.70
2:B:1248:NDP:H41N	3:B:1249:J01:H7C1	1.73	0.70
1:C:209:ILE:HD12	1:C:214:LYS:HE2	1.77	0.66
1:A:216:GLN:HB3	1:A:218:GLN:HE21	1.59	0.66
1:A:141:SER:O	2:A:1248:NDP:H6N	1.97	0.65
1:D:68:ARG:HH12	1:D:69:GLN:HE21	1.48	0.60
1:B:117:LEU:HA	1:B:120:MET:HE2	1.84	0.59
2:B:1248:NDP:C4N	3:B:1249:J01:H7C1	2.34	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:ILE:HD12	1:A:214:LYS:HE2	1.85	0.56
2:A:1248:NDP:C3N	3:A:1249:J01:H7C1	2.36	0.56
2:D:1248:NDP:H41N	3:D:1249:J01:H7C1	1.87	0.56
1:C:14:GLY:HA2	2:C:1248:NDP:H1B	1.87	0.55
2:C:1248:NDP:C3N	3:C:1249:J01:H7C1	2.36	0.55
2:D:1248:NDP:C3N	3:D:1249:J01:H7C1	2.37	0.55
1:A:218:GLN:CD	1:A:218:GLN:H	2.11	0.54
1:B:141:SER:O	2:B:1248:NDP:H6N	2.08	0.52
1:C:141:SER:O	2:C:1248:NDP:H6N	2.10	0.52
1:D:14:GLY:HA2	2:D:1248:NDP:H1B	1.92	0.51
1:D:141:SER:O	2:D:1248:NDP:H6N	2.11	0.51
1:C:195:HIS:HE1	4:C:2070:HOH:O	1.95	0.50
1:C:25:ARG:NH1	1:C:50:GLU:OE2	2.45	0.50
1:C:91:ASN:HD22	1:C:140:MET:HG2	1.78	0.49
2:D:1248:NDP:C4N	3:D:1249:J01:H7C1	2.43	0.49
1:B:9:VAL:HG11	1:B:84:GLY:O	2.13	0.48
2:B:1248:NDP:C3N	3:B:1249:J01:H7C1	2.43	0.48
1:B:231:PRO:HB2	1:B:233:HIS:CE1	2.48	0.48
2:A:1248:NDP:H41N	3:A:1249:J01:C7	2.35	0.48
1:C:143:ILE:HG12	1:C:147:VAL:CG2	2.44	0.48
1:C:193:ARG:NE	1:C:206:GLU:OE2	2.44	0.47
1:A:117:LEU:HD13	1:C:112:ILE:HD12	1.95	0.47
1:B:203:GLU:O	1:B:207:GLN:HG2	2.15	0.47
1:D:79:VAL:HG21	1:D:129:HIS:CE1	2.50	0.46
1:D:91:ASN:HD22	1:D:140:MET:HG2	1.82	0.45
1:B:79:VAL:HG21	1:B:129:HIS:CD2	2.51	0.45
1:D:170:ARG:O	1:D:174:THR:HB	2.17	0.45
1:B:91:ASN:HD22	1:B:140:MET:HG2	1.83	0.43
1:A:79:VAL:HG21	1:A:129:HIS:CE1	2.53	0.43
1:D:41:VAL:O	1:D:45:ARG:HG3	2.19	0.43
1:A:214:LYS:HE3	4:A:2049:HOH:O	2.17	0.42
1:A:195:HIS:HE1	4:A:2062:HOH:O	2.02	0.42
1:B:139:GLN:HG3	1:B:166:SER:OG	2.20	0.42
1:A:207:GLN:OE1	1:A:207:GLN:HA	2.20	0.41
1:A:112:ILE:HD12	1:C:117:LEU:HD13	2.03	0.41
1:B:42:GLU:HG2	1:B:43:LYS:N	2.36	0.41
1:C:108:TRP:CD1	1:C:154:VAL:HG21	2.56	0.41
1:A:108:TRP:CZ3	1:C:120:MET:HB3	2.56	0.41
1:A:21:GLU:OE2	1:A:25:ARG:NH2	2.52	0.40
1:B:170:ARG:O	1:B:174:THR:HB	2.21	0.40
1:B:14:GLY:HA2	2:B:1248:NDP:H1B	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/247 (98%)	238 (98%)	5 (2%)	0	100	100
1	B	243/247 (98%)	238 (98%)	5 (2%)	0	100	100
1	C	245/247 (99%)	239 (98%)	6 (2%)	0	100	100
1	D	243/247 (98%)	238 (98%)	5 (2%)	0	100	100
All	All	974/988 (99%)	953 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/190 (99%)	187 (100%)	1 (0%)	92	95
1	B	188/190 (99%)	185 (98%)	3 (2%)	70	76
1	C	190/190 (100%)	186 (98%)	4 (2%)	61	66
1	D	188/190 (99%)	185 (98%)	3 (2%)	70	76
All	All	754/760 (99%)	743 (98%)	11 (2%)	72	78

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	218	GLN
1	B	9	VAL
1	B	41	VAL
1	B	69	GLN
1	C	42	GLU
1	C	102	ASP
1	C	218	GLN
1	C	225	ARG
1	D	11	LEU
1	D	41	VAL
1	D	117	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	218	GLN
1	B	91	ASN
1	B	129	HIS
1	C	91	ASN
1	C	195	HIS
1	D	69	GLN
1	D	91	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NDP	A	1248	-	42,52,52	1.54	5 (11%)	55,80,80	1.89	4 (7%)
3	J01	A	1249	-	9,15,15	1.81	3 (33%)	7,22,22	3.93	4 (57%)
2	NDP	B	1248	-	42,52,52	1.58	5 (11%)	55,80,80	1.93	3 (5%)
3	J01	B	1249	-	9,15,15	2.03	3 (33%)	7,22,22	4.06	4 (57%)
2	NDP	C	1248	-	42,52,52	1.53	5 (11%)	55,80,80	1.93	4 (7%)
3	J01	C	1249	-	9,15,15	1.74	3 (33%)	7,22,22	3.85	5 (71%)
2	NDP	D	1248	-	42,52,52	1.56	5 (11%)	55,80,80	1.81	3 (5%)
3	J01	D	1249	-	9,15,15	2.00	2 (22%)	7,22,22	3.62	4 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1248	-	-	0/30/77/77	0/5/5/5
3	J01	A	1249	-	-	0/2/31/31	0/2/2/2
2	NDP	B	1248	-	-	0/30/77/77	0/5/5/5
3	J01	B	1249	-	-	0/2/31/31	0/2/2/2
2	NDP	C	1248	-	-	0/30/77/77	0/5/5/5
3	J01	C	1249	-	-	0/2/31/31	0/2/2/2
2	NDP	D	1248	-	-	0/30/77/77	0/5/5/5
3	J01	D	1249	-	-	0/2/31/31	0/2/2/2

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1249	J01	C4-C5	-2.34	1.49	1.52
3	B	1249	J01	C4-C5	-2.14	1.49	1.52
3	C	1249	J01	C4-C5	-2.04	1.50	1.52
2	C	1248	NDP	C2N-C3N	2.09	1.39	1.34
2	B	1248	NDP	C2N-C3N	2.18	1.40	1.34
2	A	1248	NDP	C2N-C3N	2.20	1.40	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1248	NDP	C2N-C3N	2.25	1.40	1.34
3	A	1249	J01	C6-C1	2.47	1.34	1.31
3	C	1249	J01	C6-C1	2.59	1.34	1.31
2	A	1248	NDP	C2A-N1A	2.71	1.39	1.33
2	C	1248	NDP	C2A-N1A	2.72	1.39	1.33
2	D	1248	NDP	C2A-N1A	2.74	1.39	1.33
2	C	1248	NDP	C6N-C5N	2.88	1.38	1.33
2	B	1248	NDP	C2A-N1A	2.98	1.39	1.33
2	A	1248	NDP	C6N-C5N	3.03	1.39	1.33
2	B	1248	NDP	C6N-C5N	3.05	1.39	1.33
2	D	1248	NDP	C6N-C5N	3.08	1.39	1.33
2	C	1248	NDP	C2A-N3A	3.45	1.38	1.32
3	D	1249	J01	C6-C1	3.46	1.35	1.31
3	B	1249	J01	C6-C1	3.49	1.35	1.31
2	B	1248	NDP	C2A-N3A	3.64	1.38	1.32
2	A	1248	NDP	C2A-N3A	3.65	1.38	1.32
2	D	1248	NDP	C2A-N3A	3.72	1.38	1.32
3	C	1249	J01	O1-C1	3.79	1.44	1.38
3	A	1249	J01	O1-C1	4.04	1.45	1.38
3	D	1249	J01	O1-C1	4.29	1.45	1.38
3	B	1249	J01	O1-C1	4.31	1.45	1.38
2	A	1248	NDP	O7N-C7N	6.71	1.41	1.24
2	D	1248	NDP	O7N-C7N	6.82	1.41	1.24
2	B	1248	NDP	O7N-C7N	6.86	1.41	1.24
2	C	1248	NDP	O7N-C7N	6.95	1.41	1.24

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1248	NDP	N3A-C2A-N1A	-12.39	119.41	128.89
2	B	1248	NDP	N3A-C2A-N1A	-12.31	119.47	128.89
2	A	1248	NDP	N3A-C2A-N1A	-11.83	119.83	128.89
2	D	1248	NDP	N3A-C2A-N1A	-11.31	120.24	128.89
3	A	1249	J01	O2-C5-C4	-5.24	131.13	136.78
3	B	1249	J01	O2-C5-C4	-5.16	131.22	136.78
3	D	1249	J01	O2-C5-C4	-4.98	131.41	136.78
3	C	1249	J01	O2-C5-C4	-4.84	131.56	136.78
3	B	1249	J01	C3-C4-C5	-4.74	81.70	85.20
3	C	1249	J01	C3-C4-C5	-4.41	81.93	85.20
3	A	1249	J01	C3-C4-C5	-4.41	81.94	85.20
3	B	1249	J01	C3-N1-C5	-4.16	86.35	93.30
3	D	1249	J01	C3-C4-C5	-4.04	82.21	85.20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1249	J01	C3-N1-C5	-3.93	86.75	93.30
3	C	1249	J01	C3-N1-C5	-3.92	86.75	93.30
3	D	1249	J01	C3-N1-C5	-3.87	86.84	93.30
2	D	1248	NDP	PN-O3-PA	-2.91	124.55	132.73
2	C	1248	NDP	PN-O3-PA	-2.86	124.70	132.73
2	B	1248	NDP	PN-O3-PA	-2.80	124.86	132.73
2	B	1248	NDP	C1B-N9A-C4A	-2.75	122.78	126.94
2	A	1248	NDP	C1B-N9A-C4A	-2.51	123.16	126.94
2	D	1248	NDP	C1B-N9A-C4A	-2.21	123.60	126.94
2	A	1248	NDP	C1D-N1N-C6N	-2.17	115.96	120.81
2	A	1248	NDP	PN-O3-PA	-2.11	126.79	132.73
2	C	1248	NDP	C1B-N9A-C4A	-2.04	123.86	126.94
2	C	1248	NDP	C4A-C5A-N7A	-2.02	107.62	109.48
3	C	1249	J01	O3-C7-C6	2.38	114.87	110.70
3	D	1249	J01	C4-C5-N1	5.59	97.38	92.56
3	C	1249	J01	C4-C5-N1	6.03	97.77	92.56
3	A	1249	J01	C4-C5-N1	6.26	97.96	92.56
3	B	1249	J01	C4-C5-N1	6.73	98.37	92.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1248	NDP	5	0
3	A	1249	J01	4	0
2	B	1248	NDP	5	0
3	B	1249	J01	3	0
2	C	1248	NDP	5	0
3	C	1249	J01	3	0
2	D	1248	NDP	5	0
3	D	1249	J01	3	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	245/247 (99%)	0.50	23 (9%) 11 14	30, 32, 36, 40	0
1	B	245/247 (99%)	0.73	35 (14%) 4 5	30, 32, 35, 41	0
1	C	246/247 (99%)	0.48	27 (10%) 7 10	30, 32, 36, 42	0
1	D	245/247 (99%)	0.50	19 (7%) 16 22	29, 32, 36, 39	0
All	All	981/988 (99%)	0.55	104 (10%) 8 11	29, 32, 36, 42	0

All (104) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	210	SER	6.1
1	A	42	GLU	4.6
1	B	6	GLN	4.5
1	A	199	THR	4.3
1	C	199	THR	4.2
1	D	204	MET	4.0
1	B	204	MET	3.9
1	D	207	GLN	3.7
1	B	81	ALA	3.7
1	B	69	GLN	3.6
1	B	102	ASP	3.6
1	A	3	SER	3.5
1	A	204	MET	3.5
1	B	80	GLU	3.5
1	D	7	GLY	3.5
1	A	200	ALA	3.4
1	D	200	ALA	3.4
1	D	199	THR	3.3
1	A	181	VAL	3.3
1	B	182	VAL	3.1
1	B	51	LEU	3.1

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	208	ARG	3.1
1	C	239	ILE	3.1
1	B	59	HIS	3.1
1	D	203	GLU	3.0
1	C	54	ALA	3.0
1	A	240	PHE	3.0
1	C	4	ALA	3.0
1	A	54	ALA	2.9
1	B	211	GLN	2.9
1	A	183	ILE	2.9
1	C	42	GLU	2.9
1	C	203	GLU	2.9
1	B	76	ALA	2.9
1	D	205	TYR	2.9
1	D	182	VAL	2.8
1	C	50	GLU	2.8
1	B	41	VAL	2.8
1	C	207	GLN	2.8
1	C	240	PHE	2.8
1	C	157	ALA	2.8
1	A	239	ILE	2.7
1	D	206	GLU	2.7
1	C	80	GLU	2.7
1	A	207	GLN	2.7
1	B	199	THR	2.7
1	C	53	ALA	2.7
1	D	81	ALA	2.7
1	B	45	ARG	2.7
1	B	205	TYR	2.7
1	C	161	GLY	2.7
1	D	208	ARG	2.6
1	B	162	VAL	2.6
1	A	201	THR	2.6
1	C	182	VAL	2.6
1	C	183	ILE	2.6
1	B	42	GLU	2.6
1	B	84	GLY	2.6
1	C	220	ILE	2.6
1	A	41	VAL	2.6
1	D	239	ILE	2.6
1	B	57	LYS	2.5
1	A	162	VAL	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	210	SER	2.5
1	C	2	PRO	2.5
1	B	207	GLN	2.5
1	A	236	VAL	2.5
1	A	203	GLU	2.5
1	C	241	ILE	2.5
1	A	144	ALA	2.4
1	C	164	ALA	2.4
1	A	197	THR	2.4
1	B	236	VAL	2.4
1	B	35	ALA	2.4
1	B	183	ILE	2.4
1	B	72	ASP	2.4
1	B	60	VAL	2.4
1	B	181	VAL	2.4
1	D	210	SER	2.4
1	B	195	HIS	2.4
1	B	58	VAL	2.3
1	A	182	VAL	2.3
1	C	198	HIS	2.3
1	D	212	ILE	2.3
1	A	45	ARG	2.3
1	C	138	VAL	2.3
1	B	103	ALA	2.3
1	B	164	ALA	2.3
1	D	211	GLN	2.3
1	D	183	ILE	2.3
1	D	69	GLN	2.2
1	B	138	VAL	2.2
1	C	180	VAL	2.2
1	D	62	GLU	2.1
1	B	212	ILE	2.1
1	A	209	ILE	2.1
1	C	204	MET	2.1
1	D	160	PHE	2.1
1	B	161	GLY	2.0
1	C	210	SER	2.0
1	C	201	THR	2.0
1	A	145	GLY	2.0
1	C	196	ILE	2.0
1	C	162	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	J01	C	1249	14/14	0.86	0.17	0.70	42,45,46,46	0
3	J01	B	1249	14/14	0.79	0.19	0.66	37,41,42,42	0
3	J01	D	1249	14/14	0.85	0.17	0.42	37,41,42,42	0
3	J01	A	1249	14/14	0.81	0.16	0.06	36,41,41,42	0
2	NDP	D	1248	48/48	0.95	0.10	-1.10	32,34,35,35	0
2	NDP	B	1248	48/48	0.95	0.10	-1.23	28,31,34,35	0
2	NDP	A	1248	48/48	0.96	0.09	-1.26	29,33,35,35	0
2	NDP	C	1248	48/48	0.96	0.08	-1.33	29,32,34,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.