



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:55 AM GMT

PDB ID : 2JBV
Title : CRYSTAL STRUCTURE OF CHOLINE OXIDASE REVEALS INSIGHTS
INTO THE CATALYTIC MECHANISM
Authors : Lountos, G.T.; Fan, F.; Gadda, G.; Orville, A.M.
Deposited on : 2006-12-13
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	546	
1	B	546	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DMS	B	1533	-	-	-	X
4	UNX	A	2501	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UNX	B	2500	-	-	-	X
4	UNX	B	2501	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHOLINE OXIDASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	527	Total	C	N	O	S	0	2	0
			4071	2532	731	784	24			
1	B	530	Total	C	N	O	S	0	0	0
			4094	2544	736	790	24			

- Molecule 2 is [(2R,3S,4R,5R)-5-(6-AMINO-9H-PURIN-9-YL)-3,4-DIHYDROXYTETRAHYDROFURAN-2-YL]METHYL (2R,3S,4S)-5-[(4AS,10AR)-7,8-DIMETHYL-2,4-DIOXO-1,3,4,4A,5,10A-HEXAHYDROBENZO[G]PTERIDIN-10(2H)-YL]-2,3,4-TRIHYDROXPENTYL DIHYDROGEN DIPHOSPHATE (three-letter code: FAO) (formula: C₂₇H₃₇N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	X	0	0
			2	2		
4	A	2	Total	X	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	484	Total	O	0	0
			484	484		
5	B	485	Total	O	0	0
			485	485		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CHOLINE OXIDASE



• Molecule 1: CHOLINE OXIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.36 Å 84.36 Å 343.45 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.86 41.30 – 1.86	Depositor EDS
% Data completeness (in resolution range)	92.6 (50.00-1.86) 92.6 (41.30-1.86)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 1.87 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.169 , 0.212 0.171 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.533	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 48.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 97325 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9252	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, DMS, FAO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/4179	0.75	7/5690 (0.1%)
1	B	0.66	0/4191	0.69	3/5702 (0.1%)
All	All	0.67	0/8370	0.72	10/11392 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	86	ARG	NE-CZ-NH2	-7.87	116.36	120.30
1	A	484	ASP	CB-CG-OD1	6.30	123.97	118.30
1	B	404	ARG	NE-CZ-NH2	-6.27	117.17	120.30
1	A	86	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	A	157	LEU	CA-CB-CG	5.38	127.67	115.30
1	B	350	MET	CG-SD-CE	-5.28	91.76	100.20
1	A	404	ARG	NE-CZ-NH2	-5.21	117.69	120.30
1	A	492	VAL	CG1-CB-CG2	5.15	119.14	110.90
1	B	396	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	A	11	ASP	CB-CG-OD1	5.09	122.88	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	238	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4071	0	3910	57	0
1	B	4094	0	3929	43	0
2	A	53	0	33	2	0
2	B	53	0	32	2	0
3	A	4	0	6	0	0
3	B	4	0	6	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	484	0	0	16	0
5	B	485	0	0	10	0
All	All	9252	0	7916	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:230:LEU:HD13	1:B:248:ILE:HD12	1.18	1.14
1:A:238:ASP:HB3	1:A:239:ALA:HB3	1.30	1.10
1:B:230:LEU:HD13	1:B:248:ILE:CD1	1.92	0.99
1:A:1:MET:HE3	1:A:3:ILE:HD11	1.47	0.96
1:A:238:ASP:CB	1:A:239:ALA:HB3	1.98	0.94
1:A:288:HIS:HE1	1:A:390:LEU:H	1.16	0.89
1:A:1:MET:CE	1:A:3:ILE:HD11	2.07	0.84
1:B:37:ALA:C	1:B:38:VAL:CA	2.45	0.84
1:A:383:ARG:CZ	5:A:2372:HOH:O	2.24	0.84
1:B:57:GLN:OE1	5:B:2041:HOH:O	1.96	0.83
1:A:288:HIS:CE1	1:A:390:LEU:H	1.98	0.81
1:A:383:ARG:NE	5:A:2372:HOH:O	2.16	0.79
1:A:238:ASP:HB3	1:A:240:ASP:H	1.49	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80:ASN:OD1	1:B:458:ARG:NH1	2.20	0.74
1:B:291:GLU:HG3	5:B:2274:HOH:O	1.86	0.74
1:B:195:ASN:HD22	1:B:196:PHE:H	1.35	0.73
1:B:511:PRO:HD2	2:B:1531:FAO:H1	1.55	0.71
1:A:156:HIS:HE1	5:A:2182:HOH:O	1.73	0.71
1:A:195:ASN:HD22	1:A:196:PHE:H	1.38	0.69
1:A:511:PRO:HD2	2:A:1528:FAO:H1	1.57	0.69
1:B:267:LEU:HD13	1:B:274:THR:HG23	1.74	0.69
1:B:238:ASP:HB3	1:B:240:ASP:H	1.57	0.69
1:B:234:GLN:NE2	5:B:2247:HOH:O	2.26	0.68
1:B:527:ARG:HB2	5:B:2477:HOH:O	1.96	0.66
1:B:394:ASP:OD1	1:B:396:ARG:HD2	1.97	0.65
1:A:383:ARG:NH2	5:A:2372:HOH:O	2.29	0.64
1:A:455:ASP:OD1	1:A:458:ARG:NH2	2.32	0.62
1:B:486:GLU:HG3	1:B:488:ARG:CZ	2.29	0.62
1:A:238:ASP:HB3	1:A:239:ALA:CB	2.20	0.62
1:A:238:ASP:HB3	1:A:240:ASP:N	2.14	0.61
1:A:1:MET:CE	5:A:2248:HOH:O	2.49	0.60
1:A:137:GLU:OE1	1:A:149:HIS:HD2	1.86	0.58
1:A:239:ALA:HB2	5:A:2260:HOH:O	2.02	0.58
1:B:156:HIS:HE1	5:B:2173:HOH:O	1.87	0.57
1:B:195:ASN:HD22	1:B:196:PHE:N	2.03	0.57
1:A:255:HIS:CE1	1:B:370:GLU:OE1	2.57	0.57
1:B:412:HIS:HE1	5:B:2119:HOH:O	1.89	0.56
1:A:359:MET:HB2	1:B:253:PHE:HZ	1.71	0.55
1:B:484:ASP:HB2	1:B:485:PRO:CD	2.37	0.55
1:B:410:GLU:OE2	1:B:415:ARG:NH1	2.39	0.55
1:A:415:ARG:HD3	5:A:2408:HOH:O	2.05	0.55
1:A:241:ARG:HG2	1:A:297:LEU:HD11	1.88	0.54
1:A:260:THR:HG22	5:A:2265:HOH:O	2.08	0.54
1:A:149:HIS:HE1	5:A:2163:HOH:O	1.91	0.53
1:B:2:HIS:HE1	1:B:228:THR:OG1	1.92	0.53
1:A:1:MET:HE1	5:A:2248:HOH:O	2.07	0.53
1:B:241:ARG:HD3	1:B:297:LEU:HG	1.90	0.53
1:B:394:ASP:OD1	1:B:396:ARG:CD	2.57	0.52
1:A:329:GLN:NE2	1:A:331:TRP:H	2.06	0.52
1:A:8:ASN:ND2	5:A:2009:HOH:O	1.57	0.52
1:B:240:ASP:O	1:B:241:ARG:HB2	2.10	0.51
1:A:191:VAL:HA	1:A:346:PRO:HD3	1.92	0.50
1:A:2:HIS:HE1	1:A:228:THR:OG1	1.94	0.50
1:A:422:ARG:NH2	1:A:450:ASP:OD2	2.43	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:450:ASP:O	1:A:454:GLN:HG3	2.12	0.49
1:B:230:LEU:CD1	1:B:248:ILE:HD12	2.13	0.48
1:A:364:HIS:HD2	5:A:2360:HOH:O	1.97	0.48
1:B:35:ASP:O	1:B:38:VAL:HG22	2.14	0.48
1:A:253:PHE:HZ	1:B:359:MET:HE2	1.79	0.47
1:B:306:HIS:HD2	5:B:2287:HOH:O	1.97	0.47
1:A:57:GLN:HG2	5:A:2235:HOH:O	2.13	0.47
1:A:358:ASP:OD2	1:B:396:ARG:NH2	2.28	0.47
1:A:195:ASN:HD22	1:A:196:PHE:N	2.08	0.46
1:B:412:HIS:HD2	1:B:413:ASP:OD1	1.99	0.46
1:A:133:TYR:O	1:A:137:GLU:HG2	2.15	0.46
1:A:157:LEU:HD22	1:A:200:ASN:HB3	1.96	0.46
1:B:133:TYR:O	1:B:137:GLU:HG2	2.16	0.46
1:B:512:ASN:HB2	2:B:1531:FAO:C2	2.46	0.45
1:A:404:ARG:HD2	5:A:2095:HOH:O	2.16	0.45
1:B:24:ALA:O	1:B:28:VAL:HG23	2.16	0.45
1:A:1:MET:HE2	5:A:2248:HOH:O	2.15	0.45
1:B:362:LEU:HD12	5:B:2340:HOH:O	2.17	0.45
1:A:137:GLU:OE1	1:A:149:HIS:CD2	2.69	0.45
1:B:306:HIS:HE1	5:B:2182:HOH:O	2.00	0.44
1:A:483:LEU:CD1	1:A:502:VAL:HG12	2.47	0.44
1:B:329:GLN:NE2	1:B:331:TRP:H	2.14	0.44
1:A:267:LEU:HD11	1:A:277:LEU:HD23	1.99	0.44
1:B:486:GLU:HG3	1:B:488:ARG:NH2	2.34	0.43
1:A:459:LYS:HB3	1:A:459:LYS:HE2	1.90	0.43
1:A:383:ARG:HD2	1:A:412:HIS:CG	2.54	0.42
1:A:238:ASP:HB2	1:A:242:ARG:H	1.84	0.42
1:A:1:MET:HE3	1:A:3:ILE:CD1	2.35	0.42
1:A:329:GLN:HE22	1:A:331:TRP:C	2.23	0.42
1:A:230:LEU:HD13	1:A:248:ILE:HD12	2.00	0.42
1:A:383:ARG:HH11	1:A:412:HIS:CD2	2.37	0.42
1:A:445:VAL:O	1:A:448:GLN:NE2	2.52	0.42
1:B:157:LEU:HD22	1:B:200:ASN:HB3	2.02	0.42
1:A:96:CYS:SG	2:A:1528:FAO:H8M1	2.60	0.41
1:A:321:GLN:HB2	1:A:436:TRP:CZ3	2.56	0.41
1:A:227:LEU:HB3	1:A:230:LEU:HD12	2.03	0.41
1:A:202:ARG:HD2	5:A:2227:HOH:O	2.21	0.41
1:B:191:VAL:HA	1:B:346:PRO:HD3	2.03	0.41
1:B:529:ALA:O	1:B:530:ARG:HB2	2.21	0.40
1:B:180:PRO:O	1:B:193:GLY:HA2	2.21	0.40
1:B:291:GLU:CG	5:B:2274:HOH:O	2.57	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:80:ASN:OD1	1:A:458:ARG:HD3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/546 (96%)	511 (97%)	14 (3%)	2 (0%)	39	22
1	B	526/546 (96%)	507 (96%)	18 (3%)	1 (0%)	52	36
All	All	1053/1092 (96%)	1018 (97%)	32 (3%)	3 (0%)	46	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	238	ASP
1	A	465	TYR
1	B	92	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	435/446 (98%)	425 (98%)	10 (2%)	58	41
1	B	435/446 (98%)	420 (97%)	15 (3%)	44	24

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	870/892 (98%)	845 (97%)	25 (3%)	50 31

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	HIS
1	A	130	TRP
1	A	156	HIS
1	A	171	LEU
1	A	195	ASN
1	A	208	SER
1	A	329	GLN
1	A	492	VAL
1	A	506	HIS
1	A	521	ARG
1	B	99	HIS
1	B	130	TRP
1	B	156	HIS
1	B	171	LEU
1	B	195	ASN
1	B	202	ARG
1	B	226	LEU
1	B	267	LEU
1	B	329	GLN
1	B	396	ARG
1	B	410	GLU
1	B	459	LYS
1	B	486	GLU
1	B	492	VAL
1	B	506	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	HIS
1	A	149	HIS
1	A	195	ASN
1	A	288	HIS
1	A	321	GLN
1	A	329	GLN
1	A	364	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	412	HIS
1	B	2	HIS
1	B	8	ASN
1	B	57	GLN
1	B	192	ASN
1	B	195	ASN
1	B	292	HIS
1	B	306	HIS
1	B	329	GLN
1	B	364	HIS
1	B	412	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are unknown - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAO	A	1528	1,4	50,58,58	1.41	6 (12%)	53,89,89	1.59	6 (11%)
3	DMS	A	1530	-	3,3,3	0.55	0	3,3,3	0.91	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAO	B	1531	1,4	50,58,58	1.37	6 (12%)	53,89,89	1.75	5 (9%)
3	DMS	B	1533	-	3,3,3	0.57	0	3,3,3	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAO	A	1528	1,4	-	0/30/78/78	0/6/6/6
3	DMS	A	1530	-	-	0/0/0/0	0/0/0/0
2	FAO	B	1531	1,4	-	0/30/78/78	0/6/6/6
3	DMS	B	1533	-	-	0/0/0/0	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1528	FAO	C4N-N5	2.08	1.48	1.45
2	B	1531	FAO	C8M-C8	2.10	1.55	1.51
2	B	1531	FAO	C7M-C7	2.11	1.55	1.51
2	A	1528	FAO	C8M-C8	2.57	1.56	1.51
2	B	1531	FAO	O4A'-C1A'	2.78	1.44	1.41
2	A	1528	FAO	C10-N10	2.82	1.47	1.45
2	B	1531	FAO	C4N-N5	2.85	1.49	1.45
2	A	1528	FAO	C1'-C2'	3.46	1.57	1.52
2	A	1528	FAO	C4N-C4	3.53	1.59	1.53
2	B	1531	FAO	C4N-C4	3.77	1.59	1.53
2	B	1531	FAO	C4-N3	3.92	1.44	1.37
2	A	1528	FAO	C4-N3	4.28	1.44	1.37

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1531	FAO	N3A-C2A-N1A	-10.38	120.95	128.89
2	A	1528	FAO	N3A-C2A-N1A	-8.58	122.33	128.89
2	B	1531	FAO	C4A-C5A-N7A	-2.47	107.21	109.48
2	B	1531	FAO	C4A'-O4A'-C1A'	-2.36	107.13	109.72
2	A	1528	FAO	O4-C4-N3	-2.29	116.62	120.93
2	B	1531	FAO	C1A'-N9A-C4A	-2.27	123.52	126.94
2	B	1531	FAO	C5A'-C4A'-C3A'	-2.07	106.99	115.21
2	A	1528	FAO	C4A-C5A-N7A	-2.02	107.62	109.48
2	A	1528	FAO	O3A'-C3A'-C4A'	-2.02	105.00	111.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1528	FAO	O2A-PA-O3P	2.09	114.59	105.09
2	A	1528	FAO	C5N-C9N-N10	2.10	122.18	119.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1528	FAO	2	0
2	B	1531	FAO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	527/546 (96%)	-0.19	9 (1%) 73 72	11, 19, 30, 37	0
1	B	530/546 (97%)	-0.01	18 (3%) 49 46	13, 21, 42, 63	0
All	All	1057/1092 (96%)	-0.10	27 (2%) 59 57	11, 20, 35, 63	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	8	ASN	5.6
1	B	10	SER	5.3
1	B	9	LEU	4.5
1	B	11	ASP	4.4
1	B	7	GLU	3.8
1	B	527	ARG	3.5
1	A	446	GLU	3.2
1	B	5	ASN	3.1
1	B	1	MET	2.7
1	A	4	ASP	2.5
1	B	239	ALA	2.5
1	B	38	VAL	2.4
1	B	51	GLY	2.4
1	A	326	GLU	2.4
1	B	36	PRO	2.4
1	A	163	LYS	2.4
1	A	325	ALA	2.4
1	A	1	MET	2.4
1	B	477	GLU	2.3
1	A	239	ALA	2.3
1	B	530	ARG	2.3
1	B	37	ALA	2.3
1	B	163	LYS	2.3
1	A	445	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	250	ASP	2.2
1	A	479	GLU	2.1
1	B	13	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	UNX	B	2501	1/1	0.98	0.26	21.75	26,26,26,26	1
4	UNX	A	2501	1/1	0.90	0.28	15.51	27,27,27,27	1
3	DMS	B	1533	4/4	0.94	0.12	3.34	41,41,41,42	0
4	UNX	B	2500	1/1	0.94	0.16	2.91	24,24,24,24	0
2	FAO	B	1531	53/53	0.97	0.12	0.21	14,19,25,25	0
3	DMS	A	1530	4/4	0.97	0.10	0.19	37,37,38,39	0
2	FAO	A	1528	53/53	0.98	0.10	-0.22	10,14,19,25	0
4	UNX	A	2500	1/1	0.96	0.17	-	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.