



wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:55 PM BST

PDB ID : 3JB7
EMDB ID: : EMD-6404
Title : In situ structures of the segmented genome and RNA polymerase complex inside a dsRNA virus
Authors : Zhang, X.; Ding, K.; Yu, X.K.; Chang, W.; Sun, J.C.; Zhou, Z.H.
Deposited on : 2015-08-03
Resolution : 4.00 Å(reported)

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/EMValidationReportHelp>

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk27241


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Clashscore | 114402 | 924 |
| Ramachandran outliers | 111179 | 726 |
| Sidechain outliers | 111093 | 686 |
| RNA backbone | 3027 | 244 |

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 1225 | 77% 21% . |
| 2 | B | 561 | 69% 20% 11% |
| 3 | C | 24 | 8% 50% 25% . 13% |
| 3 | D | 24 | 25% 38% . 33% |
| 4 | t | 6 | 83% 17% |
| 5 | m | 5 | 100% |

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14176 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CPV RNA-dependent RNA polymerase.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|------|------|----|---------|-------|
| 1 | A | 1198 | Total | C | N | O | S | 0 | 0 |
| | | | 9561 | 6066 | 1654 | 1805 | 36 | | |

- Molecule 2 is a protein called Viral structural protein 4.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 2 | B | 498 | Total | C | N | O | S | 0 | 0 |
| | | | 3985 | 2553 | 666 | 749 | 17 | | |

- Molecule 3 is a protein called VP1 CSP.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 3 | C | 21 | Total | C | N | O | S | 0 | 0 |
| | | | 170 | 105 | 30 | 34 | 1 | | |
| 3 | D | 16 | Total | C | N | O | | 0 | 0 |
| | | | 129 | 81 | 22 | 26 | | | |


- Molecule 4 is a RNA chain called RNA (5'-R(P*GP*GP*GP*GP*G)-3').

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|-------|
| 4 | t | 6 | Total | C | N | O | P | 0 | 0 |
| | | | 138 | 60 | 30 | 42 | 6 | | |

- Molecule 5 is a RNA chain called RNA (5'-R(P*CP*CP*CP*CP*C)-3').


| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|----|----|----|---|---------|-------|
| 5 | m | 5 | Total | C | N | O | P | 0 | 0 |
| | | | 100 | 45 | 15 | 35 | 5 | | |

- Molecule 6 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).




- # CTP


| Mol | Chain | Residues | Atoms | | | | | AltConf |
|-----|-------|----------|-------|---|---|----|---|---------|
| 7 | A | 1 | Total | C | N | O | P | 0 |
| | | | 29 | 9 | 3 | 14 | 3 | |




- Molecule 3: VP1 CSP



- Molecule 3: VP1 CSP



- Molecule 4: RNA (5'-R(P*GP*GP*GP*GP*GP*G)-3')



- Molecule 5: RNA (5'-R(P*CP*CP*CP*CP*C)-3')



There are no outlier residues recorded for this chain.

4 Experimental information

| Property | Value | Source |
|--------------------------------------|---------------------|-----------|
| Reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, Not provided | Depositor |
| Number of images | 81887 | Depositor |
| Resolution determination method | FSC 0.143 | Depositor |
| CTF correction method | Each particle | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 40 | Depositor |
| Minimum defocus (nm) | 900 | Depositor |
| Maximum defocus (nm) | 3400 | Depositor |
| Magnification | 105000 | Depositor |
| Image detector | GATAN K2 (4k x 4k) | Depositor |

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, CTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|-----------------|-------------|----------------|
| | | RMSZ | $\# Z > 2$ | RMSZ | $\# Z > 2$ |
| 1 | A | 0.26 | 0/9763 | 0.46 | 0/13213 |
| 2 | B | 0.26 | 0/4071 | 0.44 | 0/5519 |
| 3 | C | 2.25 | 12/173 (6.9%) | 1.72 | 8/234 (3.4%) |
| 3 | D | 2.19 | 2/131 (1.5%) | 0.77 | 1/177 (0.6%) |
| 4 | t | 0.19 | 0/155 | 0.68 | 0/242 |
| 5 | m | 0.14 | 0/109 | 0.73 | 0/166 |
| All | All | 0.41 | 14/14402 (0.1%) | 0.50 | 9/19551 (0.0%) |

The worst 5 of 14 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|--------|--------|-------------|----------|
| 3 | D | 111 | PRO | N-CA | -20.85 | 1.11 | 1.47 |
| 3 | D | 111 | PRO | N-CD | 11.71 | 1.64 | 1.47 |
| 3 | C | 125 | PHE | CG-CD1 | -7.89 | 1.26 | 1.38 |
| 3 | C | 125 | PHE | CA-CB | -7.41 | 1.37 | 1.53 |
| 3 | C | 132 | PRO | C-O | -7.05 | 1.09 | 1.23 |

The worst 5 of 9 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|------------|-------|-------------|----------|
| 3 | C | 120 | VAL | CG1-CB-CG2 | -9.75 | 95.30 | 110.90 |
| 3 | C | 130 | LEU | N-CA-C | 8.43 | 133.76 | 111.00 |
| 3 | C | 130 | LEU | CB-CG-CD2 | -8.15 | 97.14 | 111.00 |
| 3 | C | 131 | HIS | C-N-CD | 7.12 | 143.36 | 128.40 |
| 3 | D | 111 | PRO | N-CD-CG | -6.77 | 93.05 | 103.20 |

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 9561 | 0 | 9465 | 174 | 0 |
| 2 | B | 3985 | 0 | 3984 | 97 | 0 |
| 3 | C | 170 | 0 | 155 | 46 | 0 |
| 3 | D | 129 | 0 | 121 | 17 | 0 |
| 4 | t | 138 | 0 | 67 | 0 | 0 |
| 5 | m | 100 | 0 | 56 | 0 | 0 |
| 6 | A | 32 | 0 | 12 | 0 | 0 |
| 6 | B | 32 | 0 | 12 | 3 | 0 |
| 7 | A | 29 | 0 | 12 | 1 | 0 |
| All | All | 14176 | 0 | 13884 | 283 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 283 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|------------------|-----------------|--------------------------|-------------------|
| 1:A:1193:ARG:HD3 | 3:D:116:SER:CB | 1.36 | 1.51 |
| 2:B:518:ARG:HH11 | 3:C:115:GLN:NE2 | 1.11 | 1.47 |
| 1:A:1193:ARG:CD | 3:D:116:SER:CB | 2.16 | 1.23 |
| 1:A:1193:ARG:HD3 | 3:D:116:SER:OG | 1.06 | 1.21 |
| 2:B:518:ARG:NH1 | 3:C:115:GLN:NE2 | 1.92 | 1.15 |

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|---------|----------|-------------|-----|
| 1 | A | 1194/1225 (98%) | 1140 (96%) | 51 (4%) | 3 (0%) | 46 | 82 |
| 2 | B | 492/561 (88%) | 477 (97%) | 15 (3%) | 0 | 100 | 100 |
| 3 | C | 19/24 (79%) | 17 (90%) | 1 (5%) | 1 (5%) | 2 | 31 |
| 3 | D | 14/24 (58%) | 14 (100%) | 0 | 0 | 100 | 100 |
| All | All | 1719/1834 (94%) | 1648 (96%) | 67 (4%) | 4 (0%) | 56 | 86 |

All (4) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 928 | VAL |
| 1 | A | 97 | GLN |
| 3 | C | 119 | ASP |
| 1 | A | 581 | PRO |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|-------------|----------|-------------|----|
| 1 | A | 1039/1063 (98%) | 1035 (100%) | 4 (0%) | 93 | 97 |
| 2 | B | 434/493 (88%) | 432 (100%) | 2 (0%) | 92 | 96 |
| 3 | C | 19/22 (86%) | 15 (79%) | 4 (21%) | 1 | 11 |
| 3 | D | 15/22 (68%) | 14 (93%) | 1 (7%) | 20 | 60 |
| All | All | 1507/1600 (94%) | 1496 (99%) | 11 (1%) | 89 | 94 |

5 of 11 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 2 | B | 510 | ILE |
| 2 | B | 544 | THR |
| 3 | C | 128 | GLU |
| 1 | A | 1170 | SER |
| 3 | C | 122 | ASN |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such

sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 481 | GLN |
| 3 | C | 115 | GLN |

5.3.3 RNA [i](#)

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|------------|-------------------|-----------------|
| 4 | t | 5/6 (83%) | 1 (20%) | 0 |
| 5 | m | 4/5 (80%) | 0 | 0 |
| All | All | 9/11 (81%) | 1 (11%) | 0 |

All (1) RNA backbone outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 4 | t | 5 | G |

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|------|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 6 | GTP | A | 1301 | - | 26,34,34 | 0.92 | 1 (3%) | 29,54,54 | 1.57 | 3 (10%) |
| 7 | CTP | A | 1302 | - | 23,30,30 | 0.77 | 0 | 28,47,47 | 1.14 | 2 (7%) |
| 6 | GTP | B | 601 | - | 26,34,34 | 0.90 | 1 (3%) | 29,54,54 | 1.51 | 4 (13%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|------|------|---------|------------|---------|
| 6 | GTP | A | 1301 | - | - | 0/18/38/38 | 0/3/3/3 |
| 7 | CTP | A | 1302 | - | - | 0/18/38/38 | 0/2/2/2 |
| 6 | GTP | B | 601 | - | - | 0/18/38/38 | 0/3/3/3 |

All (2) bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|------|------|-------|------|-------------|----------|
| 6 | B | 601 | GTP | C6-N1 | 2.84 | 1.38 | 1.33 |
| 6 | A | 1301 | GTP | C6-N1 | 2.93 | 1.38 | 1.33 |

The worst 5 of 9 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|------|------|----------|-------|-------------|----------|
| 6 | B | 601 | GTP | N3-C2-N1 | -5.43 | 120.17 | 127.56 |
| 6 | A | 1301 | GTP | N3-C2-N1 | -5.41 | 120.20 | 127.56 |
| 6 | A | 1301 | GTP | C5-C6-N1 | -2.91 | 119.72 | 123.52 |
| 6 | B | 601 | GTP | C5-C6-N1 | -2.68 | 120.02 | 123.52 |
| 7 | A | 1302 | CTP | C5-C4-N3 | -2.52 | 118.59 | 121.79 |

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|------|------|---------|--------------|
| 7 | A | 1302 | CTP | 1 | 0 |
| 6 | B | 601 | GTP | 3 | 0 |

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.