



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:43 PM GMT

PDB ID : 4JBC
Title : Crystal Structure of the computationally designed serine hydrolase 3mmj_2, Northeast Structural Genomics Consortium (NESG) Target OR318
Authors : Kuzin, A.; Lew, S.; Rajagopalan, S.; Seetharaman, J.; Maglaqui, M.; Xiao, R.; Lee, D.; Everett, J.K.; Acton, T.B.; Montelione, G.T.; Tong, L.; Baker, D.; Hunt, J.F.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2013-02-19
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	<div> <div>4%</div> <div>82%</div> <div>12%</div> <div>• •</div> </div>
1	B	323	<div> <div>7%</div> <div>76%</div> <div>19%</div> <div>• •</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called designed serine hydrolase 3mmj_2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	Se	0	0	0
			2543	1621	455	461	6			
1	B	313	Total	C	N	O	Se	0	0	0
			2534	1616	453	459	6			

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	272	Total 272	O 272	0	0
3	B	238	Total 238	O 238	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: designed serine hydrolase 3mmj_2



- Molecule 1: designed serine hydrolase 3mmj_2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.24Å 194.55Å 46.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.73 – 2.00 29.73 – 2.01	Depositor EDS
% Data completeness (in resolution range)	84.5 (29.73-2.00) 84.7 (29.73-2.01)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.83 (at 2.01Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1269)	Depositor
R, R_{free}	0.202 , 0.261 0.202 , 0.260	Depositor DCC
R_{free} test set	1988 reflections (4.65%)	DCC
Wilson B-factor (Å ²)	15.8	Xtriage
Anisotropy	0.745	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	24 of 42898 reflections (0.056%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5607	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2611	0.60	0/3529
1	B	0.41	0/2602	0.57	0/3517
All	All	0.42	0/5213	0.59	0/7046

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2543	0	2468	32	0
1	B	2534	0	2460	36	0
2	A	5	0	0	0	0
2	B	15	0	0	1	0
3	A	272	0	0	5	0
3	B	238	0	0	2	0
All	All	5607	0	4928	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:41:THR:HG22	1:B:43:ALA:H	1.45	0.81
1:A:234:MSE:HE2	1:A:288:PHE:CG	2.20	0.75
1:A:234:MSE:HE2	1:A:288:PHE:CD1	2.27	0.68
1:A:201:ILE:HD13	1:A:234:MSE:HE1	1.74	0.68
1:A:173:THR:HG23	1:A:176:GLU:H	1.58	0.67
1:A:110:LEU:HD22	1:A:115:VAL:HG21	1.77	0.66
1:B:173:THR:HG23	1:B:176:GLU:H	1.60	0.64
1:A:77:PHE:CE1	1:A:82:LEU:HG	2.33	0.64
1:B:110:LEU:HD22	1:B:115:VAL:HG21	1.80	0.62
1:B:260:TYR:HE2	1:B:281:LYS:HD2	1.64	0.61
1:A:3:THR:HG22	1:A:4:VAL:HA	1.83	0.59
1:B:32:LYS:NZ	3:B:569:HOH:O	2.35	0.59
1:A:298:ASP:HB2	1:A:300:TYR:N	2.18	0.58
1:A:195:TRP:CZ2	1:A:234:MSE:HE3	2.39	0.57
1:B:237:MSE:HG2	1:B:248:ILE:HD13	1.86	0.57
1:A:195:TRP:CH2	1:A:234:MSE:HE3	2.41	0.56
1:A:41:THR:HG22	1:A:43:ALA:H	1.70	0.56
1:B:249:LEU:HD21	1:B:285:ILE:HD13	1.87	0.56
1:B:71:ILE:HD11	1:B:235:THR:HB	1.87	0.55
1:A:298:ASP:HB2	1:A:300:TYR:H	1.72	0.55
1:B:287:GLN:HG3	1:B:313:ALA:HB2	1.88	0.53
1:A:262:GLU:HA	1:A:282:ILE:HD13	1.91	0.52
1:B:314:LYS:HB2	1:B:315:ALA:HB2	1.91	0.51
1:A:41:THR:CG2	1:A:43:ALA:H	2.23	0.51
1:B:195:TRP:HZ3	1:B:285:ILE:HG22	1.77	0.50
1:B:287:GLN:OE1	1:B:290:ARG:NH1	2.45	0.49
1:B:223:SER:OG	2:B:401:PO4:O1	2.30	0.49
1:B:66:MSE:SE	1:B:251:ARG:HG2	2.63	0.49
1:A:121:ARG:NH2	3:A:713:HOH:O	2.29	0.49
1:B:195:TRP:CZ3	1:B:285:ILE:HG22	2.49	0.48
1:A:77:PHE:CD1	1:A:82:LEU:HG	2.51	0.46
1:A:209:ARG:NH2	3:A:772:HOH:O	2.41	0.46
1:B:260:TYR:CE2	1:B:281:LYS:HD2	2.45	0.46
1:A:33:GLU:HG2	1:A:37:ARG:NH2	2.31	0.46
1:A:2:GLN:HB3	1:A:3:THR:H	1.55	0.45
1:B:164:GLU:OE1	1:B:166:ARG:NH1	2.49	0.45
1:A:201:ILE:CD1	1:A:234:MSE:HE1	2.45	0.45
1:A:298:ASP:HB3	1:A:299:GLY:HA3	1.99	0.45
1:A:66:MSE:HE2	1:A:69:LEU:HD12	1.99	0.45
1:A:69:LEU:HD13	1:A:71:ILE:HG12	1.98	0.45
1:A:26:ARG:HG3	1:A:154:LEU:HG	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:ASP:HB3	1:A:274:LYS:HE3	2.00	0.44
1:A:206:ALA:O	1:A:210:THR:HG23	2.17	0.44
1:B:109:TYR:HB2	1:B:172:GLN:HB2	2.00	0.43
1:B:195:TRP:CD2	1:B:196:PRO:HD2	2.54	0.43
1:B:156:LYS:HG3	1:B:157:HIS:CD2	2.53	0.43
1:B:206:ALA:O	1:B:210:THR:HG23	2.19	0.43
1:A:139:HIS:HB3	3:A:691:HOH:O	2.19	0.42
1:B:269:ASP:N	1:B:269:ASP:OD1	2.50	0.42
1:B:167:ARG:NE	3:B:600:HOH:O	2.40	0.42
1:B:175:GLN:O	1:B:179:GLU:HG3	2.20	0.42
1:B:147:LYS:HD3	1:B:147:LYS:HA	1.80	0.41
1:B:205:LEU:CD2	1:B:238:LEU:HD21	2.50	0.41
1:A:2:GLN:OE1	1:A:3:THR:OG1	2.24	0.41
1:B:77:PHE:CE1	1:B:82:LEU:HG	2.55	0.41
1:A:263:PHE:O	1:A:265:ILE:N	2.54	0.41
1:B:195:TRP:CE2	1:B:284:MSE:HB3	2.55	0.41
1:B:89:LEU:HD22	1:B:216:TRP:CE2	2.55	0.41
1:A:174:ASN:ND2	3:A:543:HOH:O	2.54	0.41
1:B:41:THR:HB	1:B:44:ASP:OD1	2.20	0.41
1:B:314:LYS:HA	1:B:315:ALA:HA	1.79	0.41
1:B:298:ASP:O	1:B:301:GLN:HG2	2.20	0.41
1:A:199:GLU:HG3	3:A:680:HOH:O	2.21	0.41
1:B:42:SER:O	1:B:66:MSE:HE2	2.22	0.40
1:B:243:VAL:CG2	1:B:248:ILE:HG13	2.50	0.40
1:A:221:SER:OG	1:A:228:THR:HG23	2.21	0.40
1:B:27:LEU:HD13	1:B:151:ILE:HD11	2.02	0.40
1:B:151:ILE:O	1:B:151:ILE:HG13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/323 (97%)	302 (97%)	9 (3%)	1 (0%)	46	41
1	B	311/323 (96%)	297 (96%)	13 (4%)	1 (0%)	46	41
All	All	623/646 (96%)	599 (96%)	22 (4%)	2 (0%)	46	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	264	PRO
1	B	264	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/263 (99%)	244 (94%)	17 (6%)	21	15
1	B	260/263 (99%)	238 (92%)	22 (8%)	13	8
All	All	521/526 (99%)	482 (92%)	39 (8%)	17	11

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	4	VAL
1	A	26	ARG
1	A	41	THR
1	A	69	LEU
1	A	82	LEU
1	A	90	ARG
1	A	110	LEU
1	A	141	LEU
1	A	151	ILE
1	A	165	VAL
1	A	217	LEU
1	A	243	VAL
1	A	262	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	271	ASP
1	A	308	LEU
1	A	314	LYS
1	B	4	VAL
1	B	26	ARG
1	B	47	ARG
1	B	52	LYS
1	B	69	LEU
1	B	71	ILE
1	B	77	PHE
1	B	82	LEU
1	B	90	ARG
1	B	110	LEU
1	B	120	GLU
1	B	130	GLN
1	B	141	LEU
1	B	151	ILE
1	B	158	LYS
1	B	217	LEU
1	B	225	VAL
1	B	238	LEU
1	B	245	LEU
1	B	268	LYS
1	B	269	ASP
1	B	308	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	401	-	4,4,4	0.46	0	6,6,6	0.28	0
2	PO4	B	401	-	4,4,4	0.44	0	6,6,6	0.27	0
2	PO4	B	402	-	4,4,4	0.48	0	6,6,6	0.27	0
2	PO4	B	403	-	4,4,4	0.44	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	401	-	-	0/0/0/0	0/0/0/0
2	PO4	B	401	-	-	0/0/0/0	0/0/0/0
2	PO4	B	402	-	-	0/0/0/0	0/0/0/0
2	PO4	B	403	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	PO4	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	308/323 (95%)	0.10	14 (4%) 37 38	10, 21, 54, 131	0
1	B	307/323 (95%)	0.21	21 (6%) 20 22	10, 23, 55, 86	0
All	All	615/646 (95%)	0.16	35 (5%) 27 29	10, 22, 55, 131	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	GLN	7.7
1	A	3	THR	7.7
1	B	315	ALA	7.0
1	B	299	GLY	4.8
1	A	157	HIS	4.1
1	B	3	THR	4.1
1	B	233	VAL	4.0
1	B	156	LYS	4.0
1	A	265	ILE	3.9
1	B	266	LYS	3.9
1	A	266	LYS	3.8
1	B	268	LYS	3.5
1	A	4	VAL	3.5
1	B	265	ILE	3.5
1	A	268	LYS	3.2
1	A	315	ALA	3.2
1	A	269	ASP	3.1
1	A	297	ALA	3.1
1	B	229	THR	2.8
1	B	267	THR	2.8
1	B	157	HIS	2.6
1	A	267	THR	2.6
1	B	230	ALA	2.6
1	B	54	HIS	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	4	VAL	2.5
1	A	156	LYS	2.5
1	B	297	ALA	2.4
1	B	227	ALA	2.4
1	A	296	ARG	2.3
1	A	314	LYS	2.3
1	B	231	PHE	2.2
1	B	222	HIS	2.2
1	B	301	GLN	2.2
1	B	58	ALA	2.2
1	B	228	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
2	PO4	A	401	5/5	0.83	0.20	1.98	57,61,65,66	0
2	PO4	B	401	5/5	0.86	0.23	1.06	64,66,69,73	0
2	PO4	B	403	5/5	0.72	0.26	-	93,95,98,98	0
2	PO4	B	402	5/5	0.92	0.22	-	65,67,69,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.