



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2016 – 01:41 PM EDT

PDB ID : 5JCN  
Title : Structure and catalytic mechanism of monodehydroascorbate reductase, MD-HAR, from *Oryza sativa* L. japonica  
Authors : Park, A.K.; Kim, H.W.  
Deposited on : 2016-04-15  
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	<div> <div>9%</div> <div>82%</div> <div>12%</div> <div>• 5%</div> </div>
1	B	451	<div> <div>8%</div> <div>80%</div> <div>12%</div> <div>•• 6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ASC	B	502	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Os09g0567300 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	0	0
			3248	2095	531	616	6			
1	B	425	Total	C	N	O	S	0	0	0
			3215	2073	526	610	6			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	HIS	-	expression tag	UNP Q652L6
A	-14	HIS	-	expression tag	UNP Q652L6
A	-13	HIS	-	expression tag	UNP Q652L6
A	-12	HIS	-	expression tag	UNP Q652L6
A	-11	HIS	-	expression tag	UNP Q652L6
A	-10	HIS	-	expression tag	UNP Q652L6
A	-9	ALA	-	expression tag	UNP Q652L6
A	-8	SER	-	expression tag	UNP Q652L6
A	-7	GLU	-	expression tag	UNP Q652L6
A	-6	ASN	-	expression tag	UNP Q652L6
A	-5	LEU	-	expression tag	UNP Q652L6
A	-4	TYR	-	expression tag	UNP Q652L6
A	-3	PHE	-	expression tag	UNP Q652L6
A	-2	GLN	-	expression tag	UNP Q652L6
A	-1	GLY	-	expression tag	UNP Q652L6
A	0	ALA	-	expression tag	UNP Q652L6
A	349	PHE	TYR	engineered mutation	UNP Q652L6
B	-15	HIS	-	expression tag	UNP Q652L6
B	-14	HIS	-	expression tag	UNP Q652L6
B	-13	HIS	-	expression tag	UNP Q652L6
B	-12	HIS	-	expression tag	UNP Q652L6
B	-11	HIS	-	expression tag	UNP Q652L6
B	-10	HIS	-	expression tag	UNP Q652L6
B	-9	ALA	-	expression tag	UNP Q652L6
B	-8	SER	-	expression tag	UNP Q652L6

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	GLU	-	expression tag	UNP Q652L6
B	-6	ASN	-	expression tag	UNP Q652L6
B	-5	LEU	-	expression tag	UNP Q652L6
B	-4	TYR	-	expression tag	UNP Q652L6
B	-3	PHE	-	expression tag	UNP Q652L6
B	-2	GLN	-	expression tag	UNP Q652L6
B	-1	GLY	-	expression tag	UNP Q652L6
B	0	ALA	-	expression tag	UNP Q652L6
B	349	PHE	TYR	engineered mutation	UNP Q652L6

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is ASCORBIC ACID (three-letter code: ASC) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		
4	B	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	58	Total 58	O 58	0	0
5	B	51	Total 51	O 51	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Os09g0567300 protein



#### • Molecule 1: Os09g0567300 protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.64Å 85.11Å 133.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.29 43.84 – 2.29	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.29) 99.3 (43.84-2.29)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.205 , 0.264 0.230 , 0.278	Depositor DCC
$R_{free}$ test set	1934 reflections (4.71%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.1	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6790	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ASC, FAD, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.78	0/3317	0.93	6/4487 (0.1%)
1	B	0.74	0/3283	0.91	6/4440 (0.1%)
All	All	0.76	0/6600	0.92	12/8927 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	63	ARG	NE-CZ-NH2	-11.78	114.41	120.30
1	A	63	ARG	NE-CZ-NH1	11.44	126.02	120.30
1	B	63	ARG	NE-CZ-NH2	-6.77	116.92	120.30
1	B	132	ASP	CB-CG-OD1	6.31	123.98	118.30
1	B	63	ARG	NE-CZ-NH1	6.17	123.39	120.30
1	B	372	ASP	C-N-CA	5.74	136.06	121.70
1	A	372	ASP	C-N-CA	5.56	135.60	121.70
1	B	373	SER	N-CA-CB	5.30	118.46	110.50
1	A	147	ARG	NE-CZ-NH1	5.27	122.93	120.30
1	A	373	SER	N-CA-CB	5.08	118.12	110.50
1	B	362	ASP	CB-CG-OD1	5.02	122.82	118.30
1	A	377	SER	CB-CA-C	5.00	119.61	110.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	376	THR	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3248	0	3249	32	0
1	B	3215	0	3209	28	0
2	A	53	0	31	1	0
2	B	53	0	31	3	0
3	A	44	0	26	1	0
3	B	44	0	26	3	0
4	A	12	0	6	1	0
4	B	12	0	6	1	0
5	A	58	0	0	1	0
5	B	51	0	0	0	0
All	All	6790	0	6584	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:ILE:HG23	1:A:179:LEU:HD12	1.48	0.95
1:B:331:LYS:O	1:B:335:SER:OG	2.00	0.78
1:A:287:GLU:OE2	1:A:291:PRO:HA	1.86	0.76
3:A:501:NAD:N1A	5:A:602:HOH:O	2.23	0.71
1:B:331:LYS:O	1:B:335:SER:CB	2.41	0.68
1:A:377:SER:HB3	1:A:378:ALA:HB2	1.83	0.61
1:A:200:MET:HB2	1:A:204:PHE:CD2	2.37	0.60
1:A:418:VAL:HG11	1:A:422:GLU:HB2	1.84	0.59
1:B:319:ALA:HB2	2:B:500:FAD:H5'2	1.85	0.58
1:A:175:ILE:HG23	1:A:179:LEU:CD1	2.29	0.58
1:B:372:ASP:HA	1:B:373:SER:HB2	1.85	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:404:GLU:O	1:A:407:ALA:HB3	2.04	0.57
1:A:344:TYR:CE2	1:A:346:PRO:HA	2.41	0.56
1:B:127:VAL:HG21	1:B:145:TYR:HB3	1.88	0.56
1:A:179:LEU:HD13	1:A:257:VAL:HG11	1.89	0.53
1:B:104:LYS:HD2	1:B:116:TYR:O	2.10	0.52
1:B:7:PHE:O	1:B:116:TYR:HA	2.11	0.51
1:B:331:LYS:O	1:B:335:SER:HB2	2.11	0.50
1:A:189:ASP:OD1	1:A:222:LYS:NZ	2.36	0.50
1:A:55:TYR:O	1:A:63:ARG:NH2	2.44	0.50
1:A:377:SER:HB3	1:A:378:ALA:CB	2.41	0.49
1:A:342:TYR:CE2	1:A:344:TYR:HB2	2.48	0.49
1:A:7:PHE:O	1:A:116:TYR:HA	2.12	0.49
1:A:123:THR:O	2:A:500:FAD:H8A	2.13	0.49
1:B:49:PRO:HB2	4:B:502:ASC:H5	1.93	0.49
1:A:4:GLU:HG3	1:A:113:THR:HG23	1.94	0.48
1:B:342:TYR:CE2	1:B:344:TYR:HB2	2.48	0.48
1:A:79:LEU:HB3	1:A:80:PRO:HD2	1.95	0.48
1:B:71:VAL:CG1	1:B:77:ARG:HG3	2.43	0.48
2:B:500:FAD:C4	3:B:501:NAD:C7N	2.92	0.48
1:A:19:GLY:HA3	1:A:69:VAL:HG21	1.95	0.48
1:B:142:ASN:HB2	1:B:240:VAL:HG12	1.96	0.48
1:A:377:SER:CB	1:A:378:ALA:HB2	2.45	0.47
1:B:31:LYS:HB2	1:B:34:GLU:OE1	2.14	0.47
1:B:372:ASP:HA	1:B:373:SER:CB	2.45	0.47
1:A:422:GLU:O	1:A:424:LEU:N	2.49	0.46
1:A:372:ASP:HA	1:A:373:SER:CB	2.46	0.46
1:A:377:SER:HB3	1:A:378:ALA:CA	2.46	0.46
1:A:175:ILE:CG2	1:A:179:LEU:HD12	2.35	0.46
1:B:37:ILE:HB	1:B:90:LEU:HD23	1.97	0.45
1:A:110:VAL:O	1:A:110:VAL:HG12	2.17	0.44
1:B:304:MET:CE	1:B:311:ARG:HD2	2.48	0.44
1:A:49:PRO:HB2	4:A:502:ASC:H5	1.98	0.44
1:A:162:LYS:HD2	1:A:254:ASP:HB3	1.99	0.44
1:B:174:TYR:HB3	3:B:501:NAD:C4N	2.47	0.44
1:B:340:VAL:HG12	1:B:341:GLU:O	2.17	0.44
1:B:334:GLU:CB	1:B:335:SER:HB2	2.48	0.44
1:A:304:MET:CE	1:A:344:TYR:HD2	2.30	0.43
1:A:377:SER:HB3	1:A:378:ALA:HA	2.00	0.43
1:B:55:TYR:O	1:B:63:ARG:NH2	2.51	0.43
1:A:215:TYR:CD1	1:A:355:LEU:HD11	2.54	0.43
1:B:209:ALA:O	1:B:213:GLU:HG3	2.19	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:174:TYR:CD2	3:B:501:NAD:C7N	3.02	0.42
1:A:418:VAL:HG11	1:A:422:GLU:CB	2.49	0.42
1:B:369:LEU:HD23	1:B:369:LEU:C	2.40	0.42
1:B:200:MET:N	1:B:201:PRO:CD	2.83	0.41
1:B:382:PHE:O	1:B:397:LEU:HD12	2.21	0.41
1:A:372:ASP:HA	1:A:373:SER:HB2	2.01	0.41
1:B:298:ASP:OD1	2:B:500:FAD:H5'1	2.20	0.41
1:A:287:GLU:HG3	1:A:294:TYR:CZ	2.56	0.41
1:B:106:LEU:HD11	1:B:119:LEU:HD13	2.02	0.41
1:B:265:LEU:HD23	1:B:265:LEU:HA	1.88	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	425/451 (94%)	392 (92%)	27 (6%)	6 (1%)	14	13
1	B	419/451 (93%)	387 (92%)	27 (6%)	5 (1%)	16	16
All	All	844/902 (94%)	779 (92%)	54 (6%)	11 (1%)	15	15

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	335	SER
1	A	289	SER
1	A	337	GLU
1	A	373	SER
1	B	373	SER
1	A	308	ASN
1	B	433	SER
1	A	268	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	334	GLU
1	A	423	GLU
1	B	305	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/353 (95%)	318 (95%)	16 (5%)	31	42
1	B	330/353 (94%)	303 (92%)	27 (8%)	14	17
All	All	664/706 (94%)	621 (94%)	43 (6%)	21	27

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	HIS
1	A	40	LYS
1	A	77	ARG
1	A	78	LEU
1	A	105	THR
1	A	127	VAL
1	A	136	GLN
1	A	144	LEU
1	A	185	ILE
1	A	249	SER
1	A	372	ASP
1	A	373	SER
1	A	422	GLU
1	A	424	LEU
1	A	426	LYS
1	A	429	LEU
1	B	6	HIS
1	B	30	VAL
1	B	31	LYS
1	B	40	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	84	SER
1	B	85	GLU
1	B	86	LYS
1	B	89	GLU
1	B	107	THR
1	B	108	SER
1	B	113	THR
1	B	144	LEU
1	B	154	LYS
1	B	184	LYS
1	B	192	MET
1	B	202	ARG
1	B	222	LYS
1	B	249	SER
1	B	304	MET
1	B	334	GLU
1	B	338	SER
1	B	341	GLU
1	B	372	ASP
1	B	388	LYS
1	B	429	LEU
1	B	433	SER
1	B	434	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	318	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	500	-	52,58,58	1.53	13 (25%)	52,89,89	2.31	12 (23%)
3	NAD	A	501	-	42,48,48	1.15	3 (7%)	46,73,73	2.24	8 (17%)
4	ASC	A	502	-	12,12,12	2.60	2 (16%)	16,17,17	4.38	10 (62%)
2	FAD	B	500	-	52,58,58	1.57	12 (23%)	52,89,89	2.32	13 (25%)
3	NAD	B	501	-	42,48,48	1.21	3 (7%)	46,73,73	2.23	8 (17%)
4	ASC	B	502	-	12,12,12	2.80	1 (8%)	16,17,17	3.66	6 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	500	-	-	0/30/50/50	0/6/6/6
3	NAD	A	501	-	-	0/22/62/62	0/5/5/5
4	ASC	A	502	-	-	0/6/22/22	0/1/1/1
2	FAD	B	500	-	-	0/30/50/50	0/6/6/6
3	NAD	B	501	-	-	0/22/62/62	0/5/5/5
4	ASC	B	502	-	-	0/6/22/22	0/1/1/1

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	FAD	C2-N3	-3.46	1.31	1.38
2	A	500	FAD	C2-N3	-3.29	1.31	1.38
2	A	500	FAD	C6-C5X	-3.21	1.36	1.41
2	B	500	FAD	C6-C5X	-2.92	1.37	1.41
2	A	500	FAD	C2B-C1B	-2.78	1.49	1.53
2	B	500	FAD	P-O2P	-2.63	1.43	1.55
2	A	500	FAD	PA-O2A	-2.45	1.44	1.55

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	FAD	PA-O2A	-2.45	1.44	1.55
2	A	500	FAD	C2-N1	-2.37	1.33	1.38
2	B	500	FAD	C5X-N5	-2.37	1.31	1.35
2	B	500	FAD	C4'-C3'	-2.36	1.48	1.53
2	B	500	FAD	C2-N1	-2.35	1.33	1.38
2	A	500	FAD	P-O2P	-2.34	1.45	1.55
2	B	500	FAD	C2B-C1B	-2.30	1.50	1.53
2	B	500	FAD	C10-N1	-2.22	1.31	1.35
2	A	500	FAD	C5X-N5	-2.19	1.31	1.35
2	A	500	FAD	C5A-N7A	-2.15	1.31	1.39
2	A	500	FAD	C2'-C3'	-2.14	1.49	1.53
3	B	501	NAD	C2D-C1D	-2.02	1.50	1.53
3	A	501	NAD	C4N-C3N	2.02	1.42	1.39
2	A	500	FAD	C8-C7	2.04	1.46	1.41
4	A	502	ASC	C5-C4	2.09	1.56	1.53
2	A	500	FAD	C4-C4X	2.12	1.45	1.41
2	B	500	FAD	C4X-C10	2.23	1.45	1.40
2	B	500	FAD	C9A-C5X	2.27	1.47	1.42
2	B	500	FAD	C4-C4X	2.27	1.45	1.41
2	A	500	FAD	C4X-C10	2.50	1.45	1.40
3	B	501	NAD	O4D-C1D	2.56	1.44	1.41
2	A	500	FAD	C9A-C5X	2.70	1.48	1.42
3	A	501	NAD	C5A-C4A	3.25	1.47	1.40
3	A	501	NAD	O4B-C1B	3.49	1.46	1.41
3	B	501	NAD	C5A-C4A	3.61	1.48	1.40
4	A	502	ASC	O4-C1	8.21	1.48	1.35
4	B	502	ASC	O4-C1	9.33	1.49	1.35

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	NAD	N3A-C2A-N1A	-10.34	120.75	128.87
3	B	501	NAD	N3A-C2A-N1A	-8.84	121.93	128.87
4	A	502	ASC	C4-O4-C1	-8.27	101.22	109.29
4	B	502	ASC	C4-O4-C1	-7.67	101.81	109.29
3	B	501	NAD	C1B-N9A-C4A	-6.07	120.03	126.81
2	B	500	FAD	N3A-C2A-N1A	-6.04	124.13	128.87
2	A	500	FAD	N3A-C2A-N1A	-5.60	124.47	128.87
4	A	502	ASC	O1-C1-C2	-5.60	122.47	129.30
4	A	502	ASC	O4-C4-C3	-5.50	99.77	104.13
2	B	500	FAD	C4-C4X-C10	-4.85	116.83	119.94
2	A	500	FAD	C4-C4X-C10	-4.58	117.01	119.94

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	502	ASC	O1-C1-C2	-4.48	123.83	129.30
4	B	502	ASC	O4-C4-C3	-4.46	100.60	104.13
3	B	501	NAD	O4B-C1B-N9A	-4.28	100.01	108.11
2	A	500	FAD	O4'-C4'-C5'	-3.97	101.43	110.09
3	A	501	NAD	C4B-O4B-C1B	-3.92	105.49	109.64
2	A	500	FAD	C4X-C4-N3	-3.65	118.75	123.52
2	B	500	FAD	C4X-C4-N3	-3.23	119.29	123.52
3	A	501	NAD	O7N-C7N-N7N	-3.10	118.16	122.58
4	A	502	ASC	O3-C3-C2	-3.06	123.38	132.56
2	B	500	FAD	O3'-C3'-C4'	-2.92	101.15	108.73
2	B	500	FAD	O4'-C4'-C3'	-2.72	101.97	108.96
2	A	500	FAD	N3-C2-N1	-2.65	123.23	127.69
4	A	502	ASC	C5-C4-C3	-2.56	110.48	114.17
2	B	500	FAD	N3-C2-N1	-2.50	123.48	127.69
2	A	500	FAD	O2'-C2'-C3'	-2.47	102.59	108.96
3	B	501	NAD	C4B-O4B-C1B	-2.34	107.17	109.64
2	A	500	FAD	O3'-C3'-C2'	-2.03	103.46	108.73
4	A	502	ASC	O2-C2-C1	2.10	127.66	121.84
3	A	501	NAD	O4B-C4B-C3B	2.13	109.48	105.16
3	B	501	NAD	C3N-C7N-N7N	2.14	120.24	117.82
2	B	500	FAD	C6-C5X-C9A	2.24	121.58	119.11
2	A	500	FAD	C6-C5X-C9A	2.38	121.73	119.11
4	A	502	ASC	O3-C3-C4	2.39	124.32	117.91
2	B	500	FAD	C1'-N10-C9A	2.60	121.84	118.83
4	B	502	ASC	O5-C5-C4	2.63	114.83	110.33
3	A	501	NAD	N6A-C6A-N1A	2.67	123.00	118.52
3	B	501	NAD	C2D-C3D-C4D	2.89	108.55	102.64
2	A	500	FAD	C1'-C2'-C3'	2.93	118.19	109.82
2	B	500	FAD	C4X-N5-C5X	3.00	120.26	116.72
2	A	500	FAD	C5X-C9A-N10	3.14	119.93	117.58
2	B	500	FAD	O3'-C3'-C2'	3.36	117.43	108.73
3	A	501	NAD	C2A-N1A-C6A	3.49	125.00	118.77
3	A	501	NAD	C3N-C7N-N7N	3.67	121.97	117.82
3	B	501	NAD	C2A-N1A-C6A	3.72	125.41	118.77
4	A	502	ASC	O5-C5-C4	4.74	118.46	110.33
2	B	500	FAD	C5X-C9A-N10	4.86	121.22	117.58
3	B	501	NAD	O4D-C1D-N1N	5.27	113.80	108.10
3	A	501	NAD	O4D-C1D-N1N	5.53	114.08	108.10
4	B	502	ASC	O4-C1-O1	5.66	127.70	121.27
4	A	502	ASC	O4-C1-O1	6.46	128.61	121.27
2	B	500	FAD	C1'-C2'-C3'	6.71	129.02	109.82
2	B	500	FAD	C4-N3-C2	6.93	120.94	115.16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	FAD	C1'-N10-C9A	7.54	127.57	118.83
2	A	500	FAD	C4-N3-C2	7.68	121.57	115.16
4	B	502	ASC	O4-C4-C5	7.74	122.90	109.69
4	A	502	ASC	O4-C4-C5	8.70	124.56	109.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	FAD	1	0
3	A	501	NAD	1	0
4	A	502	ASC	1	0
2	B	500	FAD	3	0
3	B	501	NAD	3	0
4	B	502	ASC	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	429/451 (95%)	0.66	41 (9%) 10 15	51, 72, 113, 135	0
1	B	425/451 (94%)	0.55	34 (8%) 15 21	49, 70, 100, 138	0
All	All	854/902 (94%)	0.60	75 (8%) 12 18	49, 71, 107, 138	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	308	ASN	6.8
1	B	6	HIS	5.9
1	A	340	VAL	5.2
1	A	96	ILE	5.1
1	A	114	PHE	5.0
1	A	424	LEU	5.0
1	B	425	LYS	4.9
1	B	306	MET	4.9
1	A	283	ASP	4.9
1	B	392	VAL	4.8
1	A	430	GLN	4.6
1	B	434	LYS	4.6
1	A	97	VAL	4.4
1	A	413	LYS	4.1
1	A	423	GLU	4.1
1	B	338	SER	4.0
1	A	427	GLU	4.0
1	B	417	PRO	3.9
1	A	433	SER	3.4
1	A	273	VAL	3.4
1	A	422	GLU	3.3
1	A	432	ALA	3.2
1	B	432	ALA	3.2
1	A	285	PHE	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	428	GLY	3.1
1	A	160	GLN	3.0
1	B	115	THR	3.0
1	A	315	HIS	3.0
1	B	416	PRO	2.9
1	A	297	GLY	2.9
1	A	310	LEU	2.8
1	B	77	ARG	2.7
1	B	307	TYR	2.7
1	B	229	ALA	2.6
1	A	271	GLY	2.6
1	B	424	LEU	2.6
1	A	223	ILE	2.6
1	B	387	ILE	2.6
1	B	388	LYS	2.5
1	B	345	LEU	2.5
1	B	110	VAL	2.5
1	A	282	THR	2.5
1	B	257	VAL	2.5
1	A	146	LEU	2.5
1	A	341	GLU	2.4
1	B	97	VAL	2.4
1	A	378	ALA	2.4
1	A	6	HIS	2.4
1	A	113	THR	2.3
1	B	433	SER	2.3
1	B	430	GLN	2.3
1	A	238	GLY	2.3
1	B	39	SER	2.3
1	B	224	VAL	2.3
1	B	105	THR	2.2
1	B	409	ALA	2.2
1	A	390	GLY	2.2
1	B	260	VAL	2.2
1	A	333	LYS	2.2
1	B	364	VAL	2.2
1	A	286	PHE	2.2
1	A	426	LYS	2.1
1	A	329	ALA	2.1
1	B	102	ALA	2.1
1	A	107	THR	2.1
1	A	115	THR	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	304	MET	2.1
1	A	105	THR	2.1
1	A	202	ARG	2.1
1	A	236	ALA	2.1
1	A	379	LYS	2.1
1	B	188	PHE	2.0
1	A	280	ILE	2.0
1	A	435	ILE	2.0
1	B	413	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	ASC	B	502	12/12	0.66	0.35	5.82	73,86,95,97	0
4	ASC	A	502	12/12	0.84	0.19	0.54	71,78,90,90	0
3	NAD	B	501	44/44	0.88	0.19	-0.08	50,71,102,109	0
3	NAD	A	501	44/44	0.91	0.18	-0.19	61,71,99,127	0
2	FAD	A	500	53/53	0.94	0.13	-0.82	17,22,27,30	0
2	FAD	B	500	53/53	0.94	0.13	-0.99	17,22,27,30	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.