



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2016 – 08:09 AM EDT

PDB ID : 5JDO  
Title : T. congolense haptoglobin-haemoglobin receptor in complex with haemoglobin  
Authors : Lane-Serff, H.; Higgins, M.K.  
Deposited on : 2016-04-17  
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027257  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027257


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1124 (3.24-3.16)
Clashscore	102246	1024 (3.22-3.18)
Ramachandran outliers	100387	1004 (3.22-3.18)
Sidechain outliers	100360	1003 (3.22-3.18)
RSRZ outliers	91569	1129 (3.24-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>.</div> </div> </div>
2	B	245	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>7%</div> <div>.</div> </div> </div>
3	C	141	<div> <div></div> <div> <div></div> <div>85%</div> <div>14%</div> <div>.</div> </div> </div>
4	D	145	<div> <div></div> <div> <div></div> <div>87%</div> <div>13%</div> <div></div> </div> </div>
5	E	140	<div> <div></div> <div> <div></div> <div>88%</div> <div>11%</div> <div>.</div> </div> </div>
6	F	144	<div> <div></div> <div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	OXY	C	202	-	-	-	X
8	OXY	E	202	-	-	-	X
8	OXY	F	202	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 8226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Haptoglobin-haemoglobin receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	1	0
			1866	1136	328	400	2			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	GLU	LYS	conflict	UNP G0UVW6
A	12	SER	ALA	conflict	UNP G0UVW6
A	14	ASP	GLU	conflict	UNP G0UVW6
A	55	ASP	GLU	conflict	UNP G0UVW6
A	64	VAL	ALA	conflict	UNP G0UVW6
A	72	ASP	ASN	conflict	UNP G0UVW6
A	80	THR	ALA	conflict	UNP G0UVW6
A	112	VAL	ALA	conflict	UNP G0UVW6
A	116	ILE	THR	conflict	UNP G0UVW6
A	137	ALA	VAL	conflict	UNP G0UVW6
A	152	ASN	ASP	conflict	UNP G0UVW6
A	157	GLU	LYS	conflict	UNP G0UVW6
A	164	GLU	ASN	conflict	UNP G0UVW6

- Molecule 2 is a protein called Haptoglobin-haemoglobin receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	245	Total	C	N	O	S	0	1	0
			1834	1119	324	389	2			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	GLU	LYS	conflict	UNP G0UVW6
B	12	SER	ALA	conflict	UNP G0UVW6
B	14	ASP	GLU	conflict	UNP G0UVW6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	55	ASP	GLU	conflict	UNP G0UVW6
B	64	VAL	ALA	conflict	UNP G0UVW6
B	72	ASP	ASN	conflict	UNP G0UVW6
B	80	THR	ALA	conflict	UNP G0UVW6
B	112	VAL	ALA	conflict	UNP G0UVW6
B	116	ILE	THR	conflict	UNP G0UVW6
B	137	ALA	VAL	conflict	UNP G0UVW6
B	152	ASN	ASP	conflict	UNP G0UVW6
B	157	GLU	LYS	conflict	UNP G0UVW6
B	164	GLU	ASN	conflict	UNP G0UVW6

- Molecule 3 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	141	Total	C	N	O	S	0	0	0
			1068	685	187	193	3			

- Molecule 4 is a protein called Hemoglobin subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	145	Total	C	N	O	S	0	0	0
			1116	719	194	200	3			


- Molecule 5 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	140	Total	C	N	O	S	0	0	0
			1057	679	183	192	3			

- Molecule 6 is a protein called Hemoglobin subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	144	Total	C	N	O	S	0	0	0
			1105	713	191	198	3			

- Molecule 7 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
7	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
7	E	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
7	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 8 is OXYGEN MOLECULE (three-letter code: OXY) (formula: O<sub>2</sub>).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	1	Total 2	O 2	0	0
8	D	1	Total 2	O 2	0	0
8	E	1	Total 2	O 2	0	0
8	F	1	Total 2	O 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Haptoglobin-haemoglobin receptor




- Molecule 2: Haptoglobin-haemoglobin receptor




- Molecule 3: Hemoglobin subunit alpha




- Molecule 4: Hemoglobin subunit beta



- Molecule 5: Hemoglobin subunit alpha



- Molecule 6: Hemoglobin subunit beta





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.75Å 127.30Å 172.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	102.41 – 3.20 67.03 – 3.20	Depositor EDS
% Data completeness (in resolution range)	97.7 (102.41-3.20) 97.7 (67.03-3.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.35 (at 3.19Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.204 , 0.235 0.217 , 0.241	Depositor DCC
$R_{free}$ test set	1346 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	82.3	Xtriage
Anisotropy	0.663	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8226	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, OXY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/1875	0.70	0/2523
2	B	0.49	0/1843	0.67	0/2481
3	C	0.50	0/1096	0.65	0/1491
4	D	0.50	0/1146	0.67	1/1556 (0.1%)
5	E	0.50	0/1085	0.66	0/1477
6	F	0.48	0/1134	0.65	0/1541
All	All	0.50	0/8179	0.67	1/11069 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	D	146	TYR	C-N-CA	5.38	135.14	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1866	0	1874	12	0
2	B	1834	0	1850	5	0
3	C	1068	0	1070	13	0
4	D	1116	0	1106	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	E	1057	0	1057	8	0
6	F	1105	0	1099	4	0
7	C	43	0	30	4	0
7	D	43	0	30	1	0
7	E	43	0	30	2	0
7	F	43	0	30	1	0
8	C	2	0	0	0	0
8	D	2	0	0	0	0
8	E	2	0	0	0	0
8	F	2	0	0	0	0
All	All	8226	0	8176	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:F:201:HEM:HHC	7:F:201:HEM:HBB2	1.60	0.84
1:A:90:ALA:HB3	1:A:240:ALA:HB3	1.64	0.80
3:C:94:VAL:HG13	7:C:201:HEM:HBC2	1.68	0.74
1:A:235:LYS:O	1:A:239:GLU:HB2	1.87	0.74
7:C:201:HEM:HBB2	7:C:201:HEM:HHC	1.77	0.66
3:C:94:VAL:CG1	7:C:201:HEM:HBC2	2.28	0.64
1:A:45:THR:HG22	4:D:92:LEU:HD11	1.82	0.61
6:F:64:HIS:O	6:F:68:VAL:HG23	2.05	0.56
5:E:59:HIS:HA	5:E:62:LYS:HE3	1.87	0.56
2:B:87:VAL:HG22	2:B:243:GLU:HB2	1.89	0.55
3:C:59:HIS:HA	3:C:62:LYS:HE3	1.88	0.54
1:A:90:ALA:HB3	1:A:240:ALA:CB	2.36	0.54
1:A:237:ALA:O	1:A:241:GLU:HB3	2.08	0.53
5:E:48:ASP:HB3	5:E:55:GLN:HG3	1.91	0.53
3:C:114:LEU:HB3	3:C:117:GLU:HB2	1.89	0.53
3:C:142:ARG:HD3	5:E:128:LYS:HD3	1.91	0.53
3:C:48:ASP:HB3	3:C:55:GLN:HG3	1.91	0.52
1:A:46:THR:HB	1:A:112:VAL:HG22	1.92	0.51
5:E:67:LEU:O	5:E:71:VAL:HG23	2.10	0.51
1:A:243:GLU:HA	1:A:246:LYS:HG2	1.93	0.51
3:C:67:LEU:O	3:C:71:VAL:HG23	2.12	0.50
1:A:22:ARG:HD3	1:A:141:SER:OG	2.13	0.49
6:F:52:PRO:O	6:F:56:MET:HG2	2.14	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:52:PRO:O	4:D:56:MET:HG2	2.14	0.47
3:C:77:MET:N	3:C:78:PRO:HD2	2.30	0.47
7:E:201:HEM:HMA2	7:E:201:HEM:HBA1	1.97	0.46
7:E:201:HEM:CMA	7:E:201:HEM:HBA1	2.46	0.45
2:B:77:LEU:HD13	2:B:78:ASP:H	1.82	0.45
4:D:42:PHE:CZ	4:D:99:VAL:HG22	2.52	0.44
5:E:95:ASP:OD1	5:E:97:VAL:HG22	2.18	0.44
5:E:77:MET:N	5:E:78:PRO:HD2	2.32	0.44
3:C:7:ASP:O	3:C:11:VAL:HG23	2.18	0.44
3:C:94:VAL:HG13	7:C:201:HEM:CBC	2.45	0.43
6:F:12:VAL:HG12	6:F:76:LEU:HD21	1.99	0.43
3:C:93:ARG:CG	3:C:93:ARG:HH11	2.31	0.43
2:B:84:ALA:HB1	2:B:88:ARG:HH12	1.84	0.43
3:C:95:ASP:OD1	3:C:97:VAL:HG22	2.18	0.43
1:A:49:SER:HA	1:A:52:LYS:HD3	2.00	0.43
6:F:22:ASP:HA	6:F:66:LYS:HD3	2.01	0.42
2:B:129:ALA:HB2	2:B:201:SER:HB3	2.02	0.42
4:D:42:PHE:CD2	7:D:201:HEM:HBC1	2.55	0.42
2:B:46:THR:HB	2:B:112:VAL:HG22	2.02	0.41
5:E:43:TYR:CE1	5:E:94:VAL:HA	2.55	0.41
1:A:129:ALA:HB2	1:A:201:SER:HB3	2.03	0.41
5:E:7:ASP:O	5:E:11:VAL:HG23	2.20	0.41
1:A:90:ALA:CB	1:A:240:ALA:CB	2.99	0.41
1:A:134:ASP:O	1:A:138:VAL:HG23	2.21	0.41
4:D:29:LEU:HD23	4:D:61:VAL:HG13	2.02	0.41
4:D:93:HIS:HA	4:D:97:LEU:HD12	2.02	0.41
3:C:43:TYR:CE1	3:C:94:VAL:HA	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/249 (100%)	236 (95%)	12 (5%)	0	100	100
2	B	244/245 (100%)	238 (98%)	6 (2%)	0	100	100
3	C	139/141 (99%)	135 (97%)	4 (3%)	0	100	100
4	D	143/145 (99%)	134 (94%)	9 (6%)	0	100	100
5	E	138/140 (99%)	133 (96%)	5 (4%)	0	100	100
6	F	142/144 (99%)	137 (96%)	5 (4%)	0	100	100
All	All	1054/1064 (99%)	1013 (96%)	41 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/195 (100%)	179 (91%)	17 (9%)	13	45
2	B	192/191 (100%)	179 (93%)	13 (7%)	20	59
3	C	113/113 (100%)	109 (96%)	4 (4%)	43	80
4	D	117/117 (100%)	108 (92%)	9 (8%)	16	54
5	E	112/112 (100%)	109 (97%)	3 (3%)	52	85
6	F	116/116 (100%)	107 (92%)	9 (8%)	16	53
All	All	846/844 (100%)	791 (94%)	55 (6%)	21	61

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	VAL
1	A	22	ARG
1	A	32	SER
1	A	35	LEU
1	A	47	GLU
1	A	49	SER
1	A	77	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	79	LYS
1	A	85	GLN
1	A	160	GLU
1	A	193	THR
1	A	201	SER
1	A	239	GLU
1	A	241	GLU
1	A	242	THR
1	A	246	LYS
1	A	249	THR
2	B	7	VAL
2	B	22	ARG
2	B	32	SER
2	B	35	LEU
2	B	50	ARG
2	B	77	LEU
2	B	150	SER
2	B	160	GLU
2	B	193	THR
2	B	201	SER
2	B	219	LEU
2	B	227	GLU
2	B	243	GLU
3	C	17	LYS
3	C	39	THR
3	C	93	ARG
3	C	140	LYS
4	D	10	SER
4	D	15	LEU
4	D	22	ASP
4	D	27	GLU
4	D	76	LEU
4	D	81	ASN
4	D	83	LYS
4	D	142	LEU
4	D	147	HIS
5	E	107	LEU
5	E	128	LYS
5	E	140	LYS
6	F	10	SER
6	F	15	LEU
6	F	22	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
6	F	27	GLU
6	F	29	LEU
6	F	76	LEU
6	F	80	ASP
6	F	83	LYS
6	F	142	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	HIS
1	A	132	ASN
2	B	85	GLN
2	B	132	ASN
2	B	153	GLN
3	C	113	HIS
5	E	73	HIS
6	F	118	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	HEM	C	201	3	24,50,50	0.80	1 (4%)	16,82,82	1.36	3 (18%)
8	OXY	C	202	-	1,1,1	0.91	0	0,0,0	0.00	-
7	HEM	D	201	4	24,50,50	0.83	0	16,82,82	1.65	4 (25%)
8	OXY	D	202	-	1,1,1	0.97	0	0,0,0	0.00	-
7	HEM	E	201	8,5	24,50,50	0.99	2 (8%)	16,82,82	1.51	4 (25%)
8	OXY	E	202	7	1,1,1	0.95	0	0,0,0	0.00	-
7	HEM	F	201	8,6	24,50,50	1.01	3 (12%)	16,82,82	1.83	2 (12%)
8	OXY	F	202	7	1,1,1	0.87	0	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	HEM	C	201	3	-	0/6/54/54	0/0/8/8
8	OXY	C	202	-	-	0/0/0/0	0/0/0/0
7	HEM	D	201	4	-	0/6/54/54	0/0/8/8
8	OXY	D	202	-	-	0/0/0/0	0/0/0/0
7	HEM	E	201	8,5	-	2/6/54/54	0/0/8/8
8	OXY	E	202	7	-	0/0/0/0	0/0/0/0
7	HEM	F	201	8,6	-	0/6/54/54	0/0/8/8
8	OXY	F	202	7	-	0/0/0/0	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	E	201	HEM	C3B-C2B	-2.87	1.36	1.40
7	F	201	HEM	C1B-NB	-2.78	1.33	1.36
7	C	201	HEM	C3B-C2B	-2.57	1.37	1.40
7	F	201	HEM	C3B-C2B	-2.40	1.37	1.40
7	E	201	HEM	C1B-NB	-2.39	1.33	1.36
7	F	201	HEM	C4D-ND	-2.02	1.34	1.36

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	F	201	HEM	CBD-CAD-C3D	-4.45	104.67	112.47
7	F	201	HEM	C3C-C4C-NC	-3.75	103.86	110.94
7	E	201	HEM	CMA-C3A-C4A	-3.44	122.46	128.31
7	D	201	HEM	C3C-CAC-CBC	-3.08	120.20	126.40
7	D	201	HEM	C3C-C4C-NC	-2.94	105.39	110.94
7	C	201	HEM	C3C-C4C-NC	-2.89	105.49	110.94
7	E	201	HEM	C3C-CAC-CBC	-2.79	120.79	126.40
7	C	201	HEM	CAA-CBA-CGA	-2.44	108.03	112.78
7	D	201	HEM	C3B-CAB-CBB	-2.30	121.76	126.40
7	E	201	HEM	C3C-C4C-NC	-2.27	106.65	110.94
7	D	201	HEM	CMA-C3A-C4A	-2.16	124.63	128.31
7	C	201	HEM	CBD-CAD-C3D	-2.08	108.81	112.47
7	E	201	HEM	CMA-C3A-C2A	2.28	130.00	125.24

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	E	201	HEM	C3A-C2A-CAA-CBA
7	E	201	HEM	C1A-C2A-CAA-CBA

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	C	201	HEM	4	0
7	D	201	HEM	1	0
7	E	201	HEM	2	0
7	F	201	HEM	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	249/249 (100%)	0.20	5 (2%) 68 54	58, 92, 156, 190	0
2	B	245/245 (100%)	0.05	2 (0%) 87 80	59, 92, 164, 185	0
3	C	141/141 (100%)	-0.06	0 100 100	49, 67, 89, 99	0
4	D	145/145 (100%)	0.02	0 100 100	54, 77, 103, 116	0
5	E	140/140 (100%)	-0.13	0 100 100	53, 72, 96, 102	0
6	F	144/144 (100%)	0.03	0 100 100	56, 78, 99, 112	0
All	All	1064/1064 (100%)	0.04	7 (0%) 89 83	49, 81, 147, 190	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	251	SER	6.1
1	A	78	ASP	3.4
2	B	83	THR	3.0
1	A	249	THR	2.5
1	A	77	LEU	2.3
1	A	84	ALA	2.1
2	B	245	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
8	OXY	C	202	2/2	0.80	1.43	41.54	124,124,124,124	0
8	OXY	F	202	2/2	0.98	0.81	6.85	96,96,96,96	0
8	OXY	E	202	2/2	0.99	0.49	6.04	164,164,164,164	0
7	HEM	F	201	43/43	0.97	0.30	0.28	54,63,77,85	0
7	HEM	E	201	43/43	0.98	0.25	0.03	39,53,69,76	0
7	HEM	D	201	43/43	0.97	0.26	0.03	61,66,80,83	0
7	HEM	C	201	43/43	0.98	0.24	-0.40	47,54,57,61	0
8	OXY	D	202	2/2	0.97	1.27	-	154,154,154,154	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.