



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 18, 2016 – 03:53 PM EDT

PDB ID : 5JH7  
Title : Tubulin-Eribulin complex  
Authors : Doodhi, H.; Prota, A.E.; Rodriguez-Garcia, R.; Xiao, H.; Custar, D.W.; Bargsten, K.; Katrukha, E.A.; Hilbert, M.; Hua, S.; Jiang, K.; Grigoriev, I.; Yang, C.-P.H.; Cox, D.; Band Horwitz, S.; Kapitein, L.C.; Akhmanova, A.; Steinmetz, M.O.  
Deposited on : 2016-04-20  
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	<div> <div></div> <div>83% 14% ..</div> </div>
1	C	450	<div> <div></div> <div>83% 14% ..</div> </div>
2	B	445	<div> <div>2%</div> <div>77% 19% .</div> </div>
2	D	445	<div> <div>3%</div> <div>74% 22% 5%</div> </div>
3	E	143	<div> <div>2%</div> <div>69% 16% 15%</div> </div>
4	F	384	<div> <div>10%</div> <div>70% 18% . 11%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	CA	A	503	-	-	-	X
7	CA	C	503	-	-	-	X
8	IMD	B	507	-	-	X	-
9	GOL	A	505	-	-	-	X

## 2 Entry composition

There are 15 unique types of molecules in this entry. The entry contains 18483 atoms, of which 118 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	439	Total	C	N	O	S	0	2	0
			3436	2174	583	655	24			
1	C	440	Total	C	N	O	S	0	0	0
			3437	2175	584	656	22			

- Molecule 2 is a protein called Tubulin beta-2B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	428	Total	C	N	O	S	0	1	0
			3360	2112	573	648	27			
2	D	424	Total	C	N	O	S	0	1	0
			3335	2095	568	644	28			

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	121	Total	C	N	O	S	0	0	0
			1000	617	181	197	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MET	ILE	conflict	UNP P63043
E	4	ALA	SER	conflict	UNP P63043


- Molecule 4 is a protein called Tubulin Tyrosine Ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	F	342	Total	C	N	O	S	0	0	0
			2800	1797	477	512	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	379	HIS	-	expression tag	UNP E1BQ43
F	380	HIS	-	expression tag	UNP E1BQ43
F	381	HIS	-	expression tag	UNP E1BQ43
F	382	HIS	-	expression tag	UNP E1BQ43
F	383	HIS	-	expression tag	UNP E1BQ43
F	384	HIS	-	expression tag	UNP E1BQ43

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula:  $C_{10}H_{16}N_5O_{14}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
5	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		
6	F	1	Total	Mg	0	0
			1	1		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Ca 1 1	0	0
7	A	1	Total Ca 1 1	0	0
7	C	2	Total Ca 2 2	0	0
7	E	1	Total Ca 1 1	0	0

- Molecule 8 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C N 5 3 2	0	0
8	B	1	Total C N 5 3 2	0	0
8	B	1	Total C N 5 3 2	0	0

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			6	3	3		
9	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 10 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>11</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	B	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
10	D	1	28	10	5	11	2	0	0


- Molecule 11 is (1S,3S,6S,9S,12S,14R,16R,18S,20R,21R,22S,26R,29S,31R,32S,33R,35R,36S)-20-[(2S)-3-amino-2-hydroxypropyl]-21-methoxy-14-methyl-8,15-dimethylidene-2,19,30,34,37,39,40,41-octaoxanonacyclo[24.9.2.1 3,32 .1 3,33 .1 6,9 .1 12,16 .0 18,22 .0 29,36 .0 31,35 ]hen tetracontan-24-one (non-preferred name) (three-letter code: 6K9) (formula: C<sub>40</sub>H<sub>59</sub>NO<sub>11</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
11	B	1	111	40	59	1	11	0	0
11	D	1	111	40	59	1	11	0	0


- Molecule 12 is methanesulfonic acid (three-letter code: 03S) (formula: CH<sub>4</sub>O<sub>3</sub>S).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
12	B	1	Total	C	O	S	0	0
			5	1	3	1		

- Molecule 13 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
13	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 14 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula:  $C_{11}H_{18}N_5O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
14	F	1	Total	C	N	O	P	0	0
			31	11	5	12	3		


- Molecule 15 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	150	Total	O	0	0
			150	150		
15	B	112	Total	O	0	0
			112	112		
15	C	231	Total	O	0	0
			231	231		
15	D	99	Total	O	0	0
			99	99		
15	E	34	Total	O	0	0
			34	34		
15	F	64	Total	O	0	0
			64	64		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain



- Molecule 1: Tubulin alpha-1B chain




- Molecule 2: Tubulin beta-2B chain







• Molecule 2: Tubulin beta-2B chain



• Molecule 3: Stathmin-4



• Molecule 4: Tubulin Tyrosine Ligase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.22Å 157.13Å 182.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.18 – 2.25 78.87 – 2.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (72.18-2.25) 100.0 (78.87-2.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 2.25Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.181 , 0.220 0.182 , 0.220	Depositor DCC
$R_{free}$ test set	7162 reflections (4.98%)	DCC
Wilson B-factor (Å <sup>2</sup> )	54.5	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 52.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	18483	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, 6K9, MG, IMD, GOL, CA, GTP, ACP, MES, 03S

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.21	0/3520	0.38	0/4778
1	C	0.22	0/3515	0.39	0/4772
2	B	0.22	0/3438	0.37	0/4658
2	D	0.22	0/3411	0.37	0/4620
3	E	0.20	0/1008	0.30	0/1337
4	F	0.21	0/2863	0.36	0/3866
All	All	0.22	0/17755	0.37	0/24031

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3436	0	3349	51	0
1	C	3437	0	3348	45	0
2	B	3360	0	3232	72	0
2	D	3335	0	3219	66	0
3	E	1000	0	1018	17	0
4	F	2800	0	2764	53	0
5	A	32	0	12	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	32	0	12	0	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
6	F	1	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	2	0	0	0	0
7	E	1	0	0	0	0
8	A	5	0	5	0	0
8	B	10	0	10	4	0
9	A	6	0	8	0	0
9	C	6	0	8	1	0
10	B	28	0	12	1	0
10	D	28	0	12	3	0
11	B	52	59	0	0	0
11	D	52	59	0	0	0
12	B	5	0	4	1	0
13	B	12	0	12	2	0
14	F	31	0	14	3	0
15	A	150	0	0	3	0
15	B	112	0	0	3	0
15	C	231	0	0	2	0
15	D	99	0	0	5	0
15	E	34	0	0	4	0
15	F	64	0	0	0	0
All	All	18365	118	17039	301	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 301 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:25:LYS:HD2	3:E:26:PRO:HD2	1.49	0.94
2:D:47:GLU:HG2	2:D:245:PRO:HG3	1.53	0.89
1:C:93:ILE:HD11	1:C:121:ARG:HG3	1.55	0.88
4:F:7:ARG:HD3	4:F:40:MET:HE3	1.58	0.86
2:D:213:CYS:HB3	2:D:219:LEU:HD11	1.59	0.83

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/450 (98%)	422 (96%)	17 (4%)	0	100	100
1	C	438/450 (97%)	427 (98%)	11 (2%)	0	100	100
2	B	427/445 (96%)	422 (99%)	5 (1%)	0	100	100
2	D	421/445 (95%)	414 (98%)	7 (2%)	0	100	100
3	E	117/143 (82%)	116 (99%)	1 (1%)	0	100	100
4	F	332/384 (86%)	318 (96%)	14 (4%)	0	100	100
All	All	2174/2317 (94%)	2119 (98%)	55 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/378 (98%)	368 (99%)	4 (1%)	80	88
1	C	371/378 (98%)	361 (97%)	10 (3%)	52	63
2	B	367/383 (96%)	365 (100%)	2 (0%)	92	95
2	D	367/383 (96%)	366 (100%)	1 (0%)	94	97
3	E	109/127 (86%)	108 (99%)	1 (1%)	84	91
4	F	307/342 (90%)	304 (99%)	3 (1%)	82	89
All	All	1893/1991 (95%)	1872 (99%)	21 (1%)	80	88

5 of 21 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	C	245	ASP
1	C	293	ASN
3	E	53	LYS
1	C	221	ARG
4	F	138	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	342	GLN
2	D	167	ASN
4	F	145	ASN
4	F	242	ASN
4	F	269	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 8 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GTP	A	501	6	26,34,34	0.95	1 (3%)	29,54,54	1.58	3 (10%)
8	IMD	A	504	-	3,5,5	0.53	0	4,5,5	0.56	0
9	GOL	A	505	-	5,5,5	0.37	0	5,5,5	0.19	0
10	GDP	B	501	-	24,30,30	1.15	2 (8%)	26,47,47	1.76	4 (15%)
11	6K9	B	503	-	57,60,60	1.99	13 (22%)	59,92,92	1.61	13 (22%)
12	03S	B	504	-	4,4,4	4.40	2 (50%)	1,6,6	3.11	1 (100%)
13	MES	B	505	-	12,12,12	2.15	1 (8%)	15,16,16	2.22	7 (46%)
8	IMD	B	506	-	3,5,5	0.51	0	4,5,5	0.56	0
8	IMD	B	507	-	3,5,5	0.51	0	4,5,5	0.55	0
5	GTP	C	501	6	26,34,34	0.94	1 (3%)	29,54,54	1.64	3 (10%)
9	GOL	C	505	-	5,5,5	0.34	0	5,5,5	0.27	0
10	GDP	D	501	-	24,30,30	1.15	2 (8%)	26,47,47	1.83	4 (15%)
11	6K9	D	502	-	57,60,60	1.98	13 (22%)	59,92,92	1.58	14 (23%)
14	ACP	F	400	6	29,33,33	2.39	9 (31%)	29,52,52	2.57	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GTP	A	501	6	-	0/18/38/38	0/3/3/3
8	IMD	A	504	-	-	0/0/0/0	0/1/1/1
9	GOL	A	505	-	-	0/4/4/4	0/0/0/0
10	GDP	B	501	-	-	0/12/32/32	0/3/3/3
11	6K9	B	503	-	-	0/31/131/131	0/0/9/9
12	03S	B	504	-	-	0/0/0/0	0/0/0/0
13	MES	B	505	-	-	0/6/14/14	0/1/1/1
8	IMD	B	506	-	-	0/0/0/0	0/1/1/1
8	IMD	B	507	-	-	0/0/0/0	0/1/1/1
5	GTP	C	501	6	-	0/18/38/38	0/3/3/3
9	GOL	C	505	-	-	0/4/4/4	0/0/0/0
10	GDP	D	501	-	-	0/12/32/32	0/3/3/3
11	6K9	D	502	-	-	0/31/131/131	0/0/9/9
14	ACP	F	400	6	-	0/15/38/38	0/3/3/3

The worst 5 of 44 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
13	B	505	MES	C8-S	-7.17	1.66	1.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	D	502	6K9	C02-C01	-6.82	1.41	1.51
11	B	503	6K9	C02-C01	-6.59	1.41	1.51
11	D	502	6K9	C39-C01	-5.29	1.43	1.51
11	B	503	6K9	C39-C01	-5.24	1.43	1.51

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	F	400	ACP	N3-C2-N1	-8.89	121.89	128.87
14	F	400	ACP	O4'-C1'-N9	-7.66	93.63	108.11
5	C	501	GTP	N3-C2-N1	-5.54	120.02	127.56
5	A	501	GTP	N3-C2-N1	-5.40	120.21	127.56
10	D	501	GDP	C5-C6-N1	-4.40	117.78	123.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	501	GTP	1	0
10	B	501	GDP	1	0
12	B	504	03S	1	0
13	B	505	MES	2	0
8	B	506	IMD	2	0
8	B	507	IMD	4	0
9	C	505	GOL	1	0
10	D	501	GDP	3	0
14	F	400	ACP	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	439/450 (97%)	-0.05	5 (1%) 82 84	41, 66, 99, 145	0
1	C	440/450 (97%)	0.18	2 (0%) 91 92	37, 53, 79, 100	0
2	B	428/445 (96%)	0.25	7 (1%) 74 77	40, 65, 113, 155	2 (0%)
2	D	424/445 (95%)	0.07	12 (2%) 56 61	43, 71, 105, 136	6 (1%)
3	E	121/143 (84%)	0.37	3 (2%) 61 65	49, 83, 106, 119	0
4	F	342/384 (89%)	0.68	40 (11%) 6 6	48, 86, 145, 172	0
All	All	2194/2317 (94%)	0.22	69 (3%) 52 57	37, 68, 115, 172	8 (0%)

The worst 5 of 69 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	282	TYR	6.3
4	F	173	ILE	6.1
1	A	439	SER	5.8
4	F	132	LEU	5.4
2	D	285	ALA	5.3

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	CA	A	503	1/1	0.67	0.40	14.29	219,219,219,219	0
9	GOL	A	505	6/6	0.85	0.18	4.44	69,85,88,89	0
7	CA	C	503	1/1	0.96	0.19	2.06	69,69,69,69	0
10	GDP	B	501	28/28	0.97	0.19	0.75	40,51,57,60	0
11	6K9	D	502	52/52	0.94	0.17	0.70	63,92,111,115	0
5	GTP	C	501	32/32	0.98	0.16	0.62	35,40,51,53	0
5	GTP	A	501	32/32	0.98	0.13	0.41	38,49,57,84	0
10	GDP	D	501	28/28	0.96	0.12	-0.22	45,61,81,86	0
8	IMD	B	507	5/5	0.94	0.12	-0.52	107,109,111,113	0
13	MES	B	505	12/12	0.94	0.13	-0.79	57,78,100,112	0
8	IMD	A	504	5/5	0.92	0.10	-0.98	85,87,90,90	0
8	IMD	B	506	5/5	0.89	0.13	-1.03	94,96,102,103	0
9	GOL	C	505	6/6	0.95	0.15	-1.22	69,88,89,91	0
7	CA	E	200	1/1	0.93	0.07	-1.30	98,98,98,98	0
14	ACP	F	400	31/31	0.92	0.10	-1.47	71,84,123,146	0
11	6K9	B	503	52/52	0.96	0.11	-1.96	42,60,74,79	0
12	03S	B	504	5/5	0.81	0.16	-	51,102,113,114	0
6	MG	A	502	1/1	0.89	0.16	-	48,48,48,48	0
7	CA	C	504	1/1	0.91	0.15	-	117,117,117,117	0
6	MG	F	401	1/1	0.96	0.04	-	88,88,88,88	0
7	CA	B	502	1/1	0.99	0.26	-	96,96,96,96	0
6	MG	C	502	1/1	0.97	0.11	-	40,40,40,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.