



# Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 03:01 PM BST

PDB ID : 1JI9  
Title : Solution structure of the alpha-domain of mouse metallothionein-3  
Authors : Oz, G.; Zangger, K.; Armitage, I.M.  
Deposited on : 2001-07-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Mogul : unknown  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : rb-20027457  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 59%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	37	<div> <div></div> <div>22%</div> <div>51%</div> <div>27%</div> </div>

## 2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 498 atoms, of which 242 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called METALLOTHIONEIN-III.

Mol	Chain	Residues	Atoms						Trace
1	A	37	Total	C	H	N	O	S	0
			494	143	242	44	54	11	

- Molecule 2 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms	
2	A	4	Total	Cd
			4	4

## 4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

### • Molecule 1: METALLOTHIONEIN-III

Chain A: 



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *HYBRID DISTANCE GEOMETRY-DYNAMICAL SIMULATED ANNEALING*.

Of the 50 calculated structures, 1 were deposited, based on the following criterion: *Minimized average structure*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
X-PLOR	refinement	3.851

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5066
Number of chemical shift lists	1
Total number of shifts	240
Number of shifts mapped to atoms	240
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	59%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.15	0/252 (0.0%)	1.45	1/330 (0.3%)
All	All	1.15	0/252 (0.0%)	1.45	1/330 (0.3%)

There are no bond-length outliers.

All angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	46	ALA	N-CA-CB	-6.16	101.48	110.10

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	252	242	228	43
All	All	256	242	228	43

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 89.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:44:LYS:O	1:A:50:VAL:HG13	0.72	1.85
1:A:43:GLU:HA	1:A:46:ALA:HB2	0.68	1.64
1:A:44:LYS:O	1:A:50:VAL:CG1	0.60	2.50
1:A:61:ALA:O	1:A:62:GLU:CB	0.59	2.50
1:A:63:LYS:CD	1:A:63:LYS:C	0.58	2.72
1:A:34:CYS:O	1:A:35:CYS:SG	0.55	2.65
1:A:33:SER:O	1:A:35:CYS:N	0.55	2.40
1:A:61:ALA:O	1:A:62:GLU:HB2	0.54	2.02
1:A:43:GLU:HA	1:A:46:ALA:CB	0.54	2.33
1:A:60:GLU:O	1:A:62:GLU:N	0.53	2.41
1:A:44:LYS:HG3	1:A:67:CYS:CA	0.50	2.37
1:A:63:LYS:HD2	1:A:63:LYS:O	0.49	2.06
1:A:33:SER:O	1:A:34:CYS:SG	0.48	2.71
1:A:37:CYS:HA	1:A:63:LYS:HA	0.47	1.87
1:A:37:CYS:HB2	1:A:64:CYS:N	0.46	2.24
1:A:60:GLU:CG	1:A:61:ALA:N	0.46	2.78
1:A:44:LYS:HG3	1:A:67:CYS:CB	0.45	2.42
1:A:60:GLU:C	1:A:62:GLU:N	0.44	2.70
1:A:49:CYS:HB3	1:A:52:LYS:CG	0.44	2.42
1:A:67:CYS:O	1:A:68:GLN:C	0.44	2.55
1:A:59:ALA:O	1:A:60:GLU:C	0.44	2.55
1:A:61:ALA:C	1:A:62:GLU:CG	0.44	2.85
1:A:44:LYS:HG3	1:A:67:CYS:HA	0.44	1.89
1:A:35:CYS:CB	1:A:38:CYS:HB2	0.43	2.42
1:A:33:SER:O	1:A:34:CYS:C	0.43	2.57
1:A:60:GLU:O	1:A:62:GLU:HG2	0.43	2.13
1:A:63:LYS:CD	1:A:63:LYS:O	0.43	2.66
1:A:60:GLU:O	1:A:61:ALA:C	0.43	2.57
1:A:48:ASP:OD1	1:A:48:ASP:O	0.43	2.36
1:A:61:ALA:O	1:A:62:GLU:HG3	0.43	2.14
1:A:57:ALA:O	1:A:58:LYS:C	0.43	2.57
1:A:51:CYS:O	1:A:52:LYS:C	0.42	2.57
1:A:50:VAL:CG2	1:A:67:CYS:SG	0.42	3.07
1:A:44:LYS:HB3	1:A:67:CYS:SG	0.42	2.54
1:A:37:CYS:CA	1:A:63:LYS:HA	0.42	2.44
1:A:47:LYS:O	1:A:48:ASP:HB2	0.41	2.15
1:A:44:LYS:HG3	1:A:67:CYS:HB3	0.41	1.91
1:A:37:CYS:HA	1:A:62:GLU:O	0.41	2.16
1:A:42:CYS:SG	1:A:44:LYS:HB2	0.40	2.56
1:A:61:ALA:O	1:A:62:GLU:CG	0.40	2.69
1:A:37:CYS:HA	1:A:62:GLU:C	0.40	2.37
1:A:59:ALA:C	1:A:61:ALA:N	0.40	2.74
1:A:44:LYS:CB	1:A:67:CYS:SG	0.40	3.10



## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	35/37 (95%)	20 (57%)	11 (31%)	4 (11%)	1	8
All	All	35/37 (95%)	20 (57%)	11 (31%)	4 (11%)	1	8

All 4 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	48	ASP
1	A	61	ALA
1	A	49	CYS
1	A	34	CYS

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	29/29 (100%)	20 (69%)	9 (31%)	2	16
All	All	29/29 (100%)	20 (69%)	9 (31%)	2	16

All 9 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	37	CYS
1	A	32	LYS
1	A	66	CYS
1	A	35	CYS
1	A	58	LYS
1	A	63	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	36	SER
1	A	51	CYS
1	A	54	GLU

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 59% for the well-defined parts and 59% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 5066

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	240
Number of shifts mapped to atoms	240
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	0	—	—
$^{13}\text{C}_\beta$	0	—	—
$^{13}\text{C}'$	0	—	—
$^{15}\text{N}$	35	$0.04 \pm 0.39$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 59%, i.e. 224 atoms were assigned a chemical shift out of a possible 377. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	108/183 (59%)	73/73 (100%)	0/74 (0%)	35/36 (97%)
Sidechain	116/194 (60%)	116/118 (98%)	0/69 (0%)	0/7 (0%)

*Continued on next page...*

*Continued from previous page...*

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	224/377 (59%)	189/191 (99%)	0/143 (0%)	35/43 (81%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 59%, i.e. 224 atoms were assigned a chemical shift out of a possible 377. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	<b>Total</b>	<b><sup>1</sup>H</b>	<b><sup>13</sup>C</b>	<b><sup>15</sup>N</b>
Backbone	108/183 (59%)	73/73 (100%)	0/74 (0%)	35/36 (97%)
Sidechain	116/194 (60%)	116/118 (98%)	0/69 (0%)	0/7 (0%)
Aromatic	0/0 (—%)	0/0 (—%)	0/0 (—%)	0/0 (—%)
Overall	224/377 (59%)	189/191 (99%)	0/143 (0%)	35/43 (81%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

