



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:00 AM GMT

PDB ID : 2JJM
Title : CRYSTAL STRUCTURE OF A FAMILY GT4 GLYCOSYLTRANSFERASE
FROM BACILLUS ANTHRACIS ORF BA1558.
Authors : Ruane, K.M.; Davies, G.J.; Martinez-Fleites, C.
Deposited on : 2008-04-15
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	<div> <div>51%</div> <div>34%</div> <div>6% • 9%</div> </div>
1	B	394	<div> <div>51%</div> <div>34%</div> <div>5% • 9%</div> </div>
1	C	394	<div> <div>52%</div> <div>33%</div> <div>6% • 9%</div> </div>
1	D	394	<div> <div>53%</div> <div>32%</div> <div>6% • 9%</div> </div>
1	E	394	<div> <div>52%</div> <div>34%</div> <div>6% • 9%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	394	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>2%</div><div>52%</div><div>33%</div><div>5%</div><div>9%</div></div>
1	G	394	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>%</div><div>52%</div><div>34%</div><div>6%</div><div>9%</div></div>
1	H	394	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>14%</div><div>51%</div><div>34%</div><div>5%</div><div>9%</div></div>
1	I	394	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>3%</div><div>52%</div><div>33%</div><div>6%</div><div>9%</div></div>
1	J	394	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>2%</div><div>52%</div><div>33%</div><div>5%</div><div>9%</div></div>
1	K	394	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>2%</div><div>53%</div><div>33%</div><div>6%</div><div>9%</div></div>
1	L	394	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>%</div><div>52%</div><div>33%</div><div>6%</div><div>9%</div></div>

2 Entry composition


There is only 1 type of molecule in this entry. The entry contains 33924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	B	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	C	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	D	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	E	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	F	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	G	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	H	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	I	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	J	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	K	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			
1	L	360	Total	C	N	O	S	0	0	1
			2827	1806	477	531	13			


● Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN




● Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN




● Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN







• Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN




• Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN




• Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN



• Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN



• Molecule 1: GLYCOSYL TRANSFERASE, GROUP 1 FAMILY PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	134.60Å 204.67Å 135.33Å 90.00° 115.49° 90.00°	Depositor
Resolution (Å)	20.00 – 3.10 78.27 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.3 (20.00-3.10) 89.5 (78.27-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.69Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.254 , 0.262 0.252 , 0.260	Depositor DCC
R_{free} test set	5887 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	88.0	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 64.0	EDS
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 161831 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	33924	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.86	8/2873 (0.3%)	0.89	10/3884 (0.3%)
1	B	0.86	7/2873 (0.2%)	0.90	10/3884 (0.3%)
1	C	0.87	8/2873 (0.3%)	0.89	10/3884 (0.3%)
1	D	0.86	8/2873 (0.3%)	0.89	10/3884 (0.3%)
1	E	0.87	8/2873 (0.3%)	0.89	10/3884 (0.3%)
1	F	0.86	8/2873 (0.3%)	0.89	9/3884 (0.2%)
1	G	0.87	8/2873 (0.3%)	0.90	9/3884 (0.2%)
1	H	0.86	6/2873 (0.2%)	0.90	9/3884 (0.2%)
1	I	0.87	8/2873 (0.3%)	0.90	10/3884 (0.3%)
1	J	0.89	8/2873 (0.3%)	0.89	8/3884 (0.2%)
1	K	0.87	7/2873 (0.2%)	0.89	10/3884 (0.3%)
1	L	0.87	7/2873 (0.2%)	0.89	10/3884 (0.3%)
All	All	0.87	91/34476 (0.3%)	0.89	115/46608 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
1	D	0	2
1	E	0	2
1	F	0	2
1	G	0	2
1	H	0	2
1	I	0	3
1	J	0	2
1	K	0	2
1	L	0	2
All	All	0	25

The worst 5 of 91 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	J	287	VAL	CA-CB	-7.51	1.39	1.54
1	F	287	VAL	CA-CB	-7.43	1.39	1.54
1	D	287	VAL	CA-CB	-7.41	1.39	1.54
1	I	287	VAL	CA-CB	-7.41	1.39	1.54
1	G	287	VAL	CA-CB	-7.39	1.39	1.54

The worst 5 of 115 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	121	GLY	N-CA-C	9.49	136.84	113.10
1	J	121	GLY	N-CA-C	9.48	136.80	113.10
1	H	121	GLY	N-CA-C	9.46	136.74	113.10
1	B	121	GLY	N-CA-C	9.45	136.72	113.10
1	D	121	GLY	N-CA-C	9.44	136.70	113.10

There are no chirality outliers.

5 of 25 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	284	PHE	Mainchain,Peptide
1	B	284	PHE	Mainchain,Peptide
1	C	284	PHE	Mainchain,Peptide
1	D	284	PHE	Mainchain,Peptide
1	E	284	PHE	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2827	0	2856	182	1
1	B	2827	0	2856	191	0
1	C	2827	0	2856	188	0
1	D	2827	0	2856	182	0
1	E	2827	0	2856	182	0
1	F	2827	0	2856	185	0
1	G	2827	0	2856	195	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	2827	0	2856	187	0
1	I	2827	0	2856	189	0
1	J	2827	0	2856	183	0
1	K	2827	0	2856	186	1
1	L	2827	0	2856	190	0
All	All	33924	0	34272	2180	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 2180 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:167:ASP:OD2	1:L:160:GLU:HA	1.15	1.27
1:A:286:LEU:HD12	1:A:286:LEU:N	1.55	1.17
1:B:286:LEU:CD1	1:B:286:LEU:H	1.54	1.16
1:K:280:GLU:O	1:K:281:LYS:HG2	1.46	1.16
1:G:280:GLU:O	1:G:281:LYS:HG2	1.47	1.15

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:GLU:CA	1:K:167:ASP:OD2[1_455]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	350/394 (89%)	323 (92%)	24 (7%)	3 (1%)	21 61
1	B	350/394 (89%)	322 (92%)	26 (7%)	2 (1%)	30 68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	350/394 (89%)	323 (92%)	25 (7%)	2 (1%)	30	68
1	D	350/394 (89%)	322 (92%)	25 (7%)	3 (1%)	21	61
1	E	350/394 (89%)	323 (92%)	24 (7%)	3 (1%)	21	61
1	F	350/394 (89%)	322 (92%)	26 (7%)	2 (1%)	30	68
1	G	350/394 (89%)	322 (92%)	26 (7%)	2 (1%)	30	68
1	H	350/394 (89%)	322 (92%)	26 (7%)	2 (1%)	30	68
1	I	350/394 (89%)	322 (92%)	25 (7%)	3 (1%)	21	61
1	J	350/394 (89%)	322 (92%)	26 (7%)	2 (1%)	30	68
1	K	350/394 (89%)	322 (92%)	26 (7%)	2 (1%)	30	68
1	L	350/394 (89%)	323 (92%)	25 (7%)	2 (1%)	30	68
All	All	4200/4728 (89%)	3868 (92%)	304 (7%)	28 (1%)	26	65

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	293	ALA
1	A	304	GLY
1	B	293	ALA
1	B	304	GLY
1	C	293	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	B	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	C	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	D	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	E	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	F	312/346 (90%)	287 (92%)	25 (8%)	15	48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	H	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	I	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	J	312/346 (90%)	286 (92%)	26 (8%)	14	46
1	K	312/346 (90%)	287 (92%)	25 (8%)	15	48
1	L	312/346 (90%)	287 (92%)	25 (8%)	15	48
All	All	3744/4152 (90%)	3443 (92%)	301 (8%)	15	48

5 of 301 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	183	ARG
1	G	278	LEU
1	L	65	PHE
1	F	255	ARG
1	G	65	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 111 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	173	ASN
1	G	330	GLN
1	L	134	ASN
1	F	214	GLN
1	G	98	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/394 (91%)	0.04	3 (0%) 87 75	62, 92, 116, 125	0
1	B	360/394 (91%)	0.25	18 (5%) 32 13	63, 94, 116, 126	0
1	C	360/394 (91%)	0.21	9 (2%) 61 37	57, 93, 116, 125	0
1	D	360/394 (91%)	0.06	2 (0%) 90 80	58, 93, 115, 126	0
1	E	360/394 (91%)	0.15	7 (1%) 70 48	56, 92, 116, 125	0
1	F	360/394 (91%)	0.17	9 (2%) 61 37	57, 93, 116, 125	0
1	G	360/394 (91%)	0.07	4 (1%) 82 66	62, 93, 116, 125	0
1	H	360/394 (91%)	0.71	54 (15%) 3 1	63, 95, 117, 126	0
1	I	360/394 (91%)	0.25	10 (2%) 56 32	57, 92, 116, 125	0
1	J	360/394 (91%)	0.15	6 (1%) 73 52	38, 87, 116, 124	0
1	K	360/394 (91%)	0.18	9 (2%) 61 37	55, 90, 116, 124	0
1	L	360/394 (91%)	0.11	4 (1%) 82 66	61, 92, 115, 124	0
All	All	4320/4728 (91%)	0.20	135 (3%) 52 28	38, 92, 116, 126	0

The worst 5 of 135 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	177	GLU	5.3
1	H	190	LYS	4.5
1	H	257	LEU	4.5
1	H	272	SER	4.2
1	H	263	ASP	4.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.