



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 08:17 PM GMT

PDB ID : 1JN0
Title : Crystal structure of the non-regulatory A4 isoform of spinach chloroplast glyceraldehyde-3-phosphate dehydrogenase complexed with NADP
Authors : Fermani, S.; Ripamonti, A.; Sabatino, P.; Zanotti, G.; Scagliarini, S.; Sparla, F.; Trost, P.; Pupillo, P.
Deposited on : 2001-07-21
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	335	
1	B	335	
1	O	335	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	O	334	Total	C	N	O	S	0	0	0
			2521	1589	439	482	11			
1	A	334	Total	C	N	O	S	0	0	0
			2521	1589	439	482	11			
1	B	334	Total	C	N	O	S	0	0	0
			2521	1589	439	482	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	?	-	ARG	DELETION	UNP P19866
O	?	-	ASP	DELETION	UNP P19866
A	?	-	ARG	DELETION	UNP P19866
A	?	-	ASP	DELETION	UNP P19866
B	?	-	ARG	DELETION	UNP P19866
B	?	-	ASP	DELETION	UNP P19866

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	O	1	Total	O	S	0	0
			5	4	1		
2	O	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	O	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	65	Total	O	0	0
			65	65		
4	B	60	Total	O	0	0
			60	60		
4	O	87	Total	O	0	0
			87	87		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE A



• Molecule 1: GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE A



● Molecule 1: GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE A



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	146.25Å 182.18Å 106.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.58 – 3.00	Depositor
% Data completeness (in resolution range)	93.0 (49.58-3.00)	Depositor
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.213 , 0.254	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7949	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	3/2562 (0.1%)	0.76	2/3479 (0.1%)
1	B	0.52	2/2562 (0.1%)	0.75	5/3479 (0.1%)
1	O	0.65	4/2562 (0.2%)	0.84	8/3479 (0.2%)
All	All	0.58	9/7686 (0.1%)	0.78	15/10437 (0.1%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	O	266	GLU	CD-OE1	-11.03	1.13	1.25
1	O	333	GLN	CD-OE1	-7.77	1.06	1.24
1	A	333	GLN	CD-OE1	-6.64	1.09	1.24
1	A	333	GLN	C-O	-6.21	1.11	1.23
1	B	333	GLN	CD-OE1	-5.85	1.11	1.24

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	19	TRP	CA-CB-CG	12.31	137.08	113.70
1	O	19	TRP	CB-CG-CD1	-9.65	114.45	127.00
1	O	19	TRP	CB-CG-CD2	8.50	137.65	126.60
1	O	333	GLN	CB-CA-C	-6.22	97.96	110.40
1	B	121	PRO	CA-N-CD	-5.99	103.11	111.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2521	0	2558	256	0
1	B	2521	0	2561	232	0
1	O	2521	0	2562	240	0
2	A	10	0	0	1	0
2	B	10	0	0	0	0
2	O	10	0	0	0	0
3	A	48	0	26	5	0
3	B	48	0	26	5	0
3	O	48	0	26	7	0
4	A	65	0	0	8	0
4	B	60	0	0	9	0
4	O	87	0	0	11	0
All	All	7949	0	7759	718	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 46.

The worst 5 of 718 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:56:ASP:H	1:O:66:VAL:CG1	1.21	1.50
1:A:56:ASP:H	1:A:66:VAL:CG1	1.25	1.47
1:B:56:ASP:N	1:B:66:VAL:HG12	1.45	1.31
1:B:56:ASP:H	1:B:66:VAL:CG1	1.46	1.28
1:O:56:ASP:N	1:O:66:VAL:CG1	2.04	1.19

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/335 (99%)	284 (86%)	36 (11%)	12 (4%)	4	24
1	B	332/335 (99%)	281 (85%)	38 (11%)	13 (4%)	4	21
1	O	332/335 (99%)	281 (85%)	38 (11%)	13 (4%)	4	21
All	All	996/1005 (99%)	846 (85%)	112 (11%)	38 (4%)	4	22

5 of 38 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	O	24	PRO
1	O	66	VAL
1	A	24	PRO
1	A	66	VAL
1	A	332	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/277 (100%)	244 (88%)	33 (12%)	6	26
1	B	277/277 (100%)	246 (89%)	31 (11%)	7	29
1	O	277/277 (100%)	239 (86%)	38 (14%)	4	20
All	All	831/831 (100%)	729 (88%)	102 (12%)	6	25

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	ASP
1	A	188	SER
1	B	250	THR
1	A	64	ILE
1	A	94	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	ASN
1	A	163	GLN
1	B	163	GLN
1	A	81	ASN
1	B	133	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NDP	A	1335	2	42,52,52	1.36	4 (9%)	55,80,80	2.28	12 (21%)
2	SO4	A	1336	3	4,4,4	0.25	0	6,6,6	0.08	0
2	SO4	A	1337	-	4,4,4	0.16	0	6,6,6	0.11	0
3	NDP	B	2335	-	42,52,52	1.36	4 (9%)	55,80,80	2.27	12 (21%)
2	SO4	B	2336	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	B	2337	-	4,4,4	0.25	0	6,6,6	0.19	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NDP	O	6335	-	42,52,52	1.36	4 (9%)	55,80,80	2.27	12 (21%)
2	SO4	O	6336	-	4,4,4	0.36	0	6,6,6	0.10	0
2	SO4	O	6337	-	4,4,4	0.14	0	6,6,6	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDP	A	1335	2	-	0/30/77/77	0/5/5/5
2	SO4	A	1336	3	-	0/0/0/0	0/0/0/0
2	SO4	A	1337	-	-	0/0/0/0	0/0/0/0
3	NDP	B	2335	-	-	0/30/77/77	0/5/5/5
2	SO4	B	2336	-	-	0/0/0/0	0/0/0/0
2	SO4	B	2337	-	-	0/0/0/0	0/0/0/0
3	NDP	O	6335	-	-	0/30/77/77	0/5/5/5
2	SO4	O	6336	-	-	0/0/0/0	0/0/0/0
2	SO4	O	6337	-	-	0/0/0/0	0/0/0/0

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1335	NDP	C4N-C5N	-4.38	1.39	1.49
3	O	6335	NDP	C4N-C5N	-4.38	1.39	1.49
3	B	2335	NDP	C4N-C5N	-4.34	1.39	1.49
3	B	2335	NDP	C2N-C3N	2.71	1.41	1.34
3	O	6335	NDP	C2N-C3N	2.73	1.41	1.34

The worst 5 of 36 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1335	NDP	N3A-C2A-N1A	-8.79	122.17	128.89
3	O	6335	NDP	N3A-C2A-N1A	-8.76	122.19	128.89
3	B	2335	NDP	N3A-C2A-N1A	-8.76	122.19	128.89
3	A	1335	NDP	C4B-O4B-C1B	-8.64	100.22	109.72
3	B	2335	NDP	C4B-O4B-C1B	-8.60	100.27	109.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1335	NDP	5	0
2	A	1336	SO4	1	0
3	B	2335	NDP	5	0
3	O	6335	NDP	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.