



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 08:17 PM GMT

PDB ID : 1JND  
Title : Crystal structure of imaginal disc growth factor-2  
Authors : Varela, P.F.; Llera, A.S.; Mariuzza, R.A.; Tormo, J.  
Deposited on : 2001-07-23  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1475 (1.34-1.26)
Clashscore	102246	1031 (1.32-1.28)
Ramachandran outliers	100387	1504 (1.34-1.26)
Sidechain outliers	100360	1503 (1.34-1.26)
RSRZ outliers	91569	1476 (1.34-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3916 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Imaginal disc growth factor-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	400	Total	C	N	O	S	0	0	0
			3166	2025	536	600	5			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	257	PHE	VAL	CONFLICT	GB 4092089

- Molecule 2 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 3 is water (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	700	Total	O	0	0
			700	700		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Imaginal disc growth factor-2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.34 Å   106.34 Å   89.99 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	100.00 – 1.30 45.78 – 1.30	Depositor EDS
% Data completeness (in resolution range)	98.2 (100.00-1.30) 90.3 (45.78-1.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.02	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.16 (at 1.30 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.176 , 0.202 0.167 , 0.193	Depositor DCC
$R_{free}$ test set	2302 reflections (1.80%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.7	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 59.6	EDS
Estimated twinning fraction	0.016 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 129904 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3916	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.84	1/3248 (0.0%)	1.41	36/4409 (0.8%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	420	LEU	C-OXT	20.34	1.61	1.23

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	139	ARG	NE-CZ-NH2	-9.63	115.49	120.30
1	A	418	TYR	CB-CG-CD2	9.63	126.78	121.00
1	A	229	ARG	NE-CZ-NH2	-9.03	115.78	120.30
1	A	414	ARG	NE-CZ-NH1	-8.78	115.91	120.30
1	A	418	TYR	CB-CG-CD1	-8.24	116.06	121.00
1	A	308	PHE	CB-CG-CD2	-8.00	115.20	120.80
1	A	419	ARG	NE-CZ-NH1	7.38	123.99	120.30
1	A	27	GLU	OE1-CD-OE2	7.33	132.10	123.30
1	A	349	ARG	NE-CZ-NH1	-6.96	116.82	120.30
1	A	408	ASP	CB-CG-OD1	-6.79	112.19	118.30
1	A	97	TYR	CB-CG-CD1	6.74	125.04	121.00
1	A	249	ARG	NE-CZ-NH2	6.51	123.56	120.30
1	A	44	ARG	NE-CZ-NH1	6.25	123.43	120.30
1	A	117	GLU	CB-CG-CD	-6.11	97.72	114.20
1	A	384	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	15	ARG	NE-CZ-NH2	-6.06	117.27	120.30
1	A	128	ASP	CB-CG-OD1	6.00	123.70	118.30
1	A	249	ARG	NE-CZ-NH1	-6.00	117.30	120.30
1	A	206	ASP	CB-CG-OD2	-5.77	113.11	118.30
1	A	122	TYR	CB-CG-CD1	-5.70	117.58	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	182	ASP	CB-CG-OD1	5.68	123.41	118.30
1	A	122	TYR	CB-CG-CD2	5.65	124.39	121.00
1	A	125	ASP	CB-CG-OD2	5.39	123.15	118.30
1	A	38	TYR	CB-CG-CD1	-5.36	117.78	121.00
1	A	356	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	A	51	TYR	CB-CG-CD1	5.34	124.20	121.00
1	A	229	ARG	NE-CZ-NH1	5.27	122.93	120.30
1	A	131	TYR	CB-CG-CD1	-5.25	117.85	121.00
1	A	58	ASP	CB-CG-OD1	-5.23	113.59	118.30
1	A	60	TYR	CB-CG-CD2	-5.16	117.90	121.00
1	A	185	ARG	NE-CZ-NH1	-5.13	117.73	120.30
1	A	97	TYR	CB-CG-CD2	-5.12	117.93	121.00
1	A	320	GLU	OE1-CD-OE2	-5.12	117.16	123.30
1	A	187	ASP	CB-CG-OD2	-5.11	113.70	118.30
1	A	9	ASP	CB-CG-OD2	-5.11	113.70	118.30
1	A	395	ASP	CB-CG-OD2	-5.01	113.79	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3166	0	3081	35	0
2	A	39	0	33	0	0
3	A	11	0	10	0	0
4	A	700	0	0	24	2
All	All	3916	0	3124	35	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:LYS:HE2	4:A:1480:HOH:O	0.95	1.13
1:A:140:LYS:CE	4:A:1278:HOH:O	1.99	1.07
1:A:140:LYS:HE3	4:A:1278:HOH:O	1.53	1.04
1:A:378:ASN:HB2	4:A:1479:HOH:O	1.70	0.89
1:A:161:ILE:HD13	4:A:1640:HOH:O	1.89	0.72
1:A:161:ILE:CD1	4:A:1640:HOH:O	2.38	0.70
1:A:102:GLU:OE2	1:A:169:HIS:HE1	1.74	0.70
1:A:163:ASP:OD2	1:A:169:HIS:HD2	1.76	0.69
1:A:107:ARG:HG2	1:A:107:ARG:HH11	1.59	0.67
1:A:55:GLU:OE1	4:A:1325:HOH:O	2.14	0.66
1:A:140:LYS:O	1:A:141:VAL:HB	1.95	0.65
1:A:248:ASP:CB	4:A:1491:HOH:O	2.43	0.65
1:A:374:ASP:OD2	4:A:1560:HOH:O	2.15	0.63
1:A:2:SER:N	4:A:1694:HOH:O	2.31	0.63
1:A:140:LYS:HE2	4:A:1278:HOH:O	1.81	0.63
1:A:92:ASP:OD1	4:A:1619:HOH:O	2.15	0.62
1:A:107:ARG:NH1	1:A:107:ARG:HG2	2.13	0.61
1:A:165:HIS:CD2	4:A:1595:HOH:O	2.56	0.58
1:A:242:HIS:HD2	4:A:1007:HOH:O	1.87	0.57
1:A:106:VAL:HG11	4:A:1609:HOH:O	2.06	0.55
1:A:248:ASP:HB3	4:A:1491:HOH:O	2.04	0.55
1:A:106:VAL:CG1	4:A:1609:HOH:O	2.57	0.53
1:A:161:ILE:HD11	4:A:1648:HOH:O	2.08	0.53
1:A:30:LEU:HD11	1:A:70:LEU:HD13	1.92	0.52
1:A:248:ASP:CG	4:A:1491:HOH:O	2.51	0.48
1:A:161:ILE:HD11	4:A:1640:HOH:O	2.11	0.48
1:A:140:LYS:HD3	1:A:140:LYS:HA	1.69	0.47
1:A:242:HIS:HE1	4:A:1158:HOH:O	1.98	0.46
1:A:107:ARG:CG	1:A:107:ARG:HH11	2.25	0.46
1:A:248:ASP:OD1	4:A:1506:HOH:O	2.21	0.44
1:A:185:ARG:HD3	4:A:1628:HOH:O	2.18	0.43
1:A:46:GLU:HG2	1:A:47:ASN:N	2.34	0.43
1:A:23:ASN:N	1:A:24:PRO:CD	2.83	0.42
1:A:247:LYS:O	1:A:248:ASP:HB2	2.21	0.41
1:A:55:GLU:HG3	4:A:1147:HOH:O	2.21	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1412:HOH:O	4:A:1567:HOH:O[6_765]	2.13	0.07
4:A:1371:HOH:O	4:A:1388:HOH:O[5_665]	2.15	0.05

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/420 (94%)	391 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	341/356 (96%)	335 (98%)	6 (2%)	66	25

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	LYS
1	A	107	ARG
1	A	140	LYS
1	A	161	ILE
1	A	270	ASN
1	A	395	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	HIS
1	A	169	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	242	HIS
1	A	270	ASN
1	A	311	GLN
1	A	331	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	A	3100	1,2	14,14,15	0.89	0	15,19,21	1.30	1 (6%)
2	NAG	A	3101	2	14,14,15	1.02	1 (7%)	15,19,21	1.48	3 (20%)
2	BMA	A	3102	3,2	11,11,12	0.75	0	14,15,17	1.38	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	3100	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	3101	2	-	0/6/23/26	0/1/1/1
2	BMA	A	3102	3,2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	3101	NAG	C1-C2	2.13	1.55	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3102	BMA	C1-O5-C5	-4.00	107.17	112.25
2	A	3100	NAG	C4-C3-C2	-3.10	106.41	111.23
2	A	3101	NAG	C2-N2-C7	-2.69	119.59	123.04
2	A	3101	NAG	C8-C7-N2	-2.56	111.21	116.11
2	A	3101	NAG	O7-C7-N2	2.58	127.12	121.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MAN	A	3103	2	11,11,12	0.76	0	14,15,17	1.96	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	A	3103	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
3	A	3103	MAN	C2-C3-C4	-3.72	104.72	111.04
3	A	3103	MAN	O3-C3-C2	-2.60	105.30	110.00
3	A	3103	MAN	O2-C2-C3	-2.21	105.68	110.12
3	A	3103	MAN	C3-C4-C5	-2.06	106.61	110.20
3	A	3103	MAN	O5-C1-C2	2.06	114.20	110.86
3	A	3103	MAN	C1-O5-C5	3.49	116.68	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	400/420 (95%)	-0.08	7 (1%) 71 69	8, 12, 27, 70	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	VAL	6.2
1	A	418	TYR	3.9
1	A	75	PRO	3.5
1	A	72	ARG	3.4
1	A	106	VAL	3.3
1	A	161	ILE	2.9
1	A	140	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	3100	14/15	0.98	0.05	-1.11	10,12,15,16	0
2	NAG	A	3101	14/15	0.93	0.14	-	16,21,26,28	0
2	BMA	A	3102	11/12	0.78	0.23	-	26,33,42,45	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MAN	A	3103	11/12	0.54	0.33	-	44,46,58,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.