



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:48 AM GMT

PDB ID : 3JQE  
Title : Crystal structure of pteridine reductase 1 (PTR1) from Trypanosoma brucei in ternary complex with cofactor (NADP+) and inhibitor 2-amino-6-(4-methoxyphenyl)-4-oxo-4,7-dihydro-3H-pyrrolo[2,3-d]pyrimidine-5-carbonitrile (DX8)  
Authors : Tulloch, L.B.; Hunter, W.N.  
Deposited on : 2009-09-06  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	<div> <div></div> <div>79% 7% 14%</div> </div>
1	B	288	<div> <div></div> <div>79% 7% 14%</div> </div>
1	C	288	<div> <div>2%</div> <div>80% 6% 14%</div> </div>
1	D	288	<div> <div></div> <div>78% 7% 14%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DX8	A	270	-	-	-	X
3	DX8	B	270	-	-	-	X
3	DX8	C	270	-	-	-	X
3	DX8	D	270	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8639 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pteridine reductase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	7	0
			1878	1181	328	358	11			
1	B	248	Total	C	N	O	S	0	15	0
			1898	1198	330	359	11			
1	C	249	Total	C	N	O	S	0	9	0
			1868	1177	328	352	11			
1	D	249	Total	C	N	O	S	0	11	0
			1888	1189	329	359	11			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP Q581W1
A	-18	GLY	-	EXPRESSION TAG	UNP Q581W1
A	-17	SER	-	EXPRESSION TAG	UNP Q581W1
A	-16	SER	-	EXPRESSION TAG	UNP Q581W1
A	-15	HIS	-	EXPRESSION TAG	UNP Q581W1
A	-14	HIS	-	EXPRESSION TAG	UNP Q581W1
A	-13	HIS	-	EXPRESSION TAG	UNP Q581W1
A	-12	HIS	-	EXPRESSION TAG	UNP Q581W1
A	-11	HIS	-	EXPRESSION TAG	UNP Q581W1
A	-10	HIS	-	EXPRESSION TAG	UNP Q581W1
A	-9	SER	-	EXPRESSION TAG	UNP Q581W1
A	-8	SER	-	EXPRESSION TAG	UNP Q581W1
A	-7	GLY	-	EXPRESSION TAG	UNP Q581W1
A	-6	LEU	-	EXPRESSION TAG	UNP Q581W1
A	-5	VAL	-	EXPRESSION TAG	UNP Q581W1
A	-4	PRO	-	EXPRESSION TAG	UNP Q581W1
A	-3	ARG	-	EXPRESSION TAG	UNP Q581W1
A	-2	GLY	-	EXPRESSION TAG	UNP Q581W1
A	-1	SER	-	EXPRESSION TAG	UNP Q581W1
A	0	HIS	-	EXPRESSION TAG	UNP Q581W1
B	-19	MET	-	EXPRESSION TAG	UNP Q581W1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	GLY	-	EXPRESSION TAG	UNP Q581W1
B	-17	SER	-	EXPRESSION TAG	UNP Q581W1
B	-16	SER	-	EXPRESSION TAG	UNP Q581W1
B	-15	HIS	-	EXPRESSION TAG	UNP Q581W1
B	-14	HIS	-	EXPRESSION TAG	UNP Q581W1
B	-13	HIS	-	EXPRESSION TAG	UNP Q581W1
B	-12	HIS	-	EXPRESSION TAG	UNP Q581W1
B	-11	HIS	-	EXPRESSION TAG	UNP Q581W1
B	-10	HIS	-	EXPRESSION TAG	UNP Q581W1
B	-9	SER	-	EXPRESSION TAG	UNP Q581W1
B	-8	SER	-	EXPRESSION TAG	UNP Q581W1
B	-7	GLY	-	EXPRESSION TAG	UNP Q581W1
B	-6	LEU	-	EXPRESSION TAG	UNP Q581W1
B	-5	VAL	-	EXPRESSION TAG	UNP Q581W1
B	-4	PRO	-	EXPRESSION TAG	UNP Q581W1
B	-3	ARG	-	EXPRESSION TAG	UNP Q581W1
B	-2	GLY	-	EXPRESSION TAG	UNP Q581W1
B	-1	SER	-	EXPRESSION TAG	UNP Q581W1
B	0	HIS	-	EXPRESSION TAG	UNP Q581W1
C	-19	MET	-	EXPRESSION TAG	UNP Q581W1
C	-18	GLY	-	EXPRESSION TAG	UNP Q581W1
C	-17	SER	-	EXPRESSION TAG	UNP Q581W1
C	-16	SER	-	EXPRESSION TAG	UNP Q581W1
C	-15	HIS	-	EXPRESSION TAG	UNP Q581W1
C	-14	HIS	-	EXPRESSION TAG	UNP Q581W1
C	-13	HIS	-	EXPRESSION TAG	UNP Q581W1
C	-12	HIS	-	EXPRESSION TAG	UNP Q581W1
C	-11	HIS	-	EXPRESSION TAG	UNP Q581W1
C	-10	HIS	-	EXPRESSION TAG	UNP Q581W1
C	-9	SER	-	EXPRESSION TAG	UNP Q581W1
C	-8	SER	-	EXPRESSION TAG	UNP Q581W1
C	-7	GLY	-	EXPRESSION TAG	UNP Q581W1
C	-6	LEU	-	EXPRESSION TAG	UNP Q581W1
C	-5	VAL	-	EXPRESSION TAG	UNP Q581W1
C	-4	PRO	-	EXPRESSION TAG	UNP Q581W1
C	-3	ARG	-	EXPRESSION TAG	UNP Q581W1
C	-2	GLY	-	EXPRESSION TAG	UNP Q581W1
C	-1	SER	-	EXPRESSION TAG	UNP Q581W1
C	0	HIS	-	EXPRESSION TAG	UNP Q581W1
D	-19	MET	-	EXPRESSION TAG	UNP Q581W1
D	-18	GLY	-	EXPRESSION TAG	UNP Q581W1
D	-17	SER	-	EXPRESSION TAG	UNP Q581W1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	EXPRESSION TAG	UNP Q581W1
D	-15	HIS	-	EXPRESSION TAG	UNP Q581W1
D	-14	HIS	-	EXPRESSION TAG	UNP Q581W1
D	-13	HIS	-	EXPRESSION TAG	UNP Q581W1
D	-12	HIS	-	EXPRESSION TAG	UNP Q581W1
D	-11	HIS	-	EXPRESSION TAG	UNP Q581W1
D	-10	HIS	-	EXPRESSION TAG	UNP Q581W1
D	-9	SER	-	EXPRESSION TAG	UNP Q581W1
D	-8	SER	-	EXPRESSION TAG	UNP Q581W1
D	-7	GLY	-	EXPRESSION TAG	UNP Q581W1
D	-6	LEU	-	EXPRESSION TAG	UNP Q581W1
D	-5	VAL	-	EXPRESSION TAG	UNP Q581W1
D	-4	PRO	-	EXPRESSION TAG	UNP Q581W1
D	-3	ARG	-	EXPRESSION TAG	UNP Q581W1
D	-2	GLY	-	EXPRESSION TAG	UNP Q581W1
D	-1	SER	-	EXPRESSION TAG	UNP Q581W1
D	0	HIS	-	EXPRESSION TAG	UNP Q581W1

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is 2-AMINO-6-(4-METHOXYPHENYL)-4-OXO-4,7-DIHYDRO-3H-PYRROLO [2,3-D]PYRIMIDINE-5-CARBONITRILE (three-letter code: DX8) (formula: C<sub>14</sub>H<sub>11</sub>N<sub>5</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			21	14	5	2		
3	B	1	Total	C	N	O	0	0
			21	14	5	2		
3	C	1	Total	C	N	O	0	0
			21	14	5	2		
3	D	1	Total	C	N	O	0	0
			21	14	5	2		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	205	Total	O	0	0
			205	205		
5	B	208	Total	O	0	0
			208	208		
5	C	220	Total	O	0	0
			220	220		
5	D	194	Total	O	0	0
			194	194		




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Pteridine reductase 1




#### • Molecule 1: Pteridine reductase 1



#### • Molecule 1: Pteridine reductase 1



#### • Molecule 1: Pteridine reductase 1



PRO	ASN	CYS	THR	SER	S152	Y164	C168	H179	Y206	E217	K220	K221	K222	K223	K224	K229	K230	E231	Q250	Y251	L265	A268
-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.58 Å 89.42 Å 82.37 Å 90.00° 115.56° 90.00°	Depositor
Resolution (Å)	34.32 – 2.00 34.31 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.2 (34.32-2.00) 94.1 (34.31-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.79 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.152 , 0.213 0.157 , 0.212	Depositor DCC
$R_{free}$ test set	3179 reflections (5.40%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtriage
Anisotropy	1.076	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.1	EDS
Estimated twinning fraction	0.025 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	2 of 62100 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8639	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.71 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.4671e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, DX8, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/1920	0.61	0/2604
1	B	0.60	0/1955	0.62	0/2652
1	C	0.59	0/1904	0.62	0/2582
1	D	0.57	0/1933	0.62	0/2621
All	All	0.59	0/7712	0.62	0/10459

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1878	0	1906	20	0
1	B	1898	0	1948	17	0
1	C	1868	0	1906	13	0
1	D	1888	0	1925	20	0
2	A	48	0	25	0	0
2	B	48	0	25	1	0
2	C	48	0	25	0	0
2	D	48	0	25	0	0
3	A	21	0	11	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	21	0	11	2	0
3	C	21	0	11	5	0
3	D	21	0	11	4	0
4	A	4	0	3	0	0
5	A	205	0	0	0	0
5	B	208	0	0	6	0
5	C	220	0	0	3	0
5	D	194	0	0	8	0
All	All	8639	0	7832	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:168:CYS:SG	3:B:270:DX8:HAAB	2.19	0.81
1:C:168:CYS:SG	3:C:270:DX8:HAAB	2.20	0.81
1:C:75:GLU:OE1	5:C:484:HOH:O	2.00	0.79
3:D:270:DX8:CAE	3:D:270:DX8:HAH	2.12	0.79
1:A:168:CYS:SG	3:A:270:DX8:HAAB	2.24	0.78
1:D:222:ARG:HD3	1:D:231:GLU:OE2	1.85	0.75
3:C:270:DX8:HAG	5:C:427:HOH:O	1.85	0.74
1:D:168:CYS:SG	3:D:270:DX8:HAAB	2.27	0.74
1:C:75:GLU:HG2	5:C:364:HOH:O	1.92	0.68
1:A:247:GLY:HA2	1:A:250[B]:GLN:HG3	1.75	0.67
1:A:194:PRO:HG3	5:B:450:HOH:O	1.96	0.66
1:D:47:GLU:OE2	5:D:309:HOH:O	2.15	0.64
1:B:153:ASN:N	5:B:362:HOH:O	2.32	0.63
1:A:22:LYS:HG2	1:A:242:ILE:HG13	1.81	0.62
1:A:250[A]:GLN:HG2	5:B:271:HOH:O	1.98	0.62
1:D:217:GLU:HG3	5:D:299:HOH:O	2.01	0.61
3:A:270:DX8:HAH	3:A:270:DX8:CAE	2.31	0.61
3:C:270:DX8:HAH	3:C:270:DX8:CAE	2.32	0.60
1:B:164:VAL:HG22	1:B:179:HIS:CD2	2.36	0.60
1:A:161:ASP:HB3	1:A:164:VAL:HG13	1.85	0.59
1:C:9:THR:HA	1:C:33:HIS:HB3	1.87	0.56
1:A:9:THR:HA	1:A:33:HIS:HB3	1.86	0.56
1:D:9:THR:HA	1:D:33:HIS:HB3	1.88	0.55
1:A:250[A]:GLN:CD	1:B:236:GLN:HE21	2.09	0.55
1:B:223:ARG:HA	1:B:229:ARG:HG3	1.89	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:9:THR:HA	1:B:33:HIS:HB3	1.89	0.54
1:D:140:GLN:HG2	5:D:441:HOH:O	2.06	0.54
3:D:270:DX8:HAH	3:D:270:DX8:NAB	2.24	0.53
3:B:270:DX8:CAE	3:B:270:DX8:HAH	2.39	0.52
1:D:53:SER:CB	5:D:355:HOH:O	2.57	0.52
1:D:229:ARG:NH2	5:D:357:HOH:O	2.42	0.52
1:B:62:ASP:OD1	1:B:64[B]:THR:HG23	2.10	0.52
1:A:233:SER:H	1:A:236:GLN:HE21	1.57	0.52
1:C:232:ALA:HB2	1:D:251:TYR:CE2	2.45	0.52
1:D:132:PHE:CZ	1:D:136:MET:CE	2.94	0.51
1:D:250:GLN:HG3	5:D:443:HOH:O	2.12	0.50
1:A:103:VAL:HG21	1:C:195:TYR:OH	2.13	0.49
3:D:270:DX8:CAE	3:D:270:DX8:CAH	2.82	0.48
1:D:164:VAL:HG22	1:D:179:HIS:CD2	2.48	0.48
1:D:53:SER:HB2	5:D:355:HOH:O	2.13	0.47
3:C:270:DX8:CAH	3:C:270:DX8:CAE	2.92	0.47
1:B:64[B]:THR:CG2	1:B:122:GLU:HG2	2.45	0.47
1:A:193:ALA:N	1:A:194:PRO:CD	2.77	0.47
1:A:95:SER:CB	3:A:270:DX8:HNAC	2.28	0.46
1:B:229:ARG:NH1	5:B:450:HOH:O	2.48	0.46
1:D:220:LYS:O	1:D:224[A]:LYS:HG3	2.16	0.46
1:D:65:ASN:HA	1:D:69:LEU:HD22	1.97	0.46
1:A:67[A]:ASN:HA	1:C:117:GLU:HG3	1.98	0.46
1:C:232:ALA:HB2	1:D:251:TYR:CD2	2.52	0.45
1:B:160:CYS:HB3	5:B:383:HOH:O	2.16	0.45
1:C:210:PRO:HD3	3:C:270:DX8:NAB	2.32	0.45
1:A:254:GLY:HA3	1:B:265:LEU:HD11	1.99	0.45
1:A:65:ASN:HA	1:A:69:LEU:HD22	1.99	0.45
1:A:95:SER:HB3	3:A:270:DX8:NAC	2.32	0.44
1:C:138:PHE:O	1:C:142:GLN:HG2	2.17	0.44
1:D:132:PHE:CZ	1:D:136:MET:HE1	2.53	0.43
1:B:223:ARG:NH1	5:B:335:HOH:O	2.50	0.43
1:A:251:TYR:CE2	1:B:232:ALA:HB2	2.54	0.43
1:A:233:SER:H	1:A:236:GLN:NE2	2.17	0.43
1:A:232:ALA:HB2	1:B:251:TYR:CE2	2.54	0.42
1:B:247:GLY:HA2	1:B:250:GLN:HG3	2.02	0.42
3:A:270:DX8:CAH	3:A:270:DX8:CAE	2.95	0.41
1:C:254:GLY:HA3	1:D:265:LEU:HD11	2.02	0.41
1:B:64[B]:THR:HG22	1:B:122:GLU:HG2	2.03	0.41
1:B:35:HIS:HB2	2:B:269:NAP:C2A	2.51	0.41
1:C:65:ASN:HA	1:C:69:LEU:HD22	2.03	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:SER:HB3	3:A:270:DX8:HNAC	1.86	0.41
1:D:9:THR:O	1:D:93:ASN:HB3	2.21	0.41
1:D:220:LYS:HE3	5:D:346:HOH:O	2.20	0.40
1:C:22:LYS:HG2	1:C:242:ILE:HG13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/288 (86%)	238 (96%)	9 (4%)	0	100	100
1	B	252/288 (88%)	245 (97%)	7 (3%)	0	100	100
1	C	246/288 (85%)	238 (97%)	8 (3%)	0	100	100
1	D	250/288 (87%)	242 (97%)	8 (3%)	0	100	100
All	All	995/1152 (86%)	963 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/231 (87%)	202 (100%)	0	100	100
1	B	207/231 (90%)	205 (99%)	2 (1%)	82	85

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	200/231 (87%)	197 (98%)	3 (2%)	72	75
1	D	204/231 (88%)	200 (98%)	4 (2%)	63	65
All	All	813/924 (88%)	804 (99%)	9 (1%)	80	83

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	229	ARG
1	B	230	ARG
1	C	164	VAL
1	C	216	GLU
1	C	250	GLN
1	D	47	GLU
1	D	53	SER
1	D	206	VAL
1	D	229	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	54	ASN
1	A	236	GLN
1	B	179	HIS
1	B	236	GLN
1	D	67[A]	ASN
1	D	166	GLN
1	D	179	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAP	A	269	-	42,52,52	1.61	3 (7%)	54,80,80	2.08	8 (14%)
3	DX8	A	270	-	22,23,23	2.09	3 (13%)	19,33,33	2.40	3 (15%)
4	ACT	A	271	-	1,3,3	0.99	0	0,3,3	0.00	-
2	NAP	B	269	-	42,52,52	1.55	3 (7%)	54,80,80	2.22	9 (16%)
3	DX8	B	270	-	22,23,23	2.16	3 (13%)	19,33,33	2.57	6 (31%)
2	NAP	C	269	-	42,52,52	1.41	3 (7%)	54,80,80	2.27	7 (12%)
3	DX8	C	270	-	22,23,23	2.11	3 (13%)	19,33,33	2.38	4 (21%)
2	NAP	D	269	-	42,52,52	1.48	3 (7%)	54,80,80	2.09	7 (12%)
3	DX8	D	270	-	22,23,23	2.25	3 (13%)	19,33,33	2.37	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	269	-	-	0/27/67/67	0/5/5/5
3	DX8	A	270	-	-	0/6/8/8	0/3/3/3
4	ACT	A	271	-	-	0/0/0/0	0/0/0/0
2	NAP	B	269	-	-	0/27/67/67	0/5/5/5
3	DX8	B	270	-	-	0/6/8/8	0/3/3/3
2	NAP	C	269	-	-	0/27/67/67	0/5/5/5
3	DX8	C	270	-	-	0/6/8/8	0/3/3/3
2	NAP	D	269	-	-	0/27/67/67	0/5/5/5
3	DX8	D	270	-	-	0/6/8/8	0/3/3/3

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	270	DX8	CAP-CAR	-8.26	1.40	1.49
3	B	270	DX8	CAP-CAR	-7.97	1.40	1.49
3	A	270	DX8	CAP-CAR	-7.89	1.40	1.49
3	C	270	DX8	CAP-CAR	-7.84	1.40	1.49
2	C	269	NAP	C2A-N1A	2.58	1.38	1.33
2	A	269	NAP	C2A-N1A	2.62	1.38	1.33
2	B	269	NAP	C2A-N1A	2.76	1.39	1.33
2	D	269	NAP	C2A-N1A	2.89	1.39	1.33
3	A	270	DX8	CAN-NAK	3.04	1.40	1.35
3	B	270	DX8	CAN-NAK	3.22	1.41	1.35
2	C	269	NAP	C2A-N3A	3.42	1.38	1.32
3	C	270	DX8	CAN-NAK	3.43	1.41	1.35
2	D	269	NAP	C2A-N3A	3.55	1.38	1.32
3	A	270	DX8	CAS-NAK	3.77	1.40	1.33
3	D	270	DX8	CAN-NAK	3.80	1.42	1.35
2	A	269	NAP	C2A-N3A	3.85	1.39	1.32
2	B	269	NAP	C2A-N3A	3.94	1.39	1.32
3	C	270	DX8	CAS-NAK	4.05	1.40	1.33
3	B	270	DX8	CAS-NAK	4.17	1.40	1.33
3	D	270	DX8	CAS-NAK	4.55	1.41	1.33
2	C	269	NAP	O7N-C7N	6.50	1.38	1.24
2	D	269	NAP	O7N-C7N	6.92	1.38	1.24
2	B	269	NAP	O7N-C7N	7.45	1.40	1.24
2	A	269	NAP	O7N-C7N	7.93	1.41	1.24

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	269	NAP	N3A-C2A-N1A	-12.44	119.37	128.89
2	D	269	NAP	N3A-C2A-N1A	-11.72	119.92	128.89
2	A	269	NAP	N3A-C2A-N1A	-11.65	119.97	128.89
2	B	269	NAP	N3A-C2A-N1A	-10.64	120.75	128.89
3	B	270	DX8	CAU-CAS-NAK	-8.56	117.71	124.19
3	D	270	DX8	CAU-CAS-NAK	-7.88	118.22	124.19
3	A	270	DX8	CAU-CAS-NAK	-7.69	118.36	124.19
3	C	270	DX8	CAU-CAS-NAK	-7.63	118.41	124.19
2	B	269	NAP	O7N-C7N-C3N	-5.26	113.85	119.59
3	C	270	DX8	NAJ-CAN-NAK	-4.99	119.84	127.44
3	D	270	DX8	NAJ-CAN-NAK	-4.80	120.14	127.44
3	A	270	DX8	NAJ-CAN-NAK	-4.76	120.20	127.44
3	B	270	DX8	NAJ-CAN-NAK	-4.41	120.73	127.44
2	C	269	NAP	O7N-C7N-C3N	-4.03	115.18	119.59

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	269	NAP	C4B-O4B-C1B	-3.61	105.75	109.72
2	D	269	NAP	O7N-C7N-C3N	-3.57	115.69	119.59
2	A	269	NAP	C1B-N9A-C4A	-3.52	121.63	126.94
2	C	269	NAP	C1B-N9A-C4A	-3.16	122.17	126.94
2	D	269	NAP	C4B-O4B-C1B	-3.16	106.25	109.72
2	C	269	NAP	C4B-O4B-C1B	-3.14	106.27	109.72
2	B	269	NAP	C4B-O4B-C1B	-3.11	106.30	109.72
2	D	269	NAP	C1B-N9A-C4A	-2.76	122.77	126.94
2	B	269	NAP	C1B-N9A-C4A	-2.63	122.97	126.94
2	A	269	NAP	C4A-C5A-N7A	-2.43	107.24	109.48
3	B	270	DX8	CAI-CAP-CAR	-2.41	116.85	120.60
2	B	269	NAP	C4A-C5A-N7A	-2.29	107.37	109.48
2	B	269	NAP	O3D-C3D-C4D	-2.05	104.89	111.05
3	C	270	DX8	CAI-CAP-CAR	-2.05	117.41	120.60
3	B	270	DX8	CAR-CAQ-CAE	2.04	127.95	125.97
2	A	269	NAP	O2A-PA-O3	2.10	114.63	105.09
2	A	269	NAP	C3N-C7N-N7N	2.36	120.40	117.82
3	B	270	DX8	CAH-CAP-CAR	2.37	124.30	120.60
2	D	269	NAP	O4D-C1D-N1N	2.62	111.01	108.13
3	D	270	DX8	CAS-NAK-CAN	2.84	119.89	115.94
3	A	270	DX8	CAS-NAK-CAN	2.92	119.99	115.94
3	B	270	DX8	CAS-NAK-CAN	2.96	120.05	115.94
2	C	269	NAP	O4B-C1B-N9A	2.96	114.30	108.10
3	C	270	DX8	CAS-NAK-CAN	3.01	120.12	115.94
2	C	269	NAP	O4D-C1D-N1N	3.50	111.97	108.13
2	D	269	NAP	C3N-C7N-N7N	3.75	121.92	117.82
2	D	269	NAP	O4B-C1B-N9A	3.77	115.99	108.10
2	A	269	NAP	O4D-C1D-N1N	3.83	112.33	108.13
2	A	269	NAP	O4B-C1B-N9A	4.06	116.60	108.10
2	B	269	NAP	O4D-C1D-N1N	4.30	112.86	108.13
2	B	269	NAP	O4B-C1B-N9A	5.04	118.64	108.10
2	B	269	NAP	C3N-C7N-N7N	5.08	123.38	117.82
2	C	269	NAP	C3N-C7N-N7N	5.76	124.12	117.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	270	DX8	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	269	NAP	1	0
3	B	270	DX8	2	0
3	C	270	DX8	5	0
3	D	270	DX8	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	249/288 (86%)	-0.19	4 (1%) 74 75	9, 15, 22, 34	3 (1%)
1	B	248/288 (86%)	-0.27	4 (1%) 74 75	9, 14, 24, 35	0
1	C	249/288 (86%)	-0.12	7 (2%) 56 57	8, 14, 25, 48	1 (0%)
1	D	249/288 (86%)	-0.16	1 (0%) 93 93	9, 14, 24, 37	1 (0%)
All	All	995/1152 (86%)	-0.19	16 (1%) 74 75	8, 14, 24, 48	5 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	113	GLY	6.1
1	B	212	ALA	5.0
1	C	212	ALA	4.6
1	C	216	GLU	4.2
1	B	113	GLY	3.9
1	B	211	VAL	3.8
1	C	195	TYR	3.3
1	A	113	GLY	2.7
1	D	2	GLU	2.5
1	C	2	GLU	2.5
1	C	143	LYS	2.3
1	A	50	LYS	2.2
1	C	217[A]	GLU	2.1
1	B	2[A]	GLU	2.1
1	A	42	VAL	2.1
1	A	46[A]	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	DX8	D	270	21/21	0.84	0.29	17.74	24,29,33,36	0
3	DX8	A	270	21/21	0.89	0.26	13.88	22,27,34,36	0
3	DX8	C	270	21/21	0.87	0.29	9.26	21,29,40,41	0
3	DX8	B	270	21/21	0.89	0.21	7.91	17,25,33,35	0
4	ACT	A	271	4/4	0.98	0.11	-0.50	16,16,17,18	0
2	NAP	C	269	48/48	0.97	0.08	-1.32	10,14,21,22	0
2	NAP	A	269	48/48	0.97	0.07	-1.34	11,14,18,19	0
2	NAP	D	269	48/48	0.98	0.07	-1.89	8,14,19,21	0
2	NAP	B	269	48/48	0.98	0.06	-2.06	10,15,20,21	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.