



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2016 – 05:04 AM EDT

PDB ID : 5JQ2
Title : Crystal structure of the Ru(bpy)2PhenA functionalized P450 BM3 L407C heme domain mutant in complex with N-palmitoylglycine
Authors : Kloos, M.
Deposited on : 2016-05-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	<div> <div>2%</div> <div> <div></div> <div>94%</div> <div>...</div> </div> </div>
1	B	469	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>5% •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	140	A	502	-	-	-	X
3	140	B	502	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called P450 BM3 L407C heme domain mutant.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	457	Total	C	N	O	S	0	1	0
			3687	2352	629	688	18			
1	B	459	Total	C	N	O	S	0	0	0
			3691	2357	627	689	18			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	407	CYS	LEU	engineered mutation	UNP P14779
A	464	HIS	-	expression tag	UNP P14779
A	465	HIS	-	expression tag	UNP P14779
A	466	HIS	-	expression tag	UNP P14779
A	467	HIS	-	expression tag	UNP P14779
A	468	HIS	-	expression tag	UNP P14779
A	469	HIS	-	expression tag	UNP P14779
B	407	CYS	LEU	engineered mutation	UNP P14779
B	464	HIS	-	expression tag	UNP P14779
B	465	HIS	-	expression tag	UNP P14779
B	466	HIS	-	expression tag	UNP P14779
B	467	HIS	-	expression tag	UNP P14779
B	468	HIS	-	expression tag	UNP P14779
B	469	HIS	-	expression tag	UNP P14779

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is N-PALMITOYLGLYCINE (three-letter code: 140) (formula: C₁₈H₃₅NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			22	18	1	3		
3	B	1	Total	C	N	O	0	0
			22	18	1	3		

- Molecule 4 is bis(2,2'-bipyridine-kappa 2 N 1 ,N 1') [2-iodo-N-(1,10-phenanthrolin-5-yl-kappa 2 N 1 ,N 10)acetamide]ruthenium(2+) (three-letter code: RU8) (formula: C₃₄H₂₆IN₇ORu).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	Ru	0	0
			43	34	7	1	1		
4	B	1	Total	C	N	O	Ru	0	0
			43	34	7	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	248	Total	O	0	0
			248	248		
5	B	243	Total	O	0	0
			243	243		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: P450 BM3 L407C heme domain mutant



- Molecule 1: P450 BM3 L407C heme domain mutant



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.88Å 112.54Å 156.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.54 – 2.00 47.54 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (47.54-2.00) 99.2 (47.54-1.90)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 1.90Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.174 , 0.218 0.183 , 0.226	Depositor DCC
R_{free} test set	3581 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8085	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, RU8, 140

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/3772	0.87	8/5098 (0.2%)
1	B	0.73	0/3777	0.87	6/5107 (0.1%)
All	All	0.73	0/7549	0.87	14/10205 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	68	ASP	CB-CG-OD2	-7.21	111.81	118.30
1	A	79	ARG	NE-CZ-NH1	6.64	123.62	120.30
1	A	161	ARG	NE-CZ-NH2	-6.64	116.98	120.30
1	B	296	ARG	NE-CZ-NH2	-6.50	117.05	120.30
1	A	148	LEU	CA-CB-CG	6.24	129.66	115.30
1	A	50[A]	ARG	NE-CZ-NH2	-5.97	117.31	120.30
1	A	50[B]	ARG	NE-CZ-NH2	-5.97	117.31	120.30
1	A	161	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	B	147	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	79	ARG	NE-CZ-NH2	-5.33	117.64	120.30
1	A	63	ASP	CB-CG-OD1	5.24	123.02	118.30
1	B	68	ASP	CB-CG-OD1	5.12	122.91	118.30
1	B	148	LEU	CA-CB-CG	5.08	127.00	115.30
1	B	338	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	136	ASP	Peptide
1	A	229	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3687	0	3656	6	0
1	B	3691	0	3662	11	0
2	A	43	0	30	4	0
2	B	43	0	30	0	0
3	A	22	0	34	0	0
3	B	22	0	34	0	0
4	A	43	0	24	0	0
4	B	43	0	24	2	0
5	A	248	0	0	2	0
5	B	243	0	0	1	0
All	All	8085	0	7494	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50[B]:ARG:NH1	5:A:601:HOH:O	2.07	0.78
1:A:136:ASP:HB2	5:B:706:HOH:O	1.92	0.69
2:A:501:HEM:HMB2	2:A:501:HEM:HBB2	1.84	0.59
1:A:148:LEU:HD11	1:A:413:VAL:HG21	1.85	0.59
1:B:382:PRO:HG2	4:B:503:RU8:H32	1.88	0.56
1:A:349:LYS:NZ	5:A:602:HOH:O	2.35	0.52
1:B:8:PRO:HB2	1:B:19:LEU:HD21	1.91	0.52
1:B:382:PRO:HB2	4:B:503:RU8:H31	1.92	0.52
2:A:501:HEM:HMC2	2:A:501:HEM:HBC2	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:501:HEM:CMB	2:A:501:HEM:HBB2	2.45	0.46
1:B:17:LEU:HD11	1:B:188:LEU:O	2.15	0.46
1:A:271:GLY:HA2	1:A:440:LYS:HG3	1.98	0.45
1:B:222:ASP:O	1:B:226:SER:HB3	2.16	0.45
1:B:290:ALA:HB1	1:B:418:LEU:HD13	2.01	0.42
1:B:122:ILE:HG22	1:B:148:LEU:HD22	2.01	0.42
2:A:501:HEM:HBC2	2:A:501:HEM:CMC	2.50	0.41
1:B:104:LEU:N	1:B:105:PRO:CD	2.84	0.41
1:B:436:THR:O	1:B:437:LEU:HB2	2.21	0.41
1:A:1:THR:O	1:A:1:THR:HG23	2.22	0.40
1:B:280:LEU:HD22	1:B:287:LEU:HA	2.02	0.40
1:B:253:ASN:O	1:B:257:GLN:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/469 (97%)	442 (97%)	14 (3%)	0	100	100
1	B	457/469 (97%)	436 (95%)	20 (4%)	1 (0%)	52	48
All	All	913/938 (97%)	878 (96%)	34 (4%)	1 (0%)	56	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	196	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	400/411 (97%)	396 (99%)	4 (1%)	82	85
1	B	401/411 (98%)	400 (100%)	1 (0%)	95	97
All	All	801/822 (97%)	796 (99%)	5 (1%)	90	93

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	ASP
1	A	215	LEU
1	A	266	HIS
1	A	370	ASP
1	B	369	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	GLN
1	A	236	HIS
1	A	239	ASN
1	B	73	GLN
1	B	110	GLN
1	B	229	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	501	1	24,50,50	1.01	2 (8%)	16,82,82	1.50	3 (18%)
3	140	A	502	-	18,21,21	0.38	0	18,22,22	1.11	1 (5%)
4	RU8	A	503	1	45,52,53	4.17	30 (66%)	51,87,88	4.52	32 (62%)
2	HEM	B	501	1	24,50,50	0.82	0	16,82,82	1.65	5 (31%)
3	140	B	502	-	18,21,21	0.40	0	18,22,22	0.87	1 (5%)
4	RU8	B	503	1	45,52,53	4.11	25 (55%)	51,87,88	4.49	29 (56%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	1	-	0/6/54/54	0/0/8/8
3	140	A	502	-	-	0/18/20/20	0/0/0/0
4	RU8	A	503	1	-	0/4/116/118	0/10/10/10
2	HEM	B	501	1	-	0/6/54/54	0/0/8/8
3	140	B	502	-	-	0/18/20/20	0/0/0/0
4	RU8	B	503	1	-	0/4/116/118	0/10/10/10

All (57) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	503	RU8	C32-N33	-10.73	1.32	1.49
4	A	503	RU8	C23-N22	-10.36	1.33	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	RU8	C32-N33	-10.34	1.33	1.49
4	A	503	RU8	C45-N41	-10.25	1.33	1.49
4	B	503	RU8	C35-N34	-9.95	1.34	1.49
4	B	503	RU8	C45-N41	-9.32	1.34	1.49
4	A	503	RU8	C35-N34	-8.29	1.36	1.49
4	B	503	RU8	C23-N22	-7.82	1.37	1.49
4	B	503	RU8	C42-C40	-6.05	1.39	1.53
4	B	503	RU8	C29-C28	-5.75	1.40	1.53
4	B	503	RU8	C38-C39	-5.75	1.40	1.53
4	B	503	RU8	C30-C29	-5.65	1.38	1.53
4	A	503	RU8	C30-C29	-5.58	1.38	1.53
4	A	503	RU8	C26-C27	-5.52	1.40	1.53
4	B	503	RU8	C26-C27	-5.38	1.41	1.53
4	A	503	RU8	C29-C28	-5.33	1.41	1.53
4	B	503	RU8	C37-C38	-5.30	1.38	1.53
4	A	503	RU8	C42-C40	-5.30	1.41	1.53
4	A	503	RU8	C25-C26	-5.10	1.39	1.53
4	B	503	RU8	C43-C42	-5.02	1.39	1.53
4	A	503	RU8	C43-C42	-4.94	1.39	1.53
4	A	503	RU8	C37-C38	-4.81	1.40	1.53
4	A	503	RU8	C38-C39	-4.76	1.42	1.53
4	B	503	RU8	C25-C26	-4.70	1.40	1.53
4	B	503	RU8	C30-C31	-3.48	1.37	1.51
4	A	503	RU8	C30-C31	-3.22	1.38	1.51
4	B	503	RU8	C45-C44	-3.15	1.39	1.51
4	A	503	RU8	C47-N46	-3.09	1.30	1.35
4	A	503	RU8	C23-C24	-3.08	1.39	1.51
4	B	503	RU8	C32-C31	-3.08	1.39	1.51
4	B	503	RU8	C43-C44	-3.07	1.39	1.51
4	B	503	RU8	C23-C24	-3.07	1.39	1.51
4	B	503	RU8	C37-C36	-3.04	1.39	1.51
4	A	503	RU8	C43-C44	-3.03	1.39	1.51
4	A	503	RU8	C45-C44	-2.97	1.40	1.51
4	A	503	RU8	C32-C31	-2.92	1.40	1.51
4	A	503	RU8	C24-C25	-2.91	1.40	1.51
4	B	503	RU8	C24-C25	-2.79	1.40	1.51
4	A	503	RU8	C35-C36	-2.72	1.41	1.51
4	A	503	RU8	C37-C36	-2.63	1.41	1.51
4	A	503	RU8	RU1-N33	-2.48	2.02	2.12
2	A	501	HEM	C4C-NC	-2.40	1.33	1.36
4	B	503	RU8	C35-C36	-2.29	1.42	1.51
4	A	503	RU8	O48-C47	-2.05	1.18	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	503	RU8	RU1-N22	-2.01	2.04	2.12
4	B	503	RU8	C14-C11	2.05	1.48	1.43
4	A	503	RU8	C06-C01	2.31	1.41	1.36
4	A	503	RU8	C14-C11	2.32	1.48	1.43
2	A	501	HEM	CMA-C3A	2.37	1.56	1.51
4	A	503	RU8	C21-C14	3.07	1.46	1.41
4	A	503	RU8	C11-N10	3.34	1.42	1.37
4	A	503	RU8	C05-C21	3.43	1.49	1.43
4	B	503	RU8	C05-C21	3.43	1.49	1.43
4	B	503	RU8	C02-C11	3.74	1.47	1.41
4	B	503	RU8	C21-C14	3.77	1.47	1.41
4	B	503	RU8	C11-N10	4.03	1.43	1.37
4	A	503	RU8	C02-C11	5.12	1.50	1.41

All (71) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	RU8	C21-C05-N46	-3.38	112.10	118.42
2	A	501	HEM	CBD-CAD-C3D	-3.04	107.14	112.47
4	A	503	RU8	C21-C14-N13	-3.02	119.54	122.76
2	B	501	HEM	C3C-C4C-NC	-2.78	105.70	110.94
4	A	503	RU8	O48-C47-C49	-2.62	117.25	122.07
3	B	502	140	O3-C3-C4	-2.52	117.60	121.97
2	B	501	HEM	C3B-CAB-CBB	-2.49	121.39	126.40
2	B	501	HEM	CBD-CAD-C3D	-2.48	108.11	112.47
4	B	503	RU8	C06-C08-N10	-2.44	118.99	122.70
2	A	501	HEM	C3C-C4C-NC	-2.24	106.72	110.94
2	B	501	HEM	C3B-C4B-NB	-2.15	106.43	109.21
4	A	503	RU8	C01-C02-C11	-2.00	114.69	117.93
4	A	503	RU8	C19-C21-C05	2.02	126.30	122.95
4	B	503	RU8	C49-C47-N46	2.06	117.98	115.01
2	B	501	HEM	CAD-CBD-CGD	2.06	116.80	112.78
2	A	501	HEM	CMC-C2C-C3C	2.21	129.41	125.09
4	B	503	RU8	C15-N13-C14	2.27	120.47	117.54
3	A	502	140	C4-C3-N	2.29	120.44	116.46
4	B	503	RU8	C14-C11-N10	2.31	119.34	116.56
4	A	503	RU8	C01-C02-C03	2.50	126.38	122.05
4	B	503	RU8	C08-C06-C01	2.76	123.24	119.40
4	B	503	RU8	C36-C35-N34	3.29	118.44	112.97
4	A	503	RU8	C49-C47-N46	3.32	119.81	115.01
4	A	503	RU8	C44-C43-C42	3.37	118.41	111.44
4	B	503	RU8	C44-C43-C42	3.38	118.42	111.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	RU8	C31-C30-C29	3.59	118.87	111.44
4	A	503	RU8	C40-C39-N34	3.60	114.48	109.12
4	A	503	RU8	C31-C30-C29	3.66	119.02	111.44
4	B	503	RU8	C44-C45-N41	3.75	119.20	112.97
4	B	503	RU8	C36-C37-C38	3.75	119.19	111.44
4	A	503	RU8	C43-C42-C40	3.94	118.69	111.20
4	A	503	RU8	C24-C25-C26	3.94	119.60	111.44
4	A	503	RU8	C15-N13-C14	3.96	122.64	117.54
4	A	503	RU8	C24-C23-N22	3.98	119.58	112.97
4	B	503	RU8	C31-C32-N33	3.99	119.61	112.97
4	A	503	RU8	C36-C35-N34	4.03	119.67	112.97
4	B	503	RU8	C24-C23-N22	4.06	119.72	112.97
4	A	503	RU8	C44-C45-N41	4.11	119.80	112.97
4	A	503	RU8	C37-C38-C39	4.14	119.07	111.20
4	A	503	RU8	C30-C29-C28	4.15	119.08	111.20
4	B	503	RU8	C24-C25-C26	4.15	120.02	111.44
4	A	503	RU8	C36-C37-C38	4.26	120.25	111.44
4	A	503	RU8	C25-C26-C27	4.28	119.34	111.20
4	B	503	RU8	C30-C29-C28	4.31	119.40	111.20
4	B	503	RU8	C37-C38-C39	4.38	119.53	111.20
4	B	503	RU8	C43-C42-C40	4.38	119.54	111.20
4	A	503	RU8	C31-C32-N33	4.38	120.26	112.97
4	B	503	RU8	C25-C26-C27	4.41	119.58	111.20
4	B	503	RU8	C28-C27-N22	4.69	116.11	109.12
4	B	503	RU8	C23-C24-C25	4.76	118.81	111.43
4	A	503	RU8	C23-C24-C25	4.98	119.16	111.43
4	B	503	RU8	C40-C39-N34	5.10	116.71	109.12
4	A	503	RU8	C32-C31-C30	5.41	119.83	111.43
4	B	503	RU8	C39-C40-N41	5.63	117.51	109.12
4	A	503	RU8	C28-C27-N22	5.64	117.52	109.12
4	A	503	RU8	C35-C36-C37	5.69	120.26	111.43
4	A	503	RU8	C39-C40-N41	5.86	117.85	109.12
4	B	503	RU8	C45-C44-C43	5.93	120.63	111.43
4	B	503	RU8	C32-C31-C30	6.04	120.80	111.43
4	A	503	RU8	C45-C44-C43	6.29	121.19	111.43
4	A	503	RU8	C27-C28-N33	6.33	118.55	109.12
4	B	503	RU8	C35-C36-C37	6.39	121.35	111.43
4	B	503	RU8	C27-C28-N33	6.94	119.46	109.12
4	A	503	RU8	C45-N41-C40	10.20	120.72	109.55
4	A	503	RU8	C32-N33-C28	10.76	121.33	109.55
4	B	503	RU8	C32-N33-C28	10.96	121.55	109.55
4	A	503	RU8	C35-N34-C39	10.97	121.56	109.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	RU8	C35-N34-C39	11.04	121.64	109.55
4	B	503	RU8	C45-N41-C40	11.07	121.67	109.55
4	B	503	RU8	C23-N22-C27	12.35	123.07	109.55
4	A	503	RU8	C23-N22-C27	12.83	123.59	109.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	4	0
4	B	503	RU8	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	457/469 (97%)	-0.42	8 (1%) 71 72	16, 25, 48, 94	0
1	B	459/469 (97%)	-0.37	9 (1%) 68 69	17, 26, 54, 103	0
All	All	916/938 (97%)	-0.40	17 (1%) 70 70	16, 26, 51, 103	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	227	GLY	7.5
1	B	228	GLU	5.4
1	A	229	GLN	4.5
1	A	1	THR	3.5
1	B	196	PRO	3.4
1	A	2	ILE	3.3
1	A	457	GLY	3.2
1	A	228	GLU	3.1
1	A	136	ASP	2.8
1	B	229	GLN	2.8
1	B	192	ASN	2.8
1	B	225	ALA	2.7
1	B	1	THR	2.6
1	B	457	GLY	2.5
1	A	227	GLY	2.4
1	B	203	ARG	2.4
1	A	203	ARG	2.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	140	A	502	22/22	0.91	0.14	3.41	29,33,38,39	0
3	140	B	502	22/22	0.91	0.14	2.68	22,27,37,43	0
4	RU8	B	503	43/44	0.97	0.14	1.26	24,27,29,31	43
2	HEM	B	501	43/43	0.97	0.10	0.72	15,18,19,22	0
2	HEM	A	501	43/43	0.98	0.10	0.18	15,17,19,26	0
4	RU8	A	503	43/44	0.98	0.11	-0.08	23,30,35,43	43

6.5 Other polymers [i](#)

There are no such residues in this entry.