



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2016 – 07:12 PM EDT

PDB ID : 5JT2
Title : BRAFV600E Kinase Domain In Complex with Chemically Linked Vemurafenib Inhibitor VEM-BISAMIDE
Authors : Grasso, M.J.; Marmorstein, R.
Deposited on : 2016-05-09
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	<div> <div>2%</div> <div> <div></div> <div>71%</div> <div>20%</div> <div>•</div> <div>8%</div> </div> </div>
1	B	280	<div> <div>3%</div> <div> <div></div> <div>67%</div> <div>21%</div> <div>•</div> <div>10%</div> </div> </div>
1	C	280	<div> <div>6%</div> <div> <div></div> <div>67%</div> <div>22%</div> <div>•</div> <div>10%</div> </div> </div>
1	D	280	<div> <div>4%</div> <div> <div></div> <div>65%</div> <div>18%</div> <div>•</div> <div>16%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8051 atoms, of which 104 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase B-raf.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			1958	1249	339	358	12			
1	B	251	Total	C	N	O	S	0	0	0
			1921	1225	331	353	12			
1	C	251	Total	C	N	O	S	0	0	0
			1925	1231	333	349	12			
1	D	234	Total	C	N	O	S	0	0	0
			1801	1152	309	329	11			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	444	GLY	-	expression tag	UNP P15056
A	445	SER	-	expression tag	UNP P15056
A	446	GLU	-	expression tag	UNP P15056
A	447	PHE	-	expression tag	UNP P15056
A	543	ALA	ILE	engineered mutation	UNP P15056
A	544	SER	ILE	engineered mutation	UNP P15056
A	551	LYS	ILE	engineered mutation	UNP P15056
A	562	ARG	GLN	engineered mutation	UNP P15056
A	588	ASN	LEU	engineered mutation	UNP P15056
A	600	GLU	VAL	engineered mutation	UNP P15056
A	630	SER	LYS	engineered mutation	UNP P15056
A	667	GLU	PHE	engineered mutation	UNP P15056
A	673	SER	TYR	engineered mutation	UNP P15056
A	688	ARG	ALA	engineered mutation	UNP P15056
A	706	SER	LEU	engineered mutation	UNP P15056
A	709	ARG	GLN	engineered mutation	UNP P15056
A	713	GLU	SER	engineered mutation	UNP P15056
A	716	GLU	LEU	engineered mutation	UNP P15056
A	720	GLU	SER	engineered mutation	UNP P15056
A	722	SER	PRO	engineered mutation	UNP P15056
A	723	GLY	LYS	engineered mutation	UNP P15056

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	444	GLY	-	expression tag	UNP P15056
B	445	SER	-	expression tag	UNP P15056
B	446	GLU	-	expression tag	UNP P15056
B	447	PHE	-	expression tag	UNP P15056
B	543	ALA	ILE	engineered mutation	UNP P15056
B	544	SER	ILE	engineered mutation	UNP P15056
B	551	LYS	ILE	engineered mutation	UNP P15056
B	562	ARG	GLN	engineered mutation	UNP P15056
B	588	ASN	LEU	engineered mutation	UNP P15056
B	600	GLU	VAL	engineered mutation	UNP P15056
B	630	SER	LYS	engineered mutation	UNP P15056
B	667	GLU	PHE	engineered mutation	UNP P15056
B	673	SER	TYR	engineered mutation	UNP P15056
B	688	ARG	ALA	engineered mutation	UNP P15056
B	706	SER	LEU	engineered mutation	UNP P15056
B	709	ARG	GLN	engineered mutation	UNP P15056
B	713	GLU	SER	engineered mutation	UNP P15056
B	716	GLU	LEU	engineered mutation	UNP P15056
B	720	GLU	SER	engineered mutation	UNP P15056
B	722	SER	PRO	engineered mutation	UNP P15056
B	723	GLY	LYS	engineered mutation	UNP P15056
C	444	GLY	-	expression tag	UNP P15056
C	445	SER	-	expression tag	UNP P15056
C	446	GLU	-	expression tag	UNP P15056
C	447	PHE	-	expression tag	UNP P15056
C	543	ALA	ILE	engineered mutation	UNP P15056
C	544	SER	ILE	engineered mutation	UNP P15056
C	551	LYS	ILE	engineered mutation	UNP P15056
C	562	ARG	GLN	engineered mutation	UNP P15056
C	588	ASN	LEU	engineered mutation	UNP P15056
C	600	GLU	VAL	engineered mutation	UNP P15056
C	630	SER	LYS	engineered mutation	UNP P15056
C	667	GLU	PHE	engineered mutation	UNP P15056
C	673	SER	TYR	engineered mutation	UNP P15056
C	688	ARG	ALA	engineered mutation	UNP P15056
C	706	SER	LEU	engineered mutation	UNP P15056
C	709	ARG	GLN	engineered mutation	UNP P15056
C	713	GLU	SER	engineered mutation	UNP P15056
C	716	GLU	LEU	engineered mutation	UNP P15056
C	720	GLU	SER	engineered mutation	UNP P15056
C	722	SER	PRO	engineered mutation	UNP P15056
C	723	GLY	LYS	engineered mutation	UNP P15056

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	444	GLY	-	expression tag	UNP P15056
D	445	SER	-	expression tag	UNP P15056
D	446	GLU	-	expression tag	UNP P15056
D	447	PHE	-	expression tag	UNP P15056
D	543	ALA	ILE	engineered mutation	UNP P15056
D	544	SER	ILE	engineered mutation	UNP P15056
D	551	LYS	ILE	engineered mutation	UNP P15056
D	562	ARG	GLN	engineered mutation	UNP P15056
D	588	ASN	LEU	engineered mutation	UNP P15056
D	600	GLU	VAL	engineered mutation	UNP P15056
D	630	SER	LYS	engineered mutation	UNP P15056
D	667	GLU	PHE	engineered mutation	UNP P15056
D	673	SER	TYR	engineered mutation	UNP P15056
D	688	ARG	ALA	engineered mutation	UNP P15056
D	706	SER	LEU	engineered mutation	UNP P15056
D	709	ARG	GLN	engineered mutation	UNP P15056
D	713	GLU	SER	engineered mutation	UNP P15056
D	716	GLU	LEU	engineered mutation	UNP P15056
D	720	GLU	SER	engineered mutation	UNP P15056
D	722	SER	PRO	engineered mutation	UNP P15056
D	723	GLY	LYS	engineered mutation	UNP P15056

- Molecule 2 is BENZAMIDINE (three-letter code: BEN) (formula: C₇H₈N₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	N	0	0
			17	7	8	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	H	N	0	0
			17	7	8	2		

- Molecule 3 is 2,2'-oxybis[N-({4-[3-(3-{[dihydroxy(propyl)-lambda 4 -sulfanyl]amino}-2,6-difluorobenzene-1-carbonyl)-1H-pyrrolo[2,3-b]pyridin-5-yl]phenyl}methyl)acetamide] (three-letter code: 6NC) (formula: C₅₂H₅₀F₄N₈O₉S₂).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	F	H	N	O	S	0	0
			119	52	4	44	8	9	2		
3	C	1	Total	C	F	H	N	O	S	0	0
			119	52	4	44	8	9	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	49	Total	O	0	0
			49	49		
4	B	63	Total	O	0	0
			63	63		
4	C	26	Total	O	0	0
			26	26		
4	D	36	Total	O	0	0
			36	36		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase B-raf




- Molecule 1: Serine/threonine-protein kinase B-raf




- Molecule 1: Serine/threonine-protein kinase B-raf





● Molecule 1: Serine/threonine-protein kinase B-raf



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.10Å 68.44Å 275.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.59 – 2.70 29.58 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.59-2.70) 96.1 (29.58-2.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 2.72Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.212 , 0.273 0.207 , 0.267	Depositor DCC
R_{free} test set	1929 reflections (5.78%)	DCC
Wilson B-factor (Å ²)	54.9	Xtriage
Anisotropy	0.223	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.033 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8051	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BEN, 6NC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/1999	0.64	0/2705
1	B	0.48	0/1960	0.66	1/2652 (0.0%)
1	C	0.44	0/1962	0.60	0/2647
1	D	0.44	0/1839	0.62	0/2489
All	All	0.46	0/7760	0.63	1/10493 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	449	ASP	CB-CA-C	-5.23	99.94	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1958	0	1864	43	0
1	B	1921	0	1828	50	0
1	C	1925	0	1846	54	0
1	D	1801	0	1705	39	0
2	A	9	8	7	0	0
2	B	9	8	7	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	75	44	0	1	0
3	C	75	44	0	2	0
4	A	49	0	0	8	0
4	B	63	0	0	9	0
4	C	26	0	0	2	0
4	D	36	0	0	5	0
All	All	7947	104	7257	177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:491:THR:HG23	1:C:494:GLN:HE21	1.29	0.95
1:C:665:ILE:HG22	1:C:666:ILE:H	1.39	0.87
1:A:668:MET:HB3	1:A:674:LEU:HB2	1.59	0.84
1:B:502:VAL:HG13	1:B:516:PHE:CE2	2.17	0.79
1:C:546:THR:HA	4:C:926:HOH:O	1.84	0.77

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/280 (89%)	240 (96%)	9 (4%)	0	100	100
1	B	243/280 (87%)	233 (96%)	8 (3%)	2 (1%)	24	51
1	C	237/280 (85%)	227 (96%)	9 (4%)	1 (0%)	39	69
1	D	222/280 (79%)	214 (96%)	7 (3%)	1 (0%)	34	63
All	All	951/1120 (85%)	914 (96%)	33 (4%)	4 (0%)	39	69

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	492	PRO
1	B	493	GLN
1	C	666	ILE
1	D	492	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/245 (80%)	191 (97%)	5 (3%)	54	83
1	B	195/245 (80%)	189 (97%)	6 (3%)	47	78
1	C	193/245 (79%)	191 (99%)	2 (1%)	82	94
1	D	181/245 (74%)	176 (97%)	5 (3%)	51	81
All	All	765/980 (78%)	747 (98%)	18 (2%)	57	85

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	636	GLN
1	B	664	GLN
1	D	454	ASP
1	B	576	ASP
1	B	617	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	524	GLN
1	C	494	GLN
1	C	539	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BEN	A	801	-	9,9,9	5.03	7 (77%)	9,11,11	0.68	0
3	6NC	A	802	-	72,82,82	1.74	13 (18%)	81,119,119	1.97	24 (29%)
2	BEN	B	801	-	9,9,9	4.83	7 (77%)	9,11,11	0.77	0
3	6NC	C	801	-	72,82,82	1.72	14 (19%)	81,119,119	2.08	22 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEN	A	801	-	-	0/4/4/4	0/1/1/1
3	6NC	A	802	-	-	0/42/58/58	0/8/8/8
2	BEN	B	801	-	-	0/4/4/4	0/1/1/1
3	6NC	C	801	-	-	0/42/58/58	0/8/8/8

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	BEN	C2-C1	-5.15	1.30	1.39
2	B	801	BEN	C2-C1	-4.64	1.31	1.39
2	A	801	BEN	C5-C6	-3.60	1.31	1.38
2	B	801	BEN	C5-C6	-3.43	1.32	1.38
3	C	801	6NC	C63-N62	-3.15	1.30	1.36

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	801	6NC	C54-C49-N46	-5.62	115.02	120.79
3	C	801	6NC	C24-C25-N26	-4.97	120.28	125.49
3	C	801	6NC	C13-C08-N05	-4.97	115.69	120.79
3	A	802	6NC	C54-C49-N46	-4.71	115.95	120.79
3	C	801	6NC	O37-C36-C35	-4.63	104.02	112.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	802	6NC	1	0
3	C	801	6NC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/280 (91%)	-0.10	5 (1%) 70 70	21, 35, 56, 69	0
1	B	251/280 (89%)	-0.13	7 (2%) 56 57	19, 35, 53, 71	0
1	C	251/280 (89%)	0.21	16 (6%) 23 21	21, 48, 69, 85	0
1	D	234/280 (83%)	0.09	10 (4%) 39 38	24, 44, 70, 84	0
All	All	993/1120 (88%)	0.01	38 (3%) 44 44	19, 39, 65, 85	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	660	ASN	6.0
1	C	656	TYR	4.5
1	A	604	TRP	4.3
1	C	661	ASN	3.9
1	C	671	ARG	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	BEN	A	801	9/9	0.97	0.18	1.15	21,26,30,32	0
3	6NC	C	801	75/75	0.93	0.18	-0.10	21,34,52,63	0
3	6NC	A	802	75/75	0.95	0.16	-0.33	18,28,45,49	0
2	BEN	B	801	9/9	0.97	0.12	-1.35	21,25,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.