



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 20, 2016 – 07:47 PM EDT

PDB ID : 5JXC
Title : SynGAP Coiled-coil trimer
Authors : Shang, Y.; Zhang, M.
Deposited on : 2016-05-13
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	92	<div> <div>3%</div> <div>72% 21% • 7%</div> </div>
1	B	92	<div> <div>%</div> <div>67% 17% 15%</div> </div>
1	C	92	<div> <div>3%</div> <div>64% 21% 15%</div> </div>
1	D	92	<div> <div>2%</div> <div>58% 23% • 15%</div> </div>
1	E	92	<div> <div>%</div> <div>70% 21% • 9%</div> </div>
1	F	92	<div> <div>%</div> <div>65% 21% • 13%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3840 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ras/Rap GTPase-activating protein SynGAP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	86	Total	C	N	O	S	0	0	0
			680	416	131	131	2			
1	B	78	Total	C	N	O	S	0	0	0
			600	372	110	116	2			
1	C	78	Total	C	N	O	S	0	0	0
			613	377	112	121	3			
1	D	78	Total	C	N	O	S	0	0	0
			628	387	112	126	3			
1	E	84	Total	C	N	O	S	0	0	0
			657	405	119	131	2			
1	F	80	Total	C	N	O	S	0	0	0
			626	386	112	125	3			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1183	GLY	-	expression tag	UNP A0A0J9YUM2
A	1184	PRO	-	expression tag	UNP A0A0J9YUM2
A	1185	GLY	-	expression tag	UNP A0A0J9YUM2
A	1186	SER	-	expression tag	UNP A0A0J9YUM2
B	1183	GLY	-	expression tag	UNP A0A0J9YUM2
B	1184	PRO	-	expression tag	UNP A0A0J9YUM2
B	1185	GLY	-	expression tag	UNP A0A0J9YUM2
B	1186	SER	-	expression tag	UNP A0A0J9YUM2
C	1183	GLY	-	expression tag	UNP A0A0J9YUM2
C	1184	PRO	-	expression tag	UNP A0A0J9YUM2
C	1185	GLY	-	expression tag	UNP A0A0J9YUM2
C	1186	SER	-	expression tag	UNP A0A0J9YUM2
D	1183	GLY	-	expression tag	UNP A0A0J9YUM2
D	1184	PRO	-	expression tag	UNP A0A0J9YUM2
D	1185	GLY	-	expression tag	UNP A0A0J9YUM2
D	1186	SER	-	expression tag	UNP A0A0J9YUM2
E	1183	GLY	-	expression tag	UNP A0A0J9YUM2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1184	PRO	-	expression tag	UNP A0A0J9YUM2
E	1185	GLY	-	expression tag	UNP A0A0J9YUM2
E	1186	SER	-	expression tag	UNP A0A0J9YUM2
F	1183	GLY	-	expression tag	UNP A0A0J9YUM2
F	1184	PRO	-	expression tag	UNP A0A0J9YUM2
F	1185	GLY	-	expression tag	UNP A0A0J9YUM2
F	1186	SER	-	expression tag	UNP A0A0J9YUM2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	4	Total O 4 4	0	0
2	B	7	Total O 7 7	0	0
2	C	4	Total O 4 4	0	0
2	D	5	Total O 5 5	0	0
2	E	11	Total O 11 11	0	0
2	F	5	Total O 5 5	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ras/Rap GTPase-activating protein SynGAP



- Molecule 1: Ras/Rap GTPase-activating protein SynGAP



- Molecule 1: Ras/Rap GTPase-activating protein SynGAP



- Molecule 1: Ras/Rap GTPase-activating protein SynGAP



- Molecule 1: Ras/Rap GTPase-activating protein SynGAP



- Molecule 1: Ras/Rap GTPase-activating protein SynGAP



GLY	R1193
PRO	L1202
GLY	K1203
SER	L1206
ASP	H1207
GLU	E1215
SER	Y1216
ARG	R1219
LEU	K1229
ASP	Q1233
	Y1234
	R1237
	L1238
	E1242
	L1245
	Q1249
	K1252
	D1253
	E1267
	L1270
	R1271
	R1272
	ASP
	HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	28.80 Å 236.66 Å 42.21 Å 90.00° 96.92° 90.00°	Depositor
Resolution (Å)	39.50 – 2.50 39.50 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.8 (39.50-2.50) 99.5 (39.50-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.98 (at 2.51 Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, R_{free}	0.223 , 0.307 0.220 , 0.304	Depositor DCC
R_{free} test set	979 reflections (5.11%)	DCC
Wilson B-factor (Å ²)	53.2	Xtriage
Anisotropy	0.508	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3840	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/684	0.53	0/915
1	B	0.43	0/603	0.55	0/808
1	C	0.41	0/616	0.58	0/825
1	D	0.42	0/632	0.57	0/846
1	E	0.39	0/661	0.55	0/886
1	F	0.44	0/629	0.54	0/842
All	All	0.41	0/3825	0.55	0/5122

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	680	0	651	23	0
1	B	600	0	575	16	0
1	C	613	0	585	19	1
1	D	628	0	600	26	0
1	E	657	0	613	14	1
1	F	626	0	595	22	0
2	A	4	0	0	3	0
2	B	7	0	0	0	0
2	C	4	0	0	0	0
2	D	5	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	11	0	0	0	0
2	F	5	0	0	2	0
All	All	3840	0	3619	84	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1256:ILE:O	1:A:1260:ILE:HD13	1.51	1.09
1:D:1230:ILE:HD11	1:E:1231:LEU:HD13	1.49	0.94
1:A:1256:ILE:O	1:A:1260:ILE:CD1	2.19	0.91
1:A:1202:LEU:HD21	1:B:1203:LYS:HG2	1.54	0.89
1:A:1268:GLU:OE2	1:A:1271:ARG:NH1	2.07	0.87

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1233:GLN:NE2	1:E:1198:GLU:OE1[2_447]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	84/92 (91%)	84 (100%)	0	0	100	100
1	B	76/92 (83%)	75 (99%)	1 (1%)	0	100	100
1	C	76/92 (83%)	76 (100%)	0	0	100	100
1	D	76/92 (83%)	76 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	82/92 (89%)	81 (99%)	1 (1%)	0	100	100
1	F	78/92 (85%)	78 (100%)	0	0	100	100
All	All	472/552 (86%)	470 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	69/88 (78%)	67 (97%)	2 (3%)	50	77
1	B	60/88 (68%)	57 (95%)	3 (5%)	30	53
1	C	63/88 (72%)	60 (95%)	3 (5%)	31	55
1	D	66/88 (75%)	61 (92%)	5 (8%)	16	30
1	E	65/88 (74%)	62 (95%)	3 (5%)	33	57
1	F	64/88 (73%)	61 (95%)	3 (5%)	32	56
All	All	387/528 (73%)	368 (95%)	19 (5%)	31	55

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	1205	ARG
1	D	1238	LEU
1	E	1258	SER
1	C	1258	SER
1	F	1206	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1210	ASN
1	C	1207	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	1255	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	86/92 (93%)	0.23	3 (3%) 48 53	45, 68, 88, 115	0
1	B	78/92 (84%)	0.17	1 (1%) 79 82	47, 65, 89, 95	0
1	C	78/92 (84%)	0.24	3 (3%) 44 49	47, 59, 83, 92	0
1	D	78/92 (84%)	0.21	2 (2%) 59 63	46, 57, 84, 94	0
1	E	84/92 (91%)	0.09	1 (1%) 81 83	47, 65, 84, 98	0
1	F	80/92 (86%)	0.29	1 (1%) 79 82	46, 68, 89, 117	0
All	All	484/552 (87%)	0.21	11 (2%) 64 67	45, 64, 88, 117	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1195	TYR	5.5
1	A	1206	LEU	2.9
1	C	1245	LEU	2.6
1	C	1234	TYR	2.5
1	A	1240	GLN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.